





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07D 417/04, A01N 43/80, C07D 487/04, A01N 43/824, C07D 513/04, 471/04, 275/06, 291/08 // (C07D 471/04, 235:00, 221:00) (C07D 487/04, 249:00, 237:00) (C07D 513/04, 285:00, 237:00)</p>	A1	<p>(11) International Publication Number: WO 95/33746</p> <p>(43) International Publication Date: 14 December 1995 (14.12.95)</p>
<p>(21) International Application Number: PCT/US95/06706</p> <p>(22) International Filing Date: 2 June 1995 (02.06.95)</p> <p>(30) Priority Data: 08/255,587 8 June 1994 (08.06.94) US</p> <p>(60) Parent Application or Grant (63) Related by Continuation US 08/255,587 (CIP) Filed on 8 June 1994 (08.06.94)</p> <p>(71) Applicant (for all designated States except US): E.I. DU PONT DE NEMOURS AND COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): KAMIREDDY, Balreddy [US/US]; 204 Golding Court, Hockessin, DE 19707-1356 (US). MURRAY, William, Mark [US/US]; 126 McMullen Circle, Bear, DE 19701-3059 (US).</p>	<p>(74) Agents: GREGORY, Theodore, C. et al.; E.I. du Pont de Nemours and Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).</p> <p>(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, IS, JP, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).</p> <p>Published With international search report.</p>	

(54) Title: CYCLIC SULFONAMIDE HERBICIDES



(57) Abstract

Compounds of formula (I) and (II), and their agriculturally-suitable salts, are disclosed which are useful for controlling undesired vegetation, wherein X is a direct bond, O, S, NH, N(C₁-C₃ alkyl), N(C₁-C₃ haloalkyl) or N(allyl); R¹ is H, F, Cl, or Br; R² is H, F, Cl, Br, CF₃, nitro, or cyano; R⁴ is H, C₁-C₃ alkyl, or halogen; R⁵ is H, C₁-C₆ alkyl, C₁-C₆ haloalkyl, halogen, S(O)₂(C₁-C₆ alkyl), or C(=O)R⁸; or R⁴ and R⁵ are taken together along with the carbon to which they are attached to form a spiro-cyclopropane ring; and R³, R⁸, and J are as defined in the disclosure. Also disclosed are compositions containing the compounds of formulae (I) and (II) and a method for controlling undesired vegetation which involves contacting the vegetation or its environment with an effective amount of a compound of formula (I) or (II).

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.


AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

TITLE
CYCLIC SULFONAMIDE HERBICIDES
BACKGROUND OF THE INVENTION

This invention relates to certain cyclic sulfonamides, their agriculturally-suitable salts and compositions, and methods of their use as general or selective preemergent or postemergent herbicides.

New compounds effective for controlling the growth of undesired vegetation are in constant demand. In the most common situation, such compounds are sought to selectively control the growth of weeds in useful crops such as cotton, rice, corn, wheat, citrus and soybeans, to name a few. Unchecked weed growth in such crops (including plantation crops) can cause significant losses, reducing profit to the farmer and increasing costs to the consumer. In other situations, herbicides are desired which will control all plant growth. Examples of areas in which complete control of all vegetation is desired are areas around railroad tracks, storage tanks and industrial storage areas. There are many products commercially available for these purposes, but the search continues for products which are more effective, less costly and environmentally safe.

U.S. 4,818,275 discloses herbicidal acyclic sulfonamides of the formula



20

wherein, *inter alia*


X and Y are Br; Cl; or F;

R is alkyl; haloalkyl; or dialkylamino; and

R¹ is H; Na; lower alkyl; or SO₂R.

25

JP 63[1988]-222167 discloses herbicidal cyclic amides and sulfonamides of the formula



wherein

X is H or halogen;

Z is O or S;


J is, *inter alia*, -CO- or -SO₂-;

L₁ is -CO- or -CR₄(R₅)-;

5 R₄ and R₅ are each independently, *inter alia*, H, C₁-C₆ alkyl, C₁-C₆ haloalkyl, or
CO₂R_a;

R₁ is, *inter alia*, H, C₂-C₆ alkoxyalkyl, C₁-C₆ alkyl, C₁-C₆ haloalkyl, C₂-C₆
alkenyl, C₂-C₆ haloalkenyl, C₃-C₈ cycloalkyl, -COR_b, -CO₂R_b, or
-CH₂CO₂R_b;

W is, *inter alia*,



(W2)

10

Q is N or CH;

E is O or S;

R₁₂ is H, C₁-C₆ alkyl, or C₁-C₆ haloalkyl when Q is N; or

15 R₁₂ is H, C₁-C₆ alkyl, C₁-C₆ haloalkyl, or C₄ alkylene which combines with Q to
form a 6-membered ring when Q is CH;

R₁₃ is H, C₁-C₆ alkyl, or C₁-C₆ haloalkyl when Q is N; or



R₁₃ is H, C₁-C₆ alkyl, C₁-C₆ haloalkyl, or C₄ alkylene which combines with Q to
form a 6-membered ring when Q is CH; and

m is 0 or 1.

20 The cyclic sulfonamides of the present invention are not disclosed in either of these
references.

SUMMARY OF THE INVENTION

25 This invention includes compounds of Formulae I and II including all geometric
and stereoisomers, agriculturally suitable salts thereof, agricultural compositions
containing them and their use for controlling undesirable vegetation:



wherein

X is a direct bond; O; S; NH; N(C₁-C₃ alkyl); N(C₁-C₃ haloalkyl); or N(allyl);

R¹ is H; F; Cl; or Br;

R² is H; F; Cl; Br; CF₃; nitro; or cyano;

5 R³ is H; C₁-C₈ alkyl; C₃-C₈ cycloalkyl; C₃-C₈ alkenyl; C₃-C₈ alkynyl; C₁-C₈ haloalkyl; C₂-C₈ alkoxyalkyl; C₃-C₈ alkoxyalkoxyalkyl; C₃-C₈ haloalkynyl; C₃-C₈ haloalkenyl; C₁-C₈ alkylsulfonyl; C₁-C₈ haloalkylsulfonyl; C₃-C₈ alkoxyalkoxyalkyl; S(O)₂NH(C₁-C₈ alkyl); C(O)R⁶; or benzyl optionally substituted on the phenyl ring with R⁷;

10 R⁴ is H; C₁-C₃ alkyl; or halogen;

R⁵ is H; C₁-C₆ alkyl; C₁-C₆ haloalkyl; halogen; S(O)₂(C₁-C₆ alkyl); or C(=O)R⁸;
or


R⁴ and R⁵ are taken together along with the carbon to which they are attached to form a spiro-cyclopropane ring;

15 R⁶ is C₁-C₆ alkyl; C₁-C₆ haloalkyl; C₁-C₆ alkoxy; NH(C₁-C₆ alkyl); phenyl optionally substituted with R⁷; benzyl; or C₂-C₈ dialkylamino;


R⁷ is C₁-C₆ alkyl; 1-2 halogen; C₁-C₆ alkoxy; or CF₃;

R⁸ is H; C₁-C₆ alkyl; C₁-C₆ alkoxy; or NH(C₁-C₆ alkyl);


20 J is




J-1




J-2




J-3




J-4




J-5




J-6




J-7




J-8




J-9




J-10




J-11




J-12




J-13




J-14




J-15



J-16



J-17



J-18

wherein the dashed line in J-5, J-6 and J-12 indicates that the left-hand ring contains only single bonds or one bond in the ring is a carbon-carbon double bond;

5 n and m are each independently 0; 1; 2; or 3; provided that m + n is 2 or 3;

Z is CR^9R^{10} ; O; S; S(O); S(O)₂; N(C₁-C₄ alkyl); or $\text{N}^{\oplus}\begin{matrix} \text{R} \\ \diagup \\ \text{O}^{\ominus} \end{matrix}$;

each R⁹ is independently H; C₁-C₃ alkyl; halogen; hydroxy; C₁-C₆ alkoxy; C₁-C₆ haloalkyl; C₁-C₆ haloalkoxy; C₂-C₆ alkylcarbonyloxy; or C₂-C₆ haloalkylcarbonyloxy;

10 each R¹⁰ is independently H; C₁-C₃ alkyl; hydroxy; or halogen;

R¹¹ and R¹² are each independently H; halogen; C₁-C₆ alkyl; C₃-C₆ alkenyl; or C₁-C₆ haloalkyl;

R¹³ is H; C₁-C₆ alkyl; C₁-C₆ haloalkyl; C₃-C₆ alkenyl; C₃-C₆ haloalkenyl; C₃-C₆ alkynyl; C₃-C₆ haloalkynyl; HC(=O); (C₁-C₄ alkyl)C(=O); or NH₂;

5 R¹⁴ is C₁-C₆ alkyl; C₁-C₆ alkylthio; C₁-C₆ haloalkyl; CF₃; or N(CH₃)₂;

W is N or CR¹⁵;

R¹⁵ is H; C₁-C₆ alkyl; halogen; or phenyl optionally substituted with C₁-C₆ alkyl, 1-2 halogen, C₁-C₆ alkoxy, or CF₃;

each Q is independently O or S;

10 Q¹ is O or S;

Z¹ is CR¹⁶R¹⁷; O; S; S(O); S(O)₂; N(C₁-C₄ alkyl); or $\overset{\oplus}{\text{N}} \begin{matrix} \text{R} \\ \diagup \\ \text{O}^{\ominus} \end{matrix}$;

each R¹⁶ is independently H; halogen; hydroxy; C₁-C₆ alkoxy; C₁-C₆ haloalkyl;

C₁-C₆ haloalkoxy; C₂-C₆ alkylcarbonyloxy; or C₂-C₆ haloalkylcarbonyloxy;

each R¹⁷ is independently H; hydroxy; or halogen; or

15 when R¹⁶ and R¹⁷ are bonded to adjacent atoms they can be taken together with

the carbons to which they are attached to form $\text{—HC—} \begin{matrix} \text{CH}_2 \\ \diagup \\ \text{CH—} \end{matrix}$;

R¹⁸ is C₁-C₆ alkyl; halogen; or C₁-C₆ haloalkyl;

R¹⁹ and R²⁰ are each independently H; C₁-C₆ alkyl; or C₁-C₆ haloalkyl;

R²¹ and R²² are each independently C₁-C₆ alkyl; C₁-C₆ haloalkyl; C₃-C₆ alkenyl;

20 C₃-C₆ haloalkenyl; C₃-C₆ alkynyl; or C₃-C₆ haloalkynyl;

R²³ is halogen or cyano;

R²⁴ is C₁-C₆ alkylsulfonyl; C₁-C₆ alkyl; C₁-C₆ haloalkyl; C₃-C₆ alkenyl; C₃-C₆ alkynyl; C₁-C₆ alkoxy; C₁-C₆ haloalkoxy; or halogen;

R²⁵ is C₁-C₆ alkyl; C₁-C₆ haloalkyl; C₃-C₆ alkenyl; or C₃-C₆ alkynyl; and

25 R²⁶ is C₁-C₆ alkyl; C₁-C₆ haloalkyl; or phenyl optionally substituted with C₁-C₆ alkyl, 1-2 halogen, 1-2 nitro, C₁-C₆ alkoxy, or CF₃;

provided that when X is O or S in Formula II, then J is other than J-6 or J-7.

In the above recitations, the term "alkyl", used either alone or in compound words such as "alkylthio" or "haloalkyl" includes straight-chain or branched alkyl, such as,

30 methyl, ethyl, *n*-propyl, *i*-propyl, or the different butyl, pentyl or hexyl isomers.

"Alkenyl" includes straight-chain or branched alkenes such as 1-propenyl, 2-propenyl, and the different butenyl, pentenyl and hexenyl isomers. "Alkenyl" also includes

polyenes such as 1,3-hexadienyl and 2,4,6-heptatrienyl. "Alkynyl" includes straight-

chain or branched alkynes such as 1-propynyl, 3-propynyl and the different butynyl,

35 pentynyl and hexynyl isomers. "Alkynyl" can also include moieties comprised of multiple

triple bonds such as 2,7-octadiynyl. "Alkoxy" includes, for example, methoxy, ethoxy, *n*-propyloxy, isopropyloxy and the different butoxy, pentoxy and hexyloxy isomers. "Alkoxyalkyl" denotes alkoxy substitution on alkyl. "Alkoxyalkoxy" denotes alkoxy substitution on alkoxy. Examples of "alkoxyalkyl" include CH₃OCH₂, CH₃OCH₂CH₂,
 5 CH₃CH₂OCH₂, CH₃CH₂CH₂CH₂OCH₂ and CH₃CH₂OCH₂CH₂. "Alkylthio" includes branched or straight-chain alkylthio moieties such as methylthio, ethylthio, and the different propylthio, butylthio, pentylthio and hexylthio isomers. Examples of "alkylsulfonyl" include CH₃S(O)₂, CH₃CH₂S(O)₂, CH₃CH₂CH₂S(O)₂, (CH₃)₂CHS(O)₂ and the different butylsulfonyl, pentylsulfonyl and hexylsulfonyl isomers. "Dialkylamino"
 10 is defined analogously to the above examples. "Cycloalkyl" includes, for example, cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl.

The term "halogen", either alone or in compound words such as "haloalkyl", includes fluorine, chlorine, bromine or iodine. The term "1-2 halogen" indicates that one or two of the available positions for that substituent may be halogen. Further, when used
 15 in compound words such as "haloalkyl", said alkyl may be partially or fully substituted with halogen atoms which may be the same or different. Examples of "haloalkyl" include F₃C, ClCH₂, CF₃CH₂ and CF₃CCl₂. The terms "haloalkenyl", "haloalkynyl" "haloalkoxy", and the like are defined analogously to the term "haloalkyl". Examples of "haloalkenyl" include (Cl)₂C=CHCH₂ and CF₃CH₂CH=CHCH₂. Examples of
 20 "haloalkynyl" include HC≡CCHCl, CF₃C≡C, CCl₃C≡C and FCH₂C≡CCH₂. Examples of "haloalkoxy" include CF₃O, CCl₃CH₂O, HCF₂CH₂CH₂O and CF₃CH₂O. Examples of "haloalkylsulfonyl" include CF₃SO₂, CCl₃SO₂, CF₃CH₂SO₂ and CF₃CF₂SO₂.

The total number of carbon atoms in a substituent group is indicated by the "C_i-C_j" prefix where *i* and *j* are numbers from 1 to 8. For example, C₁-C₃ alkylsulfonyl
 25 designates methylsulfonyl through propylsulfonyl; C₂ alkoxyalkoxy designates CH₃OCH₂O; C₃ alkoxyalkoxy designates, for example, CH₃CH(OCH₃)O, CH₃OCH₂CH₂O or CH₃CH₂OCH₂O; and C₄ alkoxyalkoxy designates the various isomers of an alkoxy group substituted with a second alkoxy group containing a total of four carbon atoms, examples including CH₃CH₂CH₂OCH₂O, and CH₃CH₂OCH₂CH₂O.
 30 Examples of "alkylcarbonyl" include C(O)CH₃, C(O)CH₂CH₂CH₃ and C(O)CH(CH₃)₂. Examples of "alkoxycarbonyl" include CH₃OC(=O), CH₃CH₂OC(=O), CH₃CH₂CH₂OC(=O), (CH₃)₂CHOC(=O) and the different butoxy-, pentoxy- or hexyloxycarbonyl isomers.

When a group contains a substituent which can be hydrogen, for example R¹ or
 35 R⁹, then, when this substituent is taken as hydrogen, it is recognized that this is equivalent to said group being unsubstituted by that substituent.

Compounds of this invention can exist as one or more stereoisomers. The various stereoisomers include enantiomers, diastereomers, atropisomers and geometric isomers.


One skilled in the art will appreciate that one stereoisomer may be more active and/or may exhibit beneficial effects when enriched relative to the other stereoisomer(s) or when separated from the other stereoisomer(s). Additionally, the skilled artisan knows how to separate and/or to selectively prepare said stereoisomers. Accordingly, the present invention comprises mixtures, individual stereoisomers, and optically active mixtures of compounds of Formulae I and II as well as agriculturally suitable salts thereof.

The salts of the compounds of the invention include acid-addition salts with inorganic or organic acids such as hydrobromic, hydrochloric, nitric, phosphoric, sulfuric, acetic, butyric, fumaric, lactic, maleic, malonic, oxalic, propionic, salicylic, tartaric, 4-toluenesulfonic or valeric acids. The salts of the compounds of the invention also include those formed with organic bases (e.g., pyridine, ammonia, or triethylamine) or inorganic bases (e.g., hydrides, hydroxides, or carbonates of sodium, potassium, lithium, calcium, magnesium or barium) when the compound contains an acidic group.

Preferred compounds, compositions containing them, and methods of their use for reasons of better activity and/or ease of synthesis are:

Preferred 1. Compounds of Formulae I and II above, and agriculturally-suitable salts thereof, wherein:

- X is a direct bond or O;
- R¹ is F or Cl;
- R² is F; Cl; or Br;
- R³ is H; C₁-C₆ alkyl; C₃-C₆ cycloalkyl; C₃-C₆ alkenyl; C₃-C₆ alkynyl; C₁-C₆ haloalkyl; C₂-C₆ alkoxyalkyl; C₃-C₆ haloalkenyl; C₁-C₆ alkylsulfonyl; C₃-C₆ alkoxycarbonylalkyl; C(O)R⁶; or benzyl optionally substituted on the phenyl ring with R⁷;
- R⁴ is H or halogen;
- R⁵ is H;
- J is J-1; J-2; J-3; J-4; J-5; J-6; J-8; J-12; J-15; or J-16;
- Z is CR⁹R¹⁰; O; S; or N(C₁-C₄ alkyl);
- each R⁹ is independently H; halogen; or C₁-C₆ haloalkoxy;
- each R¹⁰ is independently H or halogen;
- each Q is O;
- Q¹ is S;
- Z¹ is CR¹⁶R¹⁷; O; S; or N(C₁-C₄ alkyl);
- each R¹⁶ is independently H; halogen; or haloalkoxy;
- each R¹⁷ is independently H or halogen; or
- when R¹⁶ and R¹⁷ are bonded to adjacent atoms they can be taken together with the carbons to which they are attached to form



R¹⁸ is *t*-butyl or CH₂CH₂CH₂F;

R²³ is Cl or cyano; and

R²⁴ is C₁-C₄ haloalkyl or C₁-C₆ haloalkoxy.

5 Preferred 2. Compounds of Preferred 1 wherein:

R² is F or Cl;

R³ is C₁-C₆ alkyl; C₃-C₆ cycloalkyl; C₃-C₆ alkenyl; C₃-C₆ alkynyl; C₃-C₆ alkoxy-carbonylalkyl; or benzyl optionally substituted on the phenyl ring with R⁷;

10 J is J-1; J-2; J-3; J-4; J-5; J-6; J-8; or J-12;

Z is CR⁹R¹⁰ or O;

each R⁹ is independently H or halogen;

R¹¹ is C₁-C₄ haloalkyl;

R¹² is C₁-C₄ alkyl;

15 R¹³ is CH₃;

R¹⁴ is CF₃;


W is CH;

Z¹ is CR¹⁶R¹⁷ or O;

each R¹⁶ is independently H or halogen;

20 each R¹⁷ is independently H or halogen; or

when R¹⁶ and R¹⁷ are bonded to adjacent atoms they can be taken together with the carbons to which they are attached to form



Preferred 3. Compounds of Preferred 2 wherein:

25 R² is Cl;

R³ is C₁-C₆ alkyl; C₃-C₆ alkenyl; or C₃-C₆ alkynyl;

J is J-1; J-2; J-3; J-4; J-6; or J-12;

Z is CR⁹R¹⁰;

R⁹ is independently H or F;

30 R¹⁰ is independently H or F;

R¹¹ is CF₂H;

R¹² is CH₃;

Z¹ is CR¹⁶R¹⁷;

R¹⁶ is independently H or F; and


35 R¹⁷ is independently H or F.

Most preferred are compounds of Preferred 1 selected from the group:
 3-[7-chloro-5-fluoro-1,3-dihydro-1-[(4-methoxyphenyl)methyl]-2,1-benzisothiazol-4-yl]-1-methyl-6-(trifluoromethyl)-2,4(1*H*,3*H*)-pyrimidinedione *S,S*-dioxide;

- 5 3-(7-chloro-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl)-1-methyl-6-(trifluoromethyl)-2,4(1*H*,3*H*)-pyrimidinedione *S,S*-dioxide;
 3-(7-chloro-1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl)-1-methyl-6-(trifluoromethyl)-2,4(1*H*,3*H*)-pyrimidinedione *S,S*-dioxide; and
 10 3-[(7-chloro-1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl)imino]tetrahydro-1*H*,3*H*-[1,3,4]thiadiazolo[3,4-*a*]pyridazin-1-one *S,S*-dioxide.

The invention further comprises anilines of Formula 1 and 2 wherein R^1 , R^2 , R^3 , R^4 , R^5 and X are as defined for compounds of Formula I and II above. The anilines are intermediates in the preparation of compounds of Formula I and II.

15




DETAILS OF THE INVENTION

The compounds of Formulae I and II can be prepared by one or more of the
 20 following methods, or variations obvious to one skilled in the art, as described below in Schemes 1-21. The definitions of Z , Z^1 , Q , Q^1 , X , W , n , m , and R^1 through R^{26} in the compounds of Formulae 1-46 below are as defined above in the Summary of the Invention. Compounds of Formulae Ia-Ir are various subsets of the compounds of Formula I, and all substituents for Formulae Ia-Ir are as defined above for Formula I.

25 Compounds of Formulae I and II are prepared from the corresponding anilines of Formulae 1 and 2, respectively, as represented in Scheme 1.


10

Scheme 1Synthesis of Anilines of Formulae 1 and 2

Anilines of Formulae 1 and 2 wherein X is a direct bond (Formula 2a) are prepared by the method illustrated in Scheme 2. The nitro aniline of Formula 3 is treated with a sulfonyl chloride in the presence of a base such as pyridine, *N,N*-dimethylaminopyridine (DMAP), or triethylamine at -20° to 25°C to obtain the sulfonamide of Formula 4. For compounds wherein R^3 is other than H, the sulfonamide nitrogen can be alkylated or acylated to give the R^3 -substituted compound of Formula 5. The alkylation is performed using an alkyl halide or alkyl sulfonate in the presence of a base such as potassium carbonate, sodium methoxide, potassium *t*-butoxide (*t*-BuOK), or sodium hydride in an anhydrous solvent such as dimethylformamide (DMF), tetrahydrofuran, or acetonitrile at ambient temperature to 80°C . Acylations to form the $\text{R}^3 = \text{C}(\text{O})\text{R}^6$ compounds are accomplished by condensing the sulfonamide of Formula 4 with the appropriate acylating agent, for example an acyl chloride, isocyanate or carbamoyl chloride.

11

Scheme 2



The cyclization to form the cyclic sulfonamide functionality is achieved via intramolecular vicarious nucleophilic substitution of hydrogen as described by K. Wojciechowski and M. Makosza in *Synthesis*, (1992), 571. Treatment of the sulfonamide of Formula 5 with a base such as sodium hydroxide, sodium methoxide, or potassium *t*-butoxide in an inert solvent such as dimethyl sulfoxide (Me₂SO), dimethylformamide or tetrahydrofuran at a temperature between -60°C and ambient

temperature affords the cyclic sulfonamides of Formulae 6 and 7. The ratio of products depends on the nature of R^2 , and the base and solvent used in the cyclization. If R^2 is other than hydrogen, only compounds of Formula 6 are obtained. When R^2 is hydrogen and a mixture of products is obtained, the isomers can be separated by chromatography or crystallization techniques.


The nitro group in the compounds of Formulae 6 and 7 can be reduced to the corresponding anilines of Formula 1 and 2a, respectively, using iron-HCl, $\text{SnCl}_2 \cdot \text{H}_2\text{O}$, or hydrogen over palladium on carbon.

In some instances, it may be desirable to introduce a non-hydrogen R^2 group in anilines of Formula 1 after cyclization. For example, introduction of a chlorine or bromine atom to give anilines of Formula 1a can be accomplished using *N*-chlorosuccinimide (NCS) or *N*-bromosuccinimide (NBS), respectively, in an inert solvent such as dimethylformamide (DMF) at a temperature between ambient temperature and 80°C .

The nitro compounds of Formula 3 are prepared as illustrated in Scheme 3. The amino group of the aniline of Formula 8 is protected with a suitable protecting group, for example as the phthalimide by reaction with phthalic anhydride, to give the protected aniline of Formula 9. For suitable amino protecting groups, see T. W. Greene and P. G. M. Wuts, *Protective Groups in Organic Synthesis*, 2nd Edition, John Wiley and Sons, Inc., New York, (1991).

Nitration of 9 using a nitric acid/sulfuric acid mixture at -20° to 20°C affords the nitro compound of Formula 10. Deprotection of the amino group (using hydrazine in the case of the phthalimide group) affords the nitroaniline of Formula 3.


Scheme 3



The anilines of Formula 8 are known or can be prepared by well-known methods (see March, J., *Advanced Organic Chemistry*, (1992), 4th Ed., John Wiley and Sons, Inc., pp 591, 1103-1105).

Anilines of Formulae 1 and 2a can also be prepared by introduction of the nitro group further along in the synthetic pathway as illustrated in Scheme 4.

Scheme 4




The aniline of Formula 8 is contacted with a sulfonyl chloride as described above to give the sulfonamide of Formula 11. *N*-Acylation with an acyl chloride or the like, or *N*-alkylation with an R³-halide or -sulfonate in the presence of a base, as described above provides the *N*-substituted sulfonamide of Formula 12. Nitration to form the nitro compound of Formula 13 can be achieved using a nitric acid/sulfuric acid mixture at -20° to 20°C. Vicarious intramolecular nucleophilic substitution followed by reduction of the nitro group using the conditions described previously affords the anilines of Formulae 1 and 2a.

Anilines of Formula 2 wherein X is other than a direct bond (Formula 2b) can be prepared starting from the corresponding amino phenol (X=O), amino thiophenol (X=S),

or diamine ($X=NH$ or $N(C_1-C_3 \text{ alkyl})$) of Formula 14 as illustrated in Scheme 5. Using conditions described above, the aniline is treated with a sulfonyl chloride to give the sulfonamide of Formula 15. Subsequent treatment with a base, for example potassium carbonate, induces cyclization. Nitration with nitric and sulfuric acid affords the corresponding nitro compound of Formula 16. *N*-Alkylation or *N*-acylation to introduce R^3 followed by reduction of the nitro group using the reaction conditions described above gives the aniline of Formula 2b.

Scheme 5




10

A similar but alternative method for the preparation of anilines of Formula 2b is outlined in Scheme 6. Starting with the nitro compound of Formula 17, the phenol ($X=O$), thiol ($X=S$), or amino ($X=NH$ or $N(C_1-C_3 \text{ alkyl})$) group is protected with a suitable protecting group (see T. W. Greene and P. G. M. Wuts, *Protective Groups in Organic Synthesis*, 2nd Edition, John Wiley and Sons, Inc., New York, (1991)). For example, a benzyl (Bn) group is a particularly suitable protecting group and is introduced by treatment with benzyl bromide (BnBr) in the presence of K_2CO_3 . Reduction of the nitro group, for example with iron and HCl as described above, affords the aniline of Formula 18. Condensation with a sulfonyl chloride followed by *N*-alkylation or *N*-acylation affords the *N*-substituted sulfonamide of Formula 19.

20

15

Scheme 6

Removal of the protecting group under the appropriate conditions (for example, by treatment with hydrogen over palladium on carbon in the case of an OBn group) and treatment with base induces cyclization to form the cyclic sulfonamide. Nitration and reduction as described above affords the aniline of Formula 2b.


Converting Anilines of Formulae 1 and 2 to Compounds of Formulae I and II

The anilines prepared by the methods outlined in Schemes 1-6 are used in the condensation with J group derivatives to form compounds of Formulae I and II. In some instances, the anilines are used directly in the condensation reactions. In other instances and depending on the nature of the J-group, the NH₂ of the aniline is first converted to another functional group prior to condensation. For example, the aniline may be converted first to a hydrazine, an isocyanate or an aryl iodide. These methods are described in more detail below.

15 Direct Coupling with the Anilines

In some instances where the aniline is used directly, the compounds of Formulae I and II are prepared by condensation of the aniline with an anhydride precursor to the J group. For example, as illustrated in Scheme 7, the anhydride of Formula 20 is condensed with the aniline of Formula 1 to give compounds of Formula Ia wherein J = J-1. This method is disclosed in EP-A-170,191.

16

Scheme 7

The anhydride of Formula 20 can be prepared by methods disclosed in EP 493,721, and WO 91/06216. Compounds of Formulae I and II wherein J = J-11 and J-18 can be prepared using similar methodology. The aniline is condensed with the appropriate J-group anhydride, diester, or other bis-electrophile to form the compound of Formula I or II. The synthesis of the J-11 precursor and the condensation reaction is described in EP 454,444 and EP 415,642. The J-18 group anhydride and the aniline condensation reaction are described in U.S. 4,003,926.


The sulfonamides of Formulae I and II wherein J = J-10 are also prepared directly from the anilines of Formula 1 and 2, respectively. The J-10 group precursor and the aniline condensation reaction are described in WO 94/03459.

Hydrazines

For some compounds of Formulae I and II, the appropriate aniline is first converted to the corresponding hydrazine, and then the hydrazine is condensed with the J-group derivative, or precursor thereof, to form the desired material. The conversion of an aniline of Formula 1 to a hydrazine of Formula 21 is illustrated in Scheme 8. Subsequent condensation of the hydrazine with the iminoether precursor to J-2 followed by cyclization with phosgene forms the cyclic sulfonamide of Formula Ib. The preparation of the iminoether J-2 precursor and the condensation procedure is described in U.S. 4,315,767.

17


Scheme 8



Anilines can be converted to the hydrazines by diazotization and then reduction as is well-known in the literature (for example, see U.S. 4,695,312).


Compounds of Formulae I and II wherein $J = J-3$ and $J-8$ are also prepared by first
 5 converting the aniline to the appropriate hydrazine, and then condensation with the appropriate J-group precursor. Methods for the preparation of the J-3 precursor and the condensation are described in U.S. 5,215,958, WO 90/02120, and U.S. 4,818,275. Methods for the preparation of the J-8 precursor and the condensation with a hydrazine are described in WO 92/12139 and U.S. 4,560,752.

For some compounds of Formulae I and II wherein $J = J-2$ and Z is CR^9R^{10} , the
 10 synthesis is carried out as illustrated below in Schemes 9-17. The retrosynthetic analysis for the synthesis of the J-2 group is shown below (Scheme 9). The formation of ring A can be accomplished by intramolecular cyclization between the nitrogen in ring B and the terminal double bond of the triazololinone with the cyclic sulfonamide group already in place.
 15




The synthesis of the triazololinone ring B is known in the art and can be prepared by methods such as those described in U.S. 4,818,275 and U.S. 4,818,276. Acidic

19

Scheme 11

- The alcohol of Formula Ic can also be prepared by hydroxybromination of the olefin in the starting triazololinone using *N*-bromosuccinimide (NBS) and water or *N*-bromoacetamide and water followed by cyclization of the resulting bromohydrin of Formula 26 using potassium carbonate in an inert solvent such as acetonitrile or acetone (Scheme 12).


Scheme 12

- Compounds of Formula Ic can also be converted to the chloro-, bromo-, and iodo-R⁹ substituted triazololinones of Formula I wherein J = J-2 using methods known to those skilled in the art. For example, the hydroxy group in compounds of Formula Ic can be acylated by known methods to prepare the alkylcarbonyloxy and haloalkylcarbonyloxy derivatives. In addition, the hydroxy or halo group can be converted by known methods to afford the R⁹ = alkoxy and haloalkoxy derivatives (March, J., *Advanced Organic Chemistry*, (1992), 4th Ed., John Wiley and Sons, Inc., pp 386-389). In fact all the R⁹ = OH or halogen compounds in Schemes 13-17 can be functionalized as is known in the art to prepare the other R⁹ substituted compounds.

- Compounds of Formulae Ie and If can be prepared as illustrated in Scheme 13. Oxidative cleavage of the olefin of the triazololinone using sodium periodate and a catalytic amount of osmium tetroxide in an inert solvent mixture such as tetrahydrofuran

(THF) and water, or using ozone in dichloromethane followed by reductive workup using dimethyl sulfide (DMS), affords the aldehyde of Formula 27. Intramolecular cyclization of the aldehyde under basic conditions using potassium carbonate or sodium hydride gives the 5-membered ring alcohol of Formula Ie. Treatment of the alcohol of Formula Ie with DAST in an inert solvent such as dichloromethane at a temperature between about -80°C and about 60°C produces the fluoride of Formula If. The fluoride If can also be made by direct cyclization and fluorination using DAST at a temperature range between about -100°C and about 60°C.

Scheme 13




10

Hydroboration of the olefin of the triazolone with 9-borabicyclo[3.3.1]nonane (9-BBN) gives the alcohol of Formula 28 after oxidative workup (Scheme 14). Oxidation of alcohol 28 with oxidizing agents such as PDC (pyridinium dichromate) produces the hemiaminal of Formula Ig presumably via the aldehyde intermediate of Formula 29. The fluoro compound of Formula Ih can be made by the treatment of the hemiaminal with DAST.

15


21

Scheme 14



Triazolones of Formula 30 can be made from 2-oxo-4-pentenoic acids and the substituted phenyl hydrazines by methods known to those skilled in the art and the methods taught herein (Scheme 10). The epoxide of Formula 31 can be prepared from the olefin of Formula 30 using an oxidant such as MCPBA. Intramolecular cyclization of the epoxide with a base such as potassium carbonate affords the alcohol of Formula Ii. The fluoro compound of Formula Ij can be made from the alcohol using DAST at a temperature between about -70°C and about room temperature in an inert solvent such as dichloromethane.


Scheme 15



Compounds of Formula Im can be prepared from the unsubstituted compounds of Formula 32 as illustrated in Scheme 16. Treatment of the bicyclic triazolone of

Formula 32 with *N*-bromosuccinimide (NBS) under allylic bromination conditions affords the mono-bromo derivative of Formula Ik.


Hydrolysis of the bromide in hot aqueous dimethyl sulfoxide (Me₂SO) affords the alcohol of Formula II. The fluoro compound of Formula Im can be prepared by
5 treatment of the alcohol with diethylaminosulfur trifluoride (DAST) as described above.



Compounds of Formula In can be prepared as illustrated in Scheme 17. Allylic
10 oxidation of the terminal alkene of the triazolinone with *t*-butyl hydroperoxide and catalytic selenium (IV) oxide in an inert solvent such as dichloromethane produces the allylic alcohol of Formula 33. Protection of the secondary alcohol as the
t-butyldimethylsilyl (TBS) ether is accomplished using *t*-butylchlorodimethylsilane and a base, preferably triethylamine and catalytic 4-(dimethylamino)pyridine. The terminal
15 olefin is converted to the primary alcohol to afford compounds of Formula 34 using 9-BBN followed by treatment with sodium perborate. Ring cyclization is accomplished using the Martin sulfurane dehydrating agent [C₆H₅(CF₃)₂O]₂S(C₆H₅)₂. Removal of the TBS group and liberation of the alcohol can be accomplished using tetrabutylammonium fluoride. The alcohol can be converted to the corresponding fluoro
20 compound using DAST, or to other R⁹ substituted compounds as described above.

23

Scheme 17




For some compounds of Formulae I and II wherein J = J-2 and R¹⁰ is other than hydrogen, the R¹⁰ substituent is more conveniently introduced after cyclization to form the bicyclic triazolone. This is especially the case when R⁹ and R¹⁰ are attached to the same carbon atom.

Isocyanates

In some instances, the appropriate aniline is first converted to the corresponding isocyanate, and then the isocyanate is condensed with the J-group derivative, or precursor thereof, to form compounds of Formulae I and II. In Scheme 18, the conversion of aniline of Formula 1 to isocyanate of Formula 35 is illustrated. Subsequent condensation of the isocyanate with the aminoester of Formula 36 forms the cyclic sulfonamide of Formula Io. The preparation of some aminoester precursors to J-6 and the condensation procedure is described in U.S. 4,179,276.

24


Scheme 18

Compounds of Formulae I and II wherein J = J-4, J-5, J-7, J-9, J-11, J-12, J-13, J-14 and J-17 are also prepared by first converting the aniline to the appropriate isocyanate, and then condensation with the appropriate J-group precursor. Methods for the preparation of the J-4 precursor and the condensation are described in WO 92/11244, EP 476,697, ZA 91/00466, JP 77,874, and U.S. 3,902,887. The synthesis of the J-5 precursor and the condensation with isocyanates is described in WO 92/13453 JP 363,278, EP 230,874, and JP 2,000,985. Methods for the preparation of the J-7 precursor and the condensation with isocyanates are described in EP 484,776. Methods for the preparation of the J-9 precursor and the condensation with isocyanates are described in U.S. 5,035,740. Methods for the preparation of the J-17 precursor and its condensation with isocyanates are described in EP 493,323. The synthesis of the J-14 precursor and the condensation with isocyanates is described in *J. Pesticide Sci.*, (1993), 18, 309. In a similar vein, the imino compounds of Formulae I and II wherein J = J-12 and J-13 can be prepared from the corresponding isocyanates of the anilines. The condensation procedure and J-group precursor preparation for compounds containing J-12 and J-13 are disclosed in EP 457,151, JP 4,145,087, EP 480,871 and DE 3,927,388.

One skilled in the art will recognize that when Q or Q¹ is S in the desired product, the appropriate isothiocyanate is used instead of the isocyanate in the synthesis.

For some compounds of Formulae I and II wherein J = J-4, J-5, J-6, J-7, J-9, J-14 and J-17, the coupling can also be accomplished starting with the aniline rather than the isocyanate. For example, the synthesis of compounds of Formula Ip (compounds of

Formula I wherein R¹⁶ and R¹⁷ are taken together to form a cyclopropane ring) is illustrated in Scheme 19.



5

Treatment of cyclopropane dicarboxylic acid of Formula 37 with urea and heating to 175-185°C affords the dicarboximide of Formula 38 as described by G. C. Crockett et al. in *Synth. Commun.* (1981), 11, 447-454. The diester of the diacid of Formula 37 is prepared by the method described by L. L. McCoy in *J. Am. Chem. Soc.*, (1958), 80, 65-68. The diacid can be obtained by saponification of the diester using well-known methods. Reduction of the dicarboximide of Formula 38 with borane in an inert solvent, such as tetrahydrofuran (THF), followed by work-up with aqueous hydrochloric acid affords the azabicyclo[3.1.0]hexane hydrochloride of Formula 39. The reduction is preferably conducted with heating, for example in THF at reflux, as described by H. C. Brown and P. Heim in *J. Org. Chem.*, (1973), 38, 912-916.

The amine hydrochloride of Formula 39 is converted via a five step sequence to the α-amino acid of Formula 43 as illustrated. Purification of the intermediates is not necessary. Neutralization of the amine hydrochloride with a base, such as concentrated aqueous potassium hydroxide, liberates the free amine. Dissolution of the amine in an inert solvent, such as diethyl ether, and treatment with a solution of *N*-chlorosuccinimide (NCS) in an inert solvent such as ether, produces the chloramine of Formula 40. The

20

solution of the chloramine is then treated with ethanolic potassium hydroxide to effect dehydrochlorination and give the imine of Formula 41. Once again, the imine is not purified but treated directly first with aqueous sodium bisulfite, and then with solid sodium cyanide to afford the aminonitrile of Formula 42. The reaction mixture is poured
5 into water and extracted with a water-immiscible solvent such as ether. The organic layers are dried and evaporated under reduced pressure to afford the aminonitrile. No additional purification is necessary. The aminonitrile can be converted to the aminoacid of Formula 43 by hydrolysis with aqueous barium hydroxide followed by neutralization with sulfuric acid. A mixture of epimers at the carboxylic acid centers is obtained, and
10 the individual diastereomers can be separated by chromatography.

The acid of Formula 43 is reacted with the aniline of Formula and a trialkylaluminum reagent (e.g., trimethylaluminum), in a non-coordinating solvent such as an aromatic hydrocarbon (e.g., benzene and toluene) or halogenated hydrocarbon (e.g., methylene chloride, chloroform, carbon tetrachloride, and dichlorobutane) to obtain the
15 amide. Generally, the reaction requires 0.1 to 48 hours at a temperature of 0°C to 25°C to proceed to completion. The amides are isolated by extraction into an organic solvent, aqueous wash, and removal of the solvent under reduced pressure. Purification can be accomplished by chromatography or recrystallization. The condensation with the amine can also be performed starting with the ester of the acid of Formula 43.

20 The tricyclic imide of Formula Ip can be prepared from the α -aminoamide by condensation with phosgene or a phosgene equivalent. Treatment of the α -aminoamide with phosgene is preferably carried out in the presence of a tertiary-amine base such as triethylamine, pyridine, or *N,N*-diisopropylethylamine, in an inert solvent such as dichloromethane or 1-chlorobutane. The phosgene can be added as a gas or as a
25 solution in an inert solvent such as toluene. Suitable temperatures range from about 0°C to the reflux temperature of the solvent. 1,1'-Carbonyldiimidazole, diphosgene ($\text{ClC}(=\text{O})\text{OCCl}_3$) and triphosgene ($\text{Cl}_3\text{COC}(=\text{O})\text{OCCl}_3$) can also be used in a similar manner.

The tricyclic imide of Formula Ip can be isolated by extraction into an organic
30 solvent, aqueous wash, and removal of the solvent under reduced pressure. Additional purification can be accomplished by chromatography or recrystallization.


Aryl Iodides

For the preparation of compounds of Formulae I and II wherein J = J-15 and J-16, the appropriate aniline is first converted to the aryl alkyne as illustrated in Scheme 20.
35 The aniline of Formula 1 is converted to the aryl iodide of Formula 44 via diazotization followed by treatment with a metal iodide salt. The aryl iodide is linked by a palladium coupling reaction to give the trimethylsilyl (TMS) alkyne. Hydrolysis of the TMS group with base affords the terminal alkyne of Formula 45. In the case of J-16, a [3+2]

cycloaddition using a sydnone as the dipole and the alkyne as the dipolarophile affords the bicyclic pyrazole compounds. Introduction of the R^{23} group affords the cyclic sulfonamides of Formula Iq. For example, treatment with *N*-chlorosuccinimide affords the $R^{23} = \text{Cl}$ compound. These methods are described in WO 93/15074.

5

Scheme 20




Methods for the similar preparation of compounds of Formulae I and II wherein $J = \text{J-15}$ are described in JP 4,059,706, WO 92/06962, and JP 3,163,063.

10

For compounds of Formulae I and II wherein $J = \text{J-5}$, the coupling can also be accomplished starting with the aniline rather than the isocyanate. For example, the synthesis of compounds of Formula Ir is illustrated in Scheme 21. Treatment of a diester of Formula 46 with an aniline of Formula 1 in the presence of a trialkylaluminum reagent (e.g., trimethylaluminum) in a non-coordinating solvent such as an aromatic hydrocarbon (e.g., benzene, toluene) or a halogenated hydrocarbon (e.g., methylene chloride, chloroform, carbon tetrachloride and dichlorobutane) affords a compound of Formula Ir.

15

Scheme 21



It is recognized that some reagents and reaction conditions described above for preparing compounds of Formulae I and II may not be compatible with certain functionalities present in the intermediates. In these instances, the incorporation of protection/deprotection sequences or functional group interconversions into the synthesis will aid in obtaining the desired products. The use and choice of the protecting groups will be apparent to one skilled in chemical synthesis (see, for example, Greene, T. W.; Wuts, P. G. M. *Protective Groups in Organic Synthesis*, 2nd ed.; Wiley: New York, 1991). One skilled in the art will recognize that, in some cases, after the introduction of a given reagent as it is depicted in any individual scheme, it may be necessary to perform additional routine synthetic steps not described in detail to complete the synthesis of compounds of Formulae I and II.

One skilled in the art will also recognize that compounds of Formulae I and II and the intermediates described herein can be subjected to various electrophilic, nucleophilic, radical, organometallic, oxidation, and reduction reactions to add substituents or modify existing substituents.

Without further elaboration, it is believed that one skilled in the art using the preceding description can utilize the present invention to its fullest extent. The following Examples are, therefore, to be construed as merely illustrative, and not limiting of the disclosure in any way whatsoever. Percentages are by weight unless otherwise indicated. Parts for chromatographic solvent mixtures are by volume unless otherwise indicated. ¹H NMR spectra are reported in ppm downfield from tetramethylsilane; s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, dd = doublet of doublets, br s = broad singlet.

25

EXAMPLE 1Step A: 1-Chloro-N-(4-fluoro-3-nitrophenyl)methanesulfonamide

To a solution of 4-fluoro-3-nitroaniline (23.4 g, 0.15 mol) in anhydrous pyridine (120 mL) was added dropwise a solution of chloromethylsulfonyl chloride (22.35 g, 0.15 mol) at 0°C under a nitrogen atmosphere. The mixture was then stirred at 0-5°C for 4 h and then the mixture was poured onto ice (1500 g) containing concentrated

30

aqueous hydrochloric acid (300 mL). The mixture was filtered, and the residue was dissolved in methylene chloride (1200 mL). After drying (MgSO₄) and concentration under reduced pressure, the crude product was dissolved in 1-chlorobutane:ether 9:1 v/v and passed through silica gel. The eluent was concentrated under reduced pressure to give the title compound of Step A.

Step B: 1-Chloro-N-ethyl-N-(4-fluoro-3-nitrophenyl)methanesulfonamide

A solution of 1-chloro-N-(4-fluoro-3-nitrophenyl)methanesulfonamide (5.93 g, 22.1 mmol), iodoethane (3.79 g, 24.3 mmol), tetrabutylammonium bromide (0.712 g, 2.21 mmol), and anhydrous K₂CO₃ (8.85 g, 64 mmol) in anhydrous dimethylformamide (40 mL) was stirred at room temperature. When the starting material had disappeared based on thin layer chromatography on silica (6 h), the reaction mixture was poured into water (200 mL) and extracted with CH₂Cl₂ (500 mL). The organic extract was washed with water (two times with 50 mL), dried (MgSO₄), and the solvent was removed under reduced pressure. The residue was purified by flash chromatography (silica gel, 1-chlorobutane and diethyl ether 100:1 v/v eluent) to give 4.68 g of the title compound of Step B as a solid melting at 80-83°C. ¹H NMR (Me₂SO-*d*₆, 400 MHz): δ 8.17 (dd, 1H, J=6.75 and 2.7 Hz), 7.85 (dd, 1H, J=9.4 and 3 Hz), 7.69 (dd, 1H, J=11 and 9 Hz), 5.25 (s, 2H), 3.83 (q, 2H), 1.04 (t, 3H).

Step C: 1-Ethyl-5-fluoro-1,3-dihydro-4-nitro-2,1-benzisothiazole 2,2-dioxide and 1-Ethyl-6-fluoro-1,3-dihydro-5-nitro-2,1-benzisothiazole 2,2-dioxide

1-Chloro-N-ethyl-N-(4-fluoro-3-nitrophenyl)methanesulfonamide (2.0 g, 6.74 mmol) in dimethyl sulfoxide (Me₂SO-*d*₆, 10 mL) was added dropwise to a vigorously stirred suspension of powdered NaOH (4.0 g) in Me₂SO-*d*₆ (30 mL) while maintaining the temperature at 25-27°C. After completion of the addition, the mixture was stirred at 25-27°C for an additional 30 min. After completion of the reaction based on thin layer chromatography on silica gel, the reaction mixture was poured into ice (400 g) and concentrated aqueous hydrochloric acid (60 mL). The precipitate was extracted into CH₂Cl₂ (300 mL), dried (MgSO₄), and subjected to flash chromatography on silica gel using ethyl acetate:hexane 1:4 v/v. Two products were isolated. The first compound to elute was 1-ethyl-6-fluoro-1,3-dihydro-5-nitro-2,1-benzisothiazole 2,2-dioxide (Compound A, 120 mg). The second compound to elute was 1-ethyl-5-fluoro-1,3-dihydro-4-nitro-2,1-benzisothiazole 2,2-dioxide (Compound B, 976 mg). Compound A: ¹H NMR (CDCl₃, 300 MHz): δ 7.33 (dd, 1H, J=10.51 and 9.75 Hz), 6.96 (dd, 1H, J=9.0 and 3.38 Hz), 4.63 (s, 2H), 3.76 (q, 2H), 1.41 (t, 3H). Compound B: m.p. 97-100°C; ¹H NMR (CDCl₃, 300 MHz): δ 7.37 (d, 1H, J=6.10 Hz), 7.27 (d, 1H, J=10.9 Hz), 4.39 (s, 2H), 3.74 (q, 2H), 1.44 (t, 3H).

Step D: 1-Ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-amine 2,2-dioxide

A mixture of 1-ethyl-5-fluoro-1,3-dihydro-4-nitro-2,1-benzisothiazole 2,2-dioxide (1.0 g, 3.8 mmol) and SnCl₂·H₂O (4.34 g, 19.2 mmol) in ethyl acetate (50 mL) was heated at reflux for 3 h. After completion of the reaction based on thin layer chromatography, the mixture was diluted with ethyl acetate (100 mL) and solid Na₂CO₃ (12.5 g) was added. The resulting mixture was vigorously stirred for 12 h and then filtered through a pad of diatomaceous earth (Celite®). The filtrate was then evaporated under reduced pressure, and the residue was subjected to flash chromatography (silica gel, chlorobutane:ether 1:10 v/v) to give 650 mg of the title compound of Step D. ¹H NMR (CDCl₃, 400 MHz): δ 6.94 (m,1H), 6.08 (m,1H), 4.10 (s,2H), 3.80 (br s,2H), 3.62 (q,2H), 1.36 (t,3H).

Step E: 2-(1-Ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazole-4-yl)-4,5,6,7-tetrahydro-1H-isoindole-1,3(2H)-dione S,S-dioxide

A mixture of 1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-amine 2,2-dioxide (506 mg, 2.2 mmol), and 3,4,5,6-tetrahydrophthalic anhydride (335 mg, 2.2 mmol) in acetic acid (10 mL) was heated at reflux for 2 days. Progress of the reaction was monitored by thin layer chromatography. After completion of the reaction, the solvent was removed under reduced pressure and the product was purified by flash chromatography (silica gel, 1-chlorobutane:ether 4:1 v/v) to give 550 mg of the title compound of Step E, a compound of the invention, as a solid melting at 189-190°C. ¹H NMR (CDCl₃, 400 MHz): δ 7.18 (dd,1H), 6.72 (dd,1H), 4.20 (s,2H), 3.68 (q,2H), 2.42 (br s,4H), 1.78 (br s,4H), 1.38 (t,3H).

EXAMPLE 2

2-(7-Chloro-1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl)-4,5,6,7-tetrahydro-1H-isoindole-1,3(2H)-dione S,S-dioxide

A mixture of 2-(1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazole-4-yl)-4,5,6,7-tetrahydro-1H-isoindole-1,3(2H)-dione S,S-dioxide (250 mg, 0.686 mmol) and N-chlorosuccinimide (96 mg, 0.72 mmol) in dimethylformamide (3 mL) was heated at 80°C for 4 h under a nitrogen atmosphere. After completion of the reaction based on thin layer chromatography, the mixture was poured into ether (25 mL) and water (5 mL). The organic phase was separated, dried (MgSO₄) and evaporated under reduced pressure. The residue was then purified by flash chromatography (silica gel, 1-chlorobutane:ether 19:1 v/v as eluent) to give 121 mg of the title compound of Example 2, a compound of the invention, as a solid melting at 190-192°C. ¹H NMR (CDCl₃, 400 MHz): δ 7.26 (dd,1H), 4.18 (s,2H), 3.96 (q,2H), 2.45 (m,4H), 1.84 (m,4H), 1.28 (t,3H).

EXAMPLE 3

Step A: 1-Ethyl-6-fluoro-1,3-dihydro-2,1-benzisothiazol-5-amine 2,2-dioxide

1-Ethyl-6-fluoro-1,3-dihydro-5-nitro-2,1-benzisothiazole 2,2-dioxide
(Compound B from Example 1, Step C, 1.0 g), was suspended in tetrahydrofuran
5 (20 mL) with 10% palladium on carbon (100 mg). The suspension was placed in Parr[®]
shaker and treated with 50 psi (3.45 x 10⁵ Pa) of hydrogen overnight. The catalyst was
then filtered off through diatomaceous earth (Celite[®]), and the filtrate was evaporated
under reduced pressure to give the title compound of Step A as a brown solid
(0.510 mg). ¹H NMR (CDCl₃): δ 6.92 (dd,1H), 6.04 (dd,1H), 4.08 (s,2H), 3.80
10 (br s,2H), 3.64 (q,2H), 1.32 (t,3H).

Step B: 2-(1-Ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-6-yl)-4,5,6,7-
tetrahydro-1H-isoindole-1,3(2H)-dione S,S-dioxide

1-Ethyl-6-fluoro-1,3-dihydro-2,1-benzisothiazol-5-amine 2,2-dioxide (123 mg) and
3,4,5,6-tetrahydrophthalic anhydride (64 mg) were condensed using the procedure
15 described in Example 1, Step E to give 64 mg of the title compound of Step B, a
compound of the invention. ¹H NMR (CDCl₃): δ 7.08 (d,1H), 6.62 (d,2H), 4.32 (s,2H),
3.64 (q,2H), 2.42 (m,4H), 1.82 (m,4H), 1.38 (t,3H).

EXAMPLE 4

Step A: 7-Chloro-1-ethyl-5-fluoro-1,3-dihydro-4-isocyanato-2,1-
20 benzisothiazole 2,2-dioxide

To a mixture of 7-chloro-1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-amine-
2,2-dioxide (1.30 g, 4.91 mmol), diphosgene (1.17 g, 1.2 eq) in p-dioxane (20 mL),
triethylamine (0.6 g, 1.2 eq) was added and the resulting reaction mixture was heated at
reflux for overnight. The precipitated triethylamine hydrochloride was filtered off and
25 the solvents were removed *in vacuo* and dried to give the title compound of Step A as a
brown semisolid. ¹H NMR (CDCl₃, 400 MHz): δ 7.21 (d,1H, J=9.5 Hz), 4.28 (s,2H),
3.94 (q,2H), 1.22 (t,3H, J=7.5 Hz).

Step B: 3-(7-Chloro-1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazole-4-yl)-6-
(trifluoromethyl)-2,4(1H,3H)-pyrimidinedione S,S-dioxide

30 A solution of ethyl 3-amino-4,4,4-trifluorocrotonate (0.899 g, 4.91 mmol) in DMF
(3 mL) was added to sodium hydride (60% suspension in oil, 0.118 g, 4.91 mmol) in
DMF (8.5 mL) at -10 to -5°C. The reaction mixture was stirred at -10 to -5°C for
30 min and was then cooled to -78°C. To the resulting solution was added a solution of
the title compound of Step A in toluene (14 mL). The reaction mixture was stirred for
35 an additional 2 h at -78°C, overnight at room temperature, and was then poured into
HCl/ice mixture and extracted with ethyl acetate (20 mL). The organic phase was
separated, washed with water (10 mL), dried (MgSO₄), filtered and the solvent was
removed on the rotary evaporator. The title compound of Step B, a compound of the

invention, was isolated by flash chromatography as a brown solid (1.18 g, m.p. >230°C, R_f 0.25 (silica gel) 1:1 1-chlorobutane and ether). ^1H NMR ($\text{Me}_2\text{SO}-d_6$, 400 MHz): δ 12.92 (broad, 1H, NH), 7.83 (d, 1H, 10.0 Hz), 6.46 (s, 1H), 4.76 (d, 1H, $J=16.5$ Hz), 4.58 (d, 1H, $J=16.5$ Hz), 3.81 (q, 2H), 1.17 (t, 3H, $J=7.0$ Hz).

5

EXAMPLE 5

3-(7-Chloro-1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl)-1-methyl-6-(trifluoromethyl)-2,4(1H,3H)-pyrimidinedione S,S-dioxide

A mixture of the title compound of Step B in Example 4 (0.33 g, 0.77 mmol), dimethyl sulfate (0.102 g, 1.05 eq) in acetone (10 mL) was heated at reflux in the presence of potassium carbonate (0.213 g) for 3 h. After completion of the reaction (TLC), the reaction mixture was cooled, the insolubles were filtered off and the solvent was removed on a rotary evaporator. The title compound of Example 5, a compound of the invention, was purified by flash chromatography to give a colorless solid melting at 147-149°C (R_f 0.55, 1:1 1-chlorobutane and ether). ^1H NMR (benzene- d_6 , 400 MHz): δ 6.55 (d, 1H, $J=9.5$ Hz), 5.86 (s, 1H), 3.19-3.60 (m, 4H), 2.77 (d, 1H, $J=1.5$ Hz), 0.963 (t, 3H). IR (KBr) 1636, 1691.

10

15

EXAMPLE 6

3-(7-Chloro-1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl)-1-ethyl-6-(trifluoromethyl)-2,4(1H,3H)-pyrimidinedione S,S-dioxide.

A mixture of the title compound of Step B in Example 4 (0.3 g, 0.70 mmol), ethyl iodide (0.13 g, 0.84 mmol), K_2CO_3 (0.12 g, 0.88 mmol) in acetone (10 mL) was heated at reflux for 40 h. The reaction mixture was cooled, the insolubles were filtered off, the solvent was removed *in vacuo* and the title compound of Example 6, a compound of the invention, was purified by flash chromatography to give a solid melting at 177-178°C. ^1H NMR: δ 7.24 (d, 1H), 6.32 (s, 1H), 4.08 (s, 2H), 4.04 (m, 4H), 1.26 (m, 6H).

20

25

EXAMPLE 7

Step A: 7-Chloro-1-ethyl-5-fluoro-1,3-dihydro-4-isothiocyanato-2,1-benzisothiazole 2,2-dioxide

To a solution of 7-chloro-1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-amine 2,2-dioxide (0.5 g, 1.88 mmol) and triethylamine (0.6 mL) in anhydrous tetrahydrofuran (THF, 10 mL), thiophosgene (0.238 g, 1.1 eq) in THF (2 mL) was added dropwise at room temperature under a nitrogen atmosphere. The resulting reaction mixture was heated at reflux for 2 days. The solvents were removed *in vacuo* to give the title compound of Step A which was used in Step B.

30

35

Step B: 3-[(7-Chloro-1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl)imino]tetrahydro-1H,3H-[1,3,4]thiadiazolo[3,4-a]pyridazin-1-one S,S-dioxide

A mixture of the title compound of Step A (0.57 g, 1.88 mmol), 1,2,3,6-tetrahydropyridazine (0.675 g, 1.97 mmol) and K_2CO_3 (1.3 g, 5 eq) in anhydrous THF was stirred at room temperature for 1 h. After completion of the reaction, the solvent was removed and the adduct was purified by flash chromatography. 1H NMR ($CDCl_3$, 300 MHz): δ 9.28 (br s, 1H), 7.08 (d, 1H), 4.21 (s, 2H), 3.86 (m, 3H), 3.56 (m, 1H), 3.04 (m, 2H), 1.82 (m, 4H), 1.26 (t, 3H).

To a solution of this adduct (0.342 g, 0.87 mmol) in anhydrous toluene (5 mL) was added a solution of phosgene (0.098 g) in toluene (3 mL) dropwise under a nitrogen atmosphere and the resulting reaction mixture was stirred at room temperature for 12 h. After completion of the reaction (TLC), the solvent was removed under reduced pressure and the title compound of Step B, a compound of the invention, was purified by flash chromatography to give a solid melting at 143-146°C. 1H NMR ($CDCl_3$, 300 MHz): δ 7.03 (d, 1H, 10.8 Hz), 4.16 (s, 2H), 3.90 (dd, 2H), 3.47 (m, 4H), 1.93 (m, 4H), 1.20 (t, 3H).

15

EXAMPLE 8

2-(7-Chloro-1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl)tetrahydroimidazo[1,5-a]pyridine-1,3(2H,5H)-dione S,S-dioxide

To a solution of pipercolinic acid (0.205 g, 1.59 mmol) in KOH (93 mg, 1.66 mmol), H_2O (5 mL) was added the title compound of Step A in Example 4 (0.44 g, 1.5 mmol) in acetone (2.5 mL). The resulting reaction mixture was stirred at room temperature for 20 h. The solvents were then removed and acidified with 3N HCl to pH 1 to obtain a brown precipitate. The mixture was extracted with ethyl acetate (50 mL) and the organic phase was separated. The organic phase was dried ($MgSO_4$) and the solvent was removed to give a brown solid. This solid was dissolved in ethanol (10 mL) and 3N HCl (10 mL) and heated at reflux for 12 h. The solvents were then removed *in vacuo* and the residue was extracted with ethyl acetate. The extracts were dried ($MgSO_4$) and concentrated to give a brown solid. The title compound of Example 8, a compound of the invention, was isolated by flash chromatography using ethyl acetate:hexane (3:7) as eluent giving a solid melting at 168-169°C. 1H NMR ($CDCl_3$, 400 MHz): δ 7.28 (d, 1H, J=9.5 Hz), 4.17-4.36 (m, 3H), 3.94 (m, 3H), 2.93 (m, 1H), 2.32 (m, 1H), 2.09 (m, 1H), 1.83 (m, 1H), 1.40-1.55 (m, 3H), 1.28 (t, 3H).

30

EXAMPLE 9

2-(7-Chloro-1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl)tetrahydro-1H-imidazo-[5,1-c][1,4]oxazine-1,3(2H)-dione S,S-dioxide

Using similar reaction conditions as described in Example 8, the title compound of Step A in Example 4 (0.44 g, 1.5 mmol) and morpholine carboxylic acid (0.218 g, 1.66 mmol) gave 0.24 g of the title compound of Example 9, a compound of the invention, as a solid melting at 200-202°C. 1H NMR ($CDCl_3$, 400 MHz): δ 7.27 (d, 1H,

35

J=9.5 Hz), 4.36 (m,2H), 4.25 (d,1H), 4.17 (m,1H), 4.03 (m,2H), 3.96 (m,2H), 3.23-3.53 (m,3H), 1.28 (t,3H).

EXAMPLE 10

5 2-(7-Chloro-1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl)-5,8-dihydro-1H-[1,2,4]triazolo[1,2-a]pyridazine-1,3(2H)-dione S,S-dioxide

A solution of trimethylaluminum (0.68 mL 2M in hexane, 1.36 mmol) was added dropwise to 7-chloro-1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-amine 2,2-dioxide (0.3 g, 1.13 mmol) in toluene (5 mL) at room temperature under a nitrogen atmosphere and the resulting reaction mixture was stirred for 10 min. A solution of diethyl 3,6-dihydro-1,2-pyridazinedicarboxylate (0.313 g, 1.36 mmol) in toluene (2 mL) was then added at room temperature and the reaction mixture was heated at reflux for 12 h. After the reaction was complete, the reaction mixture was poured into a solution of ethyl acetate (40 mL) containing 1N HCl (10 mL) and stirred for 10 min. The organic phase was separated, dried, and the solvent was removed on a rotary evaporator and followed by flash chromatography to afford the title compound of Example 10, a compound of the invention, as a semisolid. ¹H NMR (CDCl₃, 300 MHz): δ 7.30 (d,1H, J=9.6 Hz), 5.99 (s,2H), 4.31 (s,2H), 4.21 (s,4H), 3.93 (dd,2H), 1.28 (t,3H).

EXAMPLE 11

20 2-(7-Chloro-1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl)tetrahydro-1H-[1,2,4]triazolo[1,2-a]pyridazine-1,3(2H)-dione S,S-dioxide

Using similar reaction conditions as described in Example 10, 7-chloro-1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-amine 2,2-dioxide (0.46 g, 1.13 mmol) and diethyl hexahydropyridazine-1,2-dicarboxylate (0.314 g, 1.36 mmol) gave the title compound of Example 11, a compound of the invention, as a solid melting at 87-90°C. ¹H NMR (CDCl₃, 400 MHz): δ 7.29 (d,1H), 4.22 (s,2H), 3.95 (q,2H), 3.65 (m,4H), 1.89 (m,4H), 1.28 (t,3H).

EXAMPLE 12

Step A: 7-Chloro-1-ethyl-5-fluoro-4-hydrazino-1,3-dihydro-2,1-benzisothiazole 2,2-dioxide

30 To a suspension of 7-chloro-1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-amine 2,2-dioxide (0.5 g, 1.89 mmol) in water (3.1 mL) and H₂SO₄ (1.65 mL) at 0° was added NaNO₂ (0.13 g, 1.89 mmol) in water (0.5 mL). The reaction mixture was stirred at 0°C for 1 h. A solution of SnCl₂·H₂O (0.85 g, 3.78 mmol) in conc. HCl (3.1 mL) was then added dropwise and the reaction mixture was stirred at room temperature for 12 h. 35 Then 50% aqueous NaOH solution was added until the reaction mixture reached pH 13. The reaction mixture was extracted with ethyl acetate (200 mL) and the organic layer was washed with brine (3 x 50 mL) and dried (MgSO₄). The solvent was removed and

the title compound of Step A was isolated by flash chromatography. ¹H NMR (CDCl₃, 300 MHz): δ 7.04 (d,1H), 5.62 (br s,1H), 4.62 (s,2H), 3.86 (m,2H), 1.26 (t,3H).

Step B:

A mixture of the title compound of Step A (0.5 g, 1.79 mmol), aqueous HCl (10%, 0.1 mL), ethanol (0.2 mL) and sodium pyruvate (0.21 g, 1.87 mmol) was heated at 40°C for 1 h. The reaction mixture was then cooled and diluted with water (2 mL) to give a brown solid which was filtered (0.5 g) and used in Step C.

Step C: 1-(7-Chloro-1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl)-2,4-dihydro-5-methyl-3H-1,2,4-triazol-3-one S,S-dioxide

A mixture of the Step B compound (0.78 g, 2.23 mmol), diphenylphosphoryl azide (0.62 g, 2.25 mmol) and triethylamine (0.24 g, 2.32 mmol) in anhydrous toluene (11 mL) was heated at reflux for 5 h. After completion of the reaction (TLC), the reaction mixture was cooled, diluted with ethyl acetate (50 mL) and washed with brine (10 mL). The organic phase was dried (MgSO₄) and the solvent was evaporated. The resulting residue was purified by flash chromatography to give the title compound of Step C as a solid melting at 195-197°C. ¹H NMR (Me₂SO-*d*₆, 400 MHz): δ 11.92 (br s,1H), 7.79 (d,1H, J=10.25 Hz), 4.71 (s,2H), 3.78 (dd,2H), 2.15 (s,3H), 1.15 (t,3H).

Step D: 2-(7-Chloro-1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl)-4-(difluoromethyl)-2,4-dihydro-5-methyl-3H-1,2,4-triazol-3-one S,S-dioxide

To a mixture of the title compound of Step C (0.348 g, 1 mmol), tetrabutylammonium bromide (TBAB, 0.25 g), NaOH (0.257 g, 6.43 mmol) in cyclohexane (10 mL) was added Freon[®] 22 (chlorodifluoromethane, 0.3 mL, 2.98 mmol) and the reaction mixture was heated at 60°C for 2 h under a Freon[®] 22 filled balloon. After the reaction was complete, the reaction mixture was cooled and the solvent was removed. The title compound of Step D, a compound of the invention, was isolated by flash chromatography as a semisolid. ¹H NMR (CDCl₃, 300 MHz): δ 7.32 (d,1H), 7.02 (t,1H), 4.26 (s,2H), 3.96 (q,2H), 2.49 (s,3H), 1.26 (t,3H).

EXAMPLE 13

Step A: 1-Chloro-N-(4-fluoro-3-nitrophenyl)-N-[(4-methoxyphenyl)methyl]methanesulfonamide

A solution of the title compound of Step A in Example 1 (3 g, 11.16 mmol), 4-methoxybenzyl chloride (2.1 g, 13.4 mmol), tetrabutylammonium bromide (0.36 g, 1.1 mmol) and anhydrous K₂CO₃ (4.5 g, 13.8 mmol) in anhydrous dimethylformamide (20 mL) was stirred at room temperature for 6 h. When the starting material had been consumed based on TLC (silica gel), the reaction mixture was poured into water (100 mL) and extracted with ethyl acetate (250 mL). The organic extract was washed with brine (100 mL), dried (MgSO₄), and the solvent was removed. Flash

chromatography of the residue afforded 2.5 g of the title compound of Step A as an oil. ^1H NMR (CDCl_3 , 300 MHz): δ 8.01 (dd, 1H, $J=6.57$ and 2.82 Hz), 7.48 (m, 1H), 7.21 (dd, 1H, $J=10.13$, 9.01 Hz), 6.80-7.12 (m, 4H), 4.93 (s, 2H), 4.57 (s, 2H), 3.77 (s, 3H).

Step B: 5-Fluoro-1,3-dihydro-1-[(4-methoxyphenyl)methyl]-4-nitro-2,1-benzisothiazole 2,2-dioxide and 5-fluoro-1,3-dihydro-1-[(4-methoxyphenyl)methyl]-6-nitro-2,1-benzisothiazole 2,2-dioxide

5 The title compound of Step A (1.66 g, 4.26 mmol) in dimethyl sulfoxide (Me_2SO , 8 mL) was added dropwise to a vigorously stirred suspension of powdered NaOH (3.32 g) in Me_2SO (15 mL) while maintaining the temperature at $25-27^\circ\text{C}$. After completion of the addition, the reaction mixture was stirred at $25-27^\circ\text{C}$ for an additional 45 min. The reaction mixture was then poured into a mixture of ice (600 g) and conc. HCl (40 mL) and extracted with ethyl acetate (300 mL). The organic extract was dried (MgSO_4) and the solvent was removed. Flash chromatography of the resulting residue afforded two products. The first compound to elute was 5-fluoro-1,3-dihydro-1-[(4-methoxyphenyl)methyl]-6-nitro-2,1-benzisothiazole 2,2-dioxide (Compound A). The second compound to elute was 5-fluoro-1,3-dihydro-1-[(4-methoxyphenyl)methyl]-4-nitro-2,1-benzisothiazole 2,2-dioxide (Compound B). Compound A: ^1H NMR (CDCl_3 , 400 MHz): δ 7.26 (m, 3H), 6.88 (m, 3H), 4.68 (s, 2H), 4.46 (s, 2H), 3.82 (s, 3H). Compound B: ^1H NMR (CDCl_3 , 400 MHz): δ 7.30 (d, 2H), 7.16 (t, 1H, $J=10$ Hz), 6.90 (d, 2H), 6.74 (dd, 1H, $J=9.0$, 3.5 Hz), 4.73 (d, 2H, $J=4.5$ Hz), 3.80 (s, 3H).

Step C: 5-Fluoro-1,3-dihydro-1-[(4-methoxyphenyl)methyl]-2,1-benzisothiazol-4-amine 2,2-dioxide

15 A mixture of 5-fluoro-1,3-dihydro-1-[(4-methoxyphenyl)methyl]-4-nitro-2,1-benzisothiazole 2,2-dioxide (1.54 g, 4.37 mmol) and $\text{SnCl}_2 \cdot \text{H}_2\text{O}$ (4.93 g, 21.9 mmol) in ethyl acetate (200 mL) was heated at reflux for 3 h. Then the reaction mixture was cooled, diluted with ethyl acetate (200 mL) and solid Na_2CO_3 (15 g) was added. The resulting mixture was vigorously stirred for 12 h and filtered through a pad of diatomaceous earth (Celite[®]). The filtrate was concentrated and the residue was subjected to flash chromatography to give 1.35 g of the title compound of Step C as brown solid melting at $154-157^\circ\text{C}$. ^1H NMR (CDCl_3 , 400 MHz): δ 7.33 (d, 2H), 6.87 (d, 2H), 6.82 (dd, 1H, $J=11.3$ and 8.75 Hz), 5.91 (dd, 1H, $J=8.5$ and 3.5 Hz), 4.65 (s, 2H), 4.22 (s, 2H), 3.79 (s, 3H), 3.75 (br s, 2H).

Step D: 7-Chloro-5-fluoro-1,3-dihydro-1-[(4-methoxyphenyl)methyl]-2,1-benzisothiazol-4-amine 2,2-dioxide

35 A mixture of the title compound of Step C (0.387 g, 1.2 mmol) and *N*-chlorosuccinimide (0.168 g, 1.26 mmol) in dimethylformamide (6 mL) was heated at 80°C for 4 h. After completion of the reaction, the mixture was poured into a mixture of ether (25 mL) and water (10 mL). The organic phase was separated, dried (MgSO_4) and

the solvent was removed on a rotary evaporator. Flash chromatography of the resulting residue afforded 0.4 g of the title compound of Step D. ¹H NMR (CDCl₃, 400 MHz): δ 7.23 (d,2H), 7.03 (d,1H), 6.74 (d,2H), 4.93 (s,2H), 3.84 (s,2H), 3.75 (s,3H).

5 Step E: 2-[7-Chloro-5-fluoro-1,3-dihydro-1-[(4-methoxyphenyl)methyl]-2,1-benzisothiazol-4-yl]-4,5,6,7-tetrahydro-1*H*-isoindole-1,3(2*H*)-dione *S,S*-dioxide and 2-(7-chloro-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl)-4,5,6,7-tetrahydro-1*H*-isoindole-1,3(2*H*)-dione *S,S*-dioxide

A mixture of the title compound of Step D (0.43 g, 1.2 mmol) and 3,4,5,6-tetrahydrophthalic anhydride (0.183 g, 1.2 mmol) in acetic acid (10 mL) was heated at reflux for 36 h. The solvent was then removed under reduced pressure and flash chromatography using 1-chlorobutane:ether (9:1) as eluent gave two compounds. 2-[7-Chloro-5-fluoro-1,3-dihydro-1-[(4-methoxyphenyl)methyl]-2,1-benzisothiazol-4-yl)-4,5,6,7-tetrahydro-1*H*-isoindole-1,3(2*H*)-dione *S,S*-dioxide, a compound of the invention, was eluted first as a solid melting at 171-174°C. ¹H NMR (CDCl₃, 300 MHz): δ 7.22-7.27 (m,3H), 6.80 (dd,2H, J=8.63 and 2.06 Hz), 4.98 (s,2H), 3.99 (s,2H), 3.77 (s,3H), 2.41 (m,4H), 1.81 (m,4H). 2-(7-Chloro-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl)-4,5,6,7-tetrahydro-1*H*-isoindole-1,3(2*H*)-dione *S,S*-dioxide, a compound of the invention, was eluted second as a solid melting at 75-79°C. ¹H NMR (CDCl₃, 300 MHz): δ 7.21 (d,1H, 9.38 Hz), 7.13 (br s,1H), 4.38 (s,2H), 2.44 (m,4H), 1.84 (m,4H).

EXAMPLE 14

2-(1-Acetyl-7-chloro-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl)-4,5,6,7-tetrahydro-1*H*-isoindole-1,3(2*H*)-dione *S,S*-dioxide

25 To a solution of 2-(7-chloro-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl)-4,5,6,7-tetrahydro-1*H*-isoindole-1,3(2*H*)-dione *S,S*-dioxide (0.208 g, 0.56 mmol) and Et₃N (0.5 mL) in anhydrous tetrahydrofuran (5 mL) was added acetyl chloride (50 mg, 0.044 mmol) dropwise at 0°C. The reaction mixture was then stirred at room temperature for 4 h. After the reaction was complete (TLC), the solvent was removed and the title compound of Example 14, a compound of the invention, was isolated by flash chromatography as a solid melting at 205-208°C. ¹H NMR (CDCl₃, 300 MHz): δ 7.37 (d,1H, J=9.57 Hz), 4.38 (s,2H), 2.59 (s,3H), 2.44 (m,4H) and 1.84 (m,4H).

EXAMPLE 15

35 Step A: 7-Chloro-5-fluoro-1,3-dihydro-4-isocyanato-1-[(4-methoxyphenyl)methyl]-2,1-benzisothiazole 2,2-dioxide

Following the procedures described in Step A in Example 4 using the title compound of Step D in Example 13 (0.68 g, 2 mmol), diphosgene (2.2 mmol) and triethylamine (3 mmol) afforded the title compound of Step A.

Step B: 3-[7-Chloro-5-fluoro-1,3-dihydro-1-[(4-methoxyphenyl)methyl]-2,1-benzisothiazol-4-yl]-6-(trifluoromethyl)-2,4(1*H*,3*H*)-pyrimidinedione *S,S*-dioxide

5 A solution of ethyl 3-amino-4,4,4-trifluorocrotonate (0.627 g, 3.42 mmol) in DMF (5 mL) was added to sodium hydride (137 mg, 60% suspension in oil) at -10 to -5°C. The reaction mixture was stirred initially at -10 to -5°C for 30 min, and then cooled to -78°C. The reaction mixture was stirred for 2 h and warmed to room temperature with continued stirring for 12 h. The reaction mixture was then poured into ice cold aqueous
10 HCl (10 mL of conc. HCl and 18 g of ice) and extracted with ethyl acetate (25 mL). The organic extract was dried (MgSO₄) and concentrated. The title compound of Step B, a compound of the invention, was isolated by flash chromatography as a solid melting at >230°C. ¹H NMR (Me₂SO-*d*₆, 400 MHz): δ 12.75 (br s, 1H), 7.80 (d, 1H), 7.27 (d, 2H), 6.88 (d, 2H), 6.45 (s, 1H), 4.89 (s, 2H), 4.86 (d, 1H), 4.54 (d, 1H), 3.72 (s, 3H).

15

EXAMPLE 16

3-[7-Chloro-5-fluoro-1,3-dihydro-1-[(4-methoxyphenyl)methyl]-2,1-benzisothiazol-4-yl]-1-methyl-6-(trifluoromethyl)-2,4(1*H*,3*H*)-pyrimidinedione *S,S*-dioxide

A mixture of the title compound of Step B in Example 15 (0.948 g, 1.82 mmol), dimethyl sulfate (0.24 g, 1.91 mmol) and K₂CO₃ (0.504 g, 3.65 mmol) in acetone
20 (10 mL) was heated at reflux for 3 h. After the reaction was complete (TLC), the reaction mixture was cooled, insolubles were filtered off and the filtrate was concentrated on a rotary evaporator to give a solid. The title compound of Example 16, a compound of the invention, was isolated by flash chromatography as a solid melting at 180-182°C. ¹H NMR (CDCl₃, 300 MHz): δ 7.27 (m, 3H), 6.81 (m, 2H), 6.31 (s, 1H),
25 4.98 (s, 2H), 3.94 (d, 2H), 3.77 (s, 3H), 3.54 (s, 3H).

EXAMPLE 17

3-(7-Chloro-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl)-1-methyl-6-(trifluoromethyl)-2,4(1*H*,3*H*)-pyrimidinedione *S,S*-dioxide

To a solution of the title compound of Example 16 (0.12 g, 0.225 mmol) in ethyl
30 acetate (3 mL) was added conc. H₂SO₄ (3 mL) at -10°C. The reaction mixture was stirred at 0°C for 1 h. After the reaction was complete (TLC), the reaction mixture was poured into a mixture of ice (60 g) and ethyl acetate (400 mL). The organic phase was separated, washed with water (2 x 50 mL), dried (MgSO₄) and concentrated. The title compound of Example 17, a compound of the invention, was isolated by flash
35 chromatography as a solid melting at >230°C. ¹H NMR (Me₂SO-*d*₆, 400 MHz): δ 11.18 (bs, 1H), 7.68 (d, 1H), 6.62 (s, 1H), 4.58 (q, 2H), 3.42 (s, 3H).

EXAMPLE 181-Acetyl-7-chloro-4-[3,6-dihydro-3-methyl-2,6-dioxo-4-(trifluoromethyl)-1(2H)-pyrimidinyl]-5-fluoro-1,3-dihydro-2,1-benzisothiazole 2,2-dioxide

To a solution of the title compound of Example 17 (0.1 g, 0.253 mmol),
5 triethylamine (38.5 mg, 0.38 mmol) and pyridine (0.1 g) in tetrahydrofuran (4 mL) was
added acetyl chloride (0.022 mL, 0.304 mmol) and the reaction mixture was stirred
initially at 0°C and then at room temperature for 1 h. After the reaction was complete,
the solvents were removed and the title compound of Example 18, a compound of the
invention, was isolated by flash chromatography as a colorless solid melting at
10 213-215°C. ¹H NMR (CDCl₃, 300 MHz): δ 7.40 (d,1H), 6.37 (s,1H), 4.33 (s,2H), 3.57
(s,3H), 2.59 (s,3H).

EXAMPLE 19Step A: 4-Fluoro-1-nitro-2-(phenylmethoxy)benzene

A mixture of 5-fluoro-2-nitrophenol (28 g, 0.178 mol), benzyl bromide (33.53 g,
15 0.195 mol) and K₂CO₃ (30.8 g, 0.223 mol) in anhydrous DMF (150 mL) was stirred at
room temperature for 12 h. After the reaction was complete, the reaction mixture was
poured into a mixture of ether (1000 mL) and water (200 mL). The organic phase was
separated, washed with water (3 x 100 mL), dried (MgSO₄) and concentrated to give the
title compound of Step A (40.5 g).

20 Step B: 4-Fluoro-2-(phenylmethoxy)benzenamine

To a mixture of iron powder (90.36 g, 1.62 mol) and acetic acid (10%, 400 mL) at
80°C was added a solution of the title compound of Step A (40 g, 0.162 mol) in ethyl
acetate (200 mL) dropwise with vigorous stirring. The reaction mixture was stirred
further at 76°C for 1 h. After the reaction was complete, the hot reaction mixture was
25 filtered through Celite® (200 g) and the filter cake was washed with ethyl acetate
(1000 mL). The filtrate was washed with water (3 x 200 mL), sodium bicarbonate
solution (5%, 3x 200 mL), dried (MgSO₄) and concentrated to give the crude product.
The product was purified by flash chromatography to give 24.76 g of the title compound
of Step B. ¹H NMR (CDCl₃, 400 MHz): δ 7.41 (m,5H), 6.60 (m,3H), 5.04 (s,2H).

30 Step C: 1-Chloro-N-(4-fluoro-2-hydroxyphenyl)methanesulfonamide

To a mixture of the title compound of Step B (13 g, 59.8 mmol) in pyridine
(60 mL) at 0°C was added chloromethylsulfonyl chloride (8.92 g, 59.8 mmol) dropwise
at 0°C under a nitrogen atmosphere. The reaction mixture was then stirred at 0°C for
1.5 h and at room temperature for 6 h. After the reaction was complete (TLC), the
35 reaction mixture was poured into ethyl acetate (750 mL) containing dilute HCl (10%,
150 mL). The organic phase was separated, washed successively with water
(2 x 100 mL), dilute HCl (2 x 100 mL) and water (2 x 100 mL). After drying (MgSO₄),
the organic layer was concentrated under reduced pressure to give 10.8 g of a solid

melting at 87-90°C. A suspension of this solid (10.5 g, 31.8 mmol) and Pd-C (10%, 1.0 g) in THF (60 mL) was hydrogenated using a Parr-shaker at 40 psi for 6 h. After the reaction was complete, the catalyst was filtered and the filtrate was concentrated to give 7.6 g of the title compound of Step C as a solid melting at 141-143°C. ¹H NMR (CDCl₃, 300 MHz): δ 7.28 (dd,1H), 6.6 (dd,1H), 6.64 (m,1H), 4.52 (s,2H).

5

Step D: 6-Fluoro-1*H*-4,2,1-benzoxathiazine 2,2-dioxide

A mixture of the title compound of Step C (0.985 g, 4.11 mmol) and K₂CO₃ (0.625 g, 4.52 mmol) in DMF (5 mL) was heated at 100°C for 4 h. After the starting material had been consumed (TLC), the reaction mixture was poured into an ethyl acetate (50 mL) and water (10 mL) mixture. The organic phase was washed with water (2 x 10 mL), dried (MgSO₄), and concentrated. The product was purified by flash chromatography to give 0.42 g of the title compound of Step D as yellow solid. ¹H NMR (CDCl₃, 300 MHz): δ 6.8 (m,3H), 6.2 (s,1H, NH), 4.95 (s,2H). IR (CDCl₃) 3347, 1506, 1148 cm⁻¹.

10

15

Step E: 6-Fluoro-1-(phenylmethyl)-1*H*-4,2,1-benzoxathiazine 2,2-dioxide

A mixture of the title compound of Step D (0.5 g, 2.46 mmol), K₂CO₃ (1.02 g, 7.38 mmol), benzyl bromide (0.463 g, 2.71 mmol) and TBAB (79 mg) in DMF (5 mL) was stirred at room temperature for 48 h. After the reaction was complete, the reaction mixture was poured into an H₂O/ethyl acetate mixture (10/100 mL). The organic phase was separated, dried and concentrated to give the title compound of Step E as a brown solid (0.76 g). ¹H NMR (CDCl₃, 300 MHz): δ 7.32 (m,6H, aromatic) 6.72 (d,1H, 11 Hz), 6.34 (d,1H, 8 Hz), 4.82 (s,2H), 4.69 (s,2H).

20

Step F: 6-Fluoro-7-nitro-1-(phenylmethyl)-1*H*-4,2,1-benzoxathiazine 2,2-dioxide and 6-Fluoro-7-nitro-1-[(4-nitrophenyl)methyl]-1*H*-4,2,1-benzoxathiazine 2,2-dioxide

25

To a solution of the title compound of Step E (0.76 g, 2.46 mmol) in acetic acid (20 mL) and H₂SO₄ (80%, 11.4 mL) was added nitric acid (2.9 mL, 70% HNO₃ + 0.3 mL H₂O) at 0°C and resulting reaction mixture was stirred initially at 0°C for 1 h and then at room temperature for 1.5 h. After the reaction was complete, the mixture was poured into ice water (50 mL) and extracted with ethyl acetate (100 mL). The organic extract was washed with water (2 x 10 mL), dried and concentrated. The two resulting products were separated by flash chromatography. The first compound eluted was 6-fluoro-7-nitro-1-(phenylmethyl)-1*H*-4,2,1-benzoxathiazine 2,2-dioxide (0.29 g) as a solid melting at 128-130°C. ¹H NMR (CDCl₃, 400 MHz): δ 7.80 (d,1H, J=7 Hz), 7.30-7.39 (m,5H), 6.93 (d,1H, J=10.5 Hz), 4.99 (s,2H), 4.88 (s,2H). The second compound eluted was 6-fluoro-7-nitro-1-[(4-nitrophenyl)methyl]-1*H*-4,2,1-benzoxathiazine 2,2-dioxide (0.296 gm) as a solid melting at 119-122°C.

30

35

Step G: 6-Fluoro-1-(phenylmethyl)-1*H*-4,2,1-benzoxathiazine-7-amine 2,2-dioxide

A mixture of 6-fluoro-7-nitro-1-(phenylmethyl)-1*H*-4,2,1-benzoxathiazine 2,2-dioxide (1.96 g, 5.79 mmol) and SnCl₂·2H₂O (6.55 g, 29 mmol) in ethyl acetate (40 mL) was heated at reflux for 3 h. After the reaction was complete, ethyl acetate (400 mL) and then Na₂CO₃ (18.5 g) were added, and the mixture was stirred for 12 h. The resulting mixture was filtered and the filtrate was concentrated under reduced pressure to give 1.20 g of the title compound of Step G.

Step H: 2-[6-Fluoro-1-(phenylmethyl)-1*H*-4,2,1-benzoxathiazin-7-yl]-4,5,6,7-tetrahydro-1*H*-isoindole-1,3(2*H*)-dione *S,S*-dioxide

A mixture of the title compound of Step G (1.2 g, 3.89 mmol), 3,4,5,6-tetrahydrophthalic anhydride (0.59 g) and acetic acid (6 mL) was heated at reflux for 48 h. After the reaction was complete, the reaction mixture was poured into an ethyl acetate (100 mL) and water (20 mL) mixture. The organic layer was separated, dried, and concentrated. Flash chromatography of the residue afforded 1.17 g of the title compound of Step H, a compound of the invention, as a colorless solid melting at 150-152°C. ¹H NMR (CDCl₃, 300 MHz): δ 7.33 (m,5H), 6.89 (m,2H), 4.92 (s,2H), 4.79 (s,2H), 2.41 (m,4H), 1.81 (m,4H).

By the procedures described herein, the following compounds of Tables 1-19 can be prepared. The following abbreviations are used: Ph = phenyl, *c* = cyclo, *t* = tertiary, Me = methyl. All alkyl groups are the normal isomers unless indicated otherwise.

Table 1

Compounds of Formula I wherein R¹=F, R²=Cl, R⁴=R⁵=H, J=J-1, Z=CH₂, R⁹=R¹⁰=H, n=1, m=2,

R ³	R ³	R ³	R ³	R ³
H	<i>c</i> -propyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ OCH ₂ CH ₂ OCH ₂
CH ₃	<i>c</i> -hexyl	CH ₃ OCH ₂ CH ₂	(4-MeO-Ph)CH ₂	Cl ₃ CCH ₂ OC(=O)
ethyl	FCH ₂ CH ₂ CH ₂	CH ₃ S(O) ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
propyl	CHF ₂	CH ₃ C(=O)	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ NHC(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂

Compounds of Formula II wherein R¹=F, R⁴=R⁵=H, J=J-1, Z=CH₂, R⁹=R¹⁰=H, n=1, m=2,

X=direct bond,

R ³	R ³	R ³	R ³	R ³
H	<i>c</i> -propyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ OCH ₂ CH ₂ OCH ₂
CH ₃	<i>c</i> -hexyl	CH ₃ OCH ₂ CH ₂	(4-MeO-Ph)CH ₂	Cl ₃ CCH ₂ OC(=O)
ethyl	FCH ₂ CH ₂ CH ₂	CH ₃ S(O) ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
propyl	CHF ₂	CH ₃ C(=O)	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ NHC(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂

<u>X=O,</u>				
R^3	R^3	R^3	R^3	R^3
H	<i>c</i> -propyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ OCH ₂ CH ₂ OCH ₂
CH ₃	<i>c</i> -hexyl	CH ₃ OCH ₂ CH ₂	(4-MeO-Ph)CH ₂	Cl ₃ CCH ₂ OC(=O)
ethyl	FCH ₂ CH ₂ CH ₂	CH ₃ S(O) ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
propyl	CHF ₂	CH ₃ C(=O)	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ NHC(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂
<u>X=S,</u>				
R^3	R^3	R^3	R^3	R^3
H	<i>c</i> -propyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ OCH ₂ CH ₂ OCH ₂
CH ₃	<i>c</i> -hexyl	CH ₃ OCH ₂ CH ₂	(4-MeO-Ph)CH ₂	Cl ₃ CCH ₂ OC(=O)
ethyl	FCH ₂ CH ₂ CH ₂	CH ₃ S(O) ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
propyl	CHF ₂	CH ₃ C(=O)	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ NHC(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂
<u>X=NH,</u>				
R^3	R^3	R^3	R^3	R^3
H	<i>c</i> -propyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ OCH ₂ CH ₂ OCH ₂
CH ₃	<i>c</i> -hexyl	CH ₃ OCH ₂ CH ₂	(4-MeO-Ph)CH ₂	Cl ₃ CCH ₂ OC(=O)
ethyl	FCH ₂ CH ₂ CH ₂	CH ₃ S(O) ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
propyl	CHF ₂	CH ₃ C(=O)	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ NHC(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂
<u>X=NCH₃,</u>				
R^3	R^3	R^3	R^3	R^3
H	<i>c</i> -propyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ OCH ₂ CH ₂ OCH ₂
CH ₃	<i>c</i> -hexyl	CH ₃ OCH ₂ CH ₂	(4-MeO-Ph)CH ₂	Cl ₃ CCH ₂ OC(=O)
ethyl	FCH ₂ CH ₂ CH ₂	CH ₃ S(O) ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
propyl	CHF ₂	CH ₃ C(=O)	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ NHC(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂

Table 2

Compounds of Formula I wherein R¹=F, R²=Cl, R⁴=R⁵=H, J=J-2, R⁹=R¹⁰=H,

Z=CH₂, m=1, n=2,

R^3	R^3	R^3	R^3	R^3
H	<i>c</i> -propyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ OCH ₂ CH ₂ OCH ₂
CH ₃	<i>c</i> -hexyl	CH ₃ OCH ₂ CH ₂	(4-MeO-Ph)CH ₂	Cl ₃ CCH ₂ OC(=O)
ethyl	FCH ₂ CH ₂ CH ₂	CH ₃ S(O) ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
propyl	CHF ₂	CH ₃ C(=O)	propargyl	(CH ₃) ₃ COC(=O)

octyl	allyl	$\text{CH}_3\text{NHC(=O)}$	$\text{CH}_3\text{NHS(O)}_2$	$\text{CH}_2=\text{CHCH}_2\text{CH}_2\text{CH}_2$
<u>Z=CHF, n=1, m=1,</u>				
R^3	R^3	R^3	R^3	R^3
H	<i>c</i> -propyl	CH_3OCH_2	$\text{PhCH}_2\text{OC(=O)}$	$\text{CH}_3\text{OCH}_2\text{CH}_2\text{OCH}_2$
CH_3	<i>c</i> -hexyl	$\text{CH}_3\text{OCH}_2\text{CH}_2$	(4-MeO-Ph) CH_2	$\text{Cl}_3\text{CCH}_2\text{OC(=O)}$
ethyl	$\text{FCH}_2\text{CH}_2\text{CH}_2$	$\text{CH}_3\text{S(O)}_2$	PhCH_2	$\text{ClCH}_2\text{CH}_2\text{CH}_2$
propyl	CHF_2	$\text{CH}_3\text{C(=O)}$	propargyl	$(\text{CH}_3)_3\text{COC(=O)}$
octyl	allyl	$\text{CH}_3\text{NHC(=O)}$	$\text{CH}_3\text{NHS(O)}_2$	$\text{CH}_2=\text{CHCH}_2\text{CH}_2\text{CH}_2$
<u>Z=O, n=2, m=1,</u>				
R^3	R^3	R^3	R^3	R^3
H	<i>c</i> -propyl	CH_3OCH_2	$\text{PhCH}_2\text{OC(=O)}$	$\text{CH}_3\text{OCH}_2\text{CH}_2\text{OCH}_2$
CH_3	<i>c</i> -hexyl	$\text{CH}_3\text{OCH}_2\text{CH}_2$	(4-MeO-Ph) CH_2	$\text{Cl}_3\text{CCH}_2\text{OC(=O)}$
ethyl	$\text{FCH}_2\text{CH}_2\text{CH}_2$	$\text{CH}_3\text{S(O)}_2$	PhCH_2	$\text{ClCH}_2\text{CH}_2\text{CH}_2$
propyl	CHF_2	$\text{CH}_3\text{C(=O)}$	propargyl	$(\text{CH}_3)_3\text{COC(=O)}$
octyl	allyl	$\text{CH}_3\text{NHC(=O)}$	$\text{CH}_3\text{NHS(O)}_2$	$\text{CH}_2=\text{CHCH}_2\text{CH}_2\text{CH}_2$

Compounds of Formula II wherein $\text{R}^1=\text{F}$, $\text{R}^4=\text{R}^5=\text{H}$, $\text{J}=\text{J}-2$, $\text{Z}=\text{CH}_2$, $\text{R}^9=\text{R}^{10}=\text{H}$, $n=1$, $m=2$,

$\text{Z}=\text{CH}_2$.

X=direct bond,

R^3	R^3	R^3	R^3	R^3
H	<i>c</i> -propyl	CH_3OCH_2	$\text{PhCH}_2\text{OC(=O)}$	$\text{CH}_3\text{OCH}_2\text{CH}_2\text{OCH}_2$
CH_3	<i>c</i> -hexyl	$\text{CH}_3\text{OCH}_2\text{CH}_2$	(4-MeO-Ph) CH_2	$\text{Cl}_3\text{CCH}_2\text{OC(=O)}$
ethyl	$\text{FCH}_2\text{CH}_2\text{CH}_2$	$\text{CH}_3\text{S(O)}_2$	PhCH_2	$\text{ClCH}_2\text{CH}_2\text{CH}_2$
propyl	CHF_2	$\text{CH}_3\text{C(=O)}$	propargyl	$(\text{CH}_3)_3\text{COC(=O)}$
octyl	allyl	$\text{CH}_3\text{NHC(=O)}$	$\text{CH}_3\text{NHS(O)}_2$	$\text{CH}_2=\text{CHCH}_2\text{CH}_2\text{CH}_2$

X=O,

R^3	R^3	R^3	R^3	R^3
H	<i>c</i> -propyl	CH_3OCH_2	$\text{PhCH}_2\text{OC(=O)}$	$\text{CH}_3\text{OCH}_2\text{CH}_2\text{OCH}_2$
CH_3	<i>c</i> -hexyl	$\text{CH}_3\text{OCH}_2\text{CH}_2$	(4-MeO-Ph) CH_2	$\text{Cl}_3\text{CCH}_2\text{OC(=O)}$
ethyl	$\text{FCH}_2\text{CH}_2\text{CH}_2$	$\text{CH}_3\text{S(O)}_2$	PhCH_2	$\text{ClCH}_2\text{CH}_2\text{CH}_2$
propyl	CHF_2	$\text{CH}_3\text{C(=O)}$	propargyl	$(\text{CH}_3)_3\text{COC(=O)}$
octyl	allyl	$\text{CH}_3\text{NHC(=O)}$	$\text{CH}_3\text{NHS(O)}_2$	$\text{CH}_2=\text{CHCH}_2\text{CH}_2\text{CH}_2$

X=S,

R^3	R^3	R^3	R^3	R^3
H	<i>c</i> -propyl	CH_3OCH_2	$\text{PhCH}_2\text{OC(=O)}$	$\text{CH}_3\text{OCH}_2\text{CH}_2\text{OCH}_2$
CH_3	<i>c</i> -hexyl	$\text{CH}_3\text{OCH}_2\text{CH}_2$	(4-MeO-Ph) CH_2	$\text{Cl}_3\text{CCH}_2\text{OC(=O)}$
ethyl	$\text{FCH}_2\text{CH}_2\text{CH}_2$	$\text{CH}_3\text{S(O)}_2$	PhCH_2	$\text{ClCH}_2\text{CH}_2\text{CH}_2$
propyl	CHF_2	$\text{CH}_3\text{C(=O)}$	propargyl	$(\text{CH}_3)_3\text{COC(=O)}$

octyl	allyl	$\text{CH}_3\text{NHC}(=\text{O})$	$\text{CH}_3\text{NHS}(\text{O})_2$	$\text{CH}_2=\text{CHCH}_2\text{CH}_2\text{CH}_2$
$\text{X}=\text{NH}$,				
R^3	R^3	R^3	R^3	R^3
H	c-propyl	CH_3OCH_2	$\text{PhCH}_2\text{OC}(=\text{O})$	$\text{CH}_3\text{OCH}_2\text{CH}_2\text{OCH}_2$
CH_3	c-hexyl	$\text{CH}_3\text{OCH}_2\text{CH}_2$	(4-MeO-Ph) CH_2	$\text{Cl}_3\text{CCH}_2\text{OC}(=\text{O})$
ethyl	$\text{FCH}_2\text{CH}_2\text{CH}_2$	$\text{CH}_3\text{S}(\text{O})_2$	PhCH_2	$\text{ClCH}_2\text{CH}_2\text{CH}_2$
propyl	CHF_2	$\text{CH}_3\text{C}(=\text{O})$	propargyl	$(\text{CH}_3)_3\text{COC}(=\text{O})$
octyl	allyl	$\text{CH}_3\text{NHC}(=\text{O})$	$\text{CH}_3\text{NHS}(\text{O})_2$	$\text{CH}_2=\text{CHCH}_2\text{CH}_2\text{CH}_2$
$\text{X}=\text{NCH}_3$,				
R^3	R^3	R^3	R^3	R^3
H	c-propyl	CH_3OCH_2	$\text{PhCH}_2\text{OC}(=\text{O})$	$\text{CH}_3\text{OCH}_2\text{CH}_2\text{OCH}_2$
CH_3	c-hexyl	$\text{CH}_3\text{OCH}_2\text{CH}_2$	(4-MeO-Ph) CH_2	$\text{Cl}_3\text{CCH}_2\text{OC}(=\text{O})$
ethyl	$\text{FCH}_2\text{CH}_2\text{CH}_2$	$\text{CH}_3\text{S}(\text{O})_2$	PhCH_2	$\text{ClCH}_2\text{CH}_2\text{CH}_2$
propyl	CHF_2	$\text{CH}_3\text{C}(=\text{O})$	propargyl	$(\text{CH}_3)_3\text{COC}(=\text{O})$
octyl	allyl	$\text{CH}_3\text{NHC}(=\text{O})$	$\text{CH}_3\text{NHS}(\text{O})_2$	$\text{CH}_2=\text{CHCH}_2\text{CH}_2\text{CH}_2$

Table 3

Compounds of Formula I wherein $\text{J}=\text{J}-3$, $\text{R}^1=\text{F}$, $\text{R}^2=\text{Cl}$, $\text{R}^4=\text{R}^5=\text{H}$, $\text{R}^{11}=\text{CF}_2\text{H}$, $\text{R}^{12}=\text{CH}_3$,

R^3	R^3	R^3	R^3	R^3
H	c-propyl	CH_3OCH_2	$\text{PhCH}_2\text{OC}(=\text{O})$	$\text{CH}_3\text{OCH}_2\text{CH}_2\text{OCH}_2$
CH_3	c-hexyl	$\text{CH}_3\text{OCH}_2\text{CH}_2$	(4-MeO-Ph) CH_2	$\text{Cl}_3\text{CCH}_2\text{OC}(=\text{O})$
ethyl	$\text{FCH}_2\text{CH}_2\text{CH}_2$	$\text{CH}_3\text{S}(\text{O})_2$	PhCH_2	$\text{ClCH}_2\text{CH}_2\text{CH}_2$
propyl	CHF_2	$\text{CH}_3\text{C}(=\text{O})$	propargyl	$(\text{CH}_3)_3\text{COC}(=\text{O})$
octyl	allyl	$\text{CH}_3\text{NHC}(=\text{O})$	$\text{CH}_3\text{NHS}(\text{O})_2$	$\text{CH}_2=\text{CHCH}_2\text{CH}_2\text{CH}_2$

Compounds of Formula II wherein $\text{J}=\text{J}-3$, $\text{R}^1=\text{F}$, $\text{R}^4=\text{R}^5=\text{H}$, $\text{R}^{11}=\text{CF}_2\text{H}$, $\text{R}^{12}=\text{CH}_3$,

$\text{X}=\text{Direct Bond}$,

R^3	R^3	R^3	R^3	R^3
H	c-hexyl	CH_3OCH_2	$\text{PhCH}_2\text{OC}(=\text{O})$	$\text{CH}_3\text{NHC}(=\text{O})$
CH_3	$\text{FCH}_2\text{CH}_2\text{CH}_2$	$\text{CH}_3\text{OCH}_2\text{CH}_2$	PhCH_2	$\text{ClCH}_2\text{CH}_2\text{CH}_2$
ethyl	CHF_2	$\text{CH}_3\text{S}(\text{O})_2$	propargyl	$(\text{CH}_3)_3\text{COC}(=\text{O})$
octyl	allyl	$\text{CH}_3\text{C}(=\text{O})$	$\text{CH}_3\text{NHS}(\text{O})_2$	$\text{CH}_2=\text{CHCH}_2\text{CH}_2\text{CH}_2$
$\text{X}=\text{O}$,				
R^3	R^3	R^3	R^3	R^3
H	c-hexyl	CH_3OCH_2	$\text{PhCH}_2\text{OC}(=\text{O})$	$\text{CH}_3\text{NHC}(=\text{O})$
CH_3	$\text{FCH}_2\text{CH}_2\text{CH}_2$	$\text{CH}_3\text{OCH}_2\text{CH}_2$	PhCH_2	$\text{ClCH}_2\text{CH}_2\text{CH}_2$
ethyl	CHF_2	$\text{CH}_3\text{S}(\text{O})_2$	propargyl	$(\text{CH}_3)_3\text{COC}(=\text{O})$
octyl	allyl	$\text{CH}_3\text{C}(=\text{O})$	$\text{CH}_3\text{NHS}(\text{O})_2$	$\text{CH}_2=\text{CHCH}_2\text{CH}_2\text{CH}_2$

Table 4

Compounds of Formula I wherein J=J-4, R¹=F, R²=Cl, R⁴=R⁵=H, Q=O, R¹³=CH₃,W=CH, R¹⁴=CF₃,

R ³	R ³	R ³	R ³	R ³
H	c-hexyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
ethyl	CHF ₂	CH ₃ S(O) ₂	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ C(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂

W=N, R¹⁴=N(CH₃)₂,

R ³	R ³	R ³	R ³	R ³
H	c-hexyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
ethyl	CHF ₂	CH ₃ S(O) ₂	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ C(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂

Compounds of Formula II wherein J=J-4, R¹=F, R⁴=R⁵=H, Q=O, R¹³=CH₃,X=Direct Bond, W=CH, R¹⁴=CF₃,

R ³	R ³	R ³	R ³	R ³
H	c-hexyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
ethyl	CHF ₂	CH ₃ S(O) ₂	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ C(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂

X=O, W=N, R¹⁴=N(CH₃)₂,

R ³	R ³	R ³	R ³	R ³
H	c-hexyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
ethyl	CHF ₂	CH ₃ S(O) ₂	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ C(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂

Table 5

Compounds of Formula I wherein J=J-5, R¹=F, R²=Cl, R⁴=R⁵=H, Q=O, R⁹=R¹⁰=H, Z=CH₂, only single bonds in left-hand ring,n=m=1,

R ³	R ³	R ³	R ³	R ³
H	c-hexyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
ethyl	CHF ₂	CH ₃ S(O) ₂	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ C(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂

m=2, n=1,

R^3	R^3	R^3	R^3	R^3
H	c-hexyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
ethyl	CHF ₂	CH ₃ S(O) ₂	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ C(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂

Compounds of Formula II wherein J=J-5, R¹=F, R⁴=R⁵=H, Q=O, R⁹=R¹⁰=H, Z=CH₂, only single bonds in left-hand ring,

X=Direct Bond, n=m=1,

R^3	R^3	R^3	R^3	R^3
H	c-hexyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
ethyl	CHF ₂	CH ₃ S(O) ₂	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ C(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂

X=O, m=2, n=1,

R^3	R^3	R^3	R^3	R^3
H	c-hexyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
ethyl	CHF ₂	CH ₃ S(O) ₂	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ C(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂

Table 6

Compounds of Formula I wherein J=J-6, R¹=F, R²=Cl, R⁴=R⁵=H, Q=O, R¹⁶=R¹⁷=H, only single bonds in left-hand ring,

Z¹=CH₂, n=m=1,

R^3	R^3	R^3	R^3	R^3
H	c-hexyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
ethyl	CHF ₂	CH ₃ S(O) ₂	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ C(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂

Z¹=CHF, n=m=1,

R^3	R^3	R^3	R^3	R^3
H	c-hexyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
ethyl	CHF ₂	CH ₃ S(O) ₂	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ C(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂

$Z^1 = \text{CF}_2, n=m=1,$

R^3	R^3	R^3	R^3	R^3
H	<i>c</i> -hexyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
ethyl	CHF ₂	CH ₃ S(O) ₂	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ C(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂

 $Z^1 = \text{O}, n=2, m=1,$

R^3	R^3	R^3	R^3	R^3
H	<i>c</i> -hexyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
ethyl	CHF ₂	CH ₃ S(O) ₂	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ C(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂

Compounds of Formula II wherein J=J-6, R¹=F, R⁴=R⁵=H, Q=O, R¹⁶=R¹⁷=H, n=m=1, only single bonds in left-hand ring,

 $X = \text{Direct Bond}, Z^1 = \text{CH}_2,$

R^3	R^3	R^3	R^3	R^3
H	<i>c</i> -hexyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
ethyl	CHF ₂	CH ₃ S(O) ₂	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ C(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂

 $X = \text{NCH}_3, Z^1 = \text{CH}_2,$

R^3	R^3	R^3	R^3	R^3
H	<i>c</i> -hexyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
ethyl	CHF ₂	CH ₃ S(O) ₂	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ C(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂

 $X = \text{Direct Bond}, Z^1 = \text{CHF},$

R^3	R^3	R^3	R^3	R^3
H	<i>c</i> -hexyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
ethyl	CHF ₂	CH ₃ S(O) ₂	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ C(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂

 $X = \text{NCH}_3, Z^1 = \text{CHF},$

R^3	R^3	R^3	R^3	R^3
H	<i>c</i> -hexyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂

ethyl	CHF ₂	CH ₃ S(O) ₂	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ C(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂

Table 7

Compounds of Formula I wherein J=J-7, R¹=F, R²=Cl, R⁴=R⁵=H, Q=O, R¹²=H,

R ³	R ¹¹	R ³	R ¹¹	R ³	R ¹¹
H	H	FCH ₂ CH ₂ CH ₂	CH ₃	CHF ₂	CH ₃
CH ₃	H	CH ₃ OCH ₂ CH ₂	CH ₃	allyl	H
ethyl	CH ₃	CH ₃ S(O) ₂	FCH ₂ CH ₂ CH ₂	propargyl	FCH ₂ CH ₂ CH ₂
ethyl	FCH ₂ CH ₂ CH ₂	CH ₃ C(=O)	FCH ₂ CH ₂ CH ₂	PhCH ₂	CH ₃

Compounds of Formula II wherein J=J-7, R¹=F, R⁴=R⁵=H, Q=O, R¹²=H,X=Direct Bond.

R ³	R ¹¹	R ³	R ¹¹	R ³	R ¹¹
H	H	FCH ₂ CH ₂ CH ₂	CH ₃	CHF ₂	CH ₃
CH ₃	H	CH ₃ OCH ₂ CH ₂	CH ₃	allyl	H
ethyl	CH ₃	CH ₃ S(O) ₂	FCH ₂ CH ₂ CH ₂	propargyl	FCH ₂ CH ₂ CH ₂
ethyl	FCH ₂ CH ₂ CH ₂	CH ₃ C(=O)	FCH ₂ CH ₂ CH ₂	PhCH ₂	CH ₃

X=NCH₃.

R ³	R ¹¹	R ³	R ¹¹	R ³	R ¹¹
H	H	FCH ₂ CH ₂ CH ₂	CH ₃	CHF ₂	CH ₃
CH ₃	H	CH ₃ OCH ₂ CH ₂	CH ₃	allyl	H
ethyl	CH ₃	CH ₃ S(O) ₂	FCH ₂ CH ₂ CH ₂	propargyl	FCH ₂ CH ₂ CH ₂
ethyl	FCH ₂ CH ₂ CH ₂	CH ₃ C(=O)	FCH ₂ CH ₂ CH ₂	PhCH ₂	CH ₃

Table 8

Compounds of Formula I wherein J=J-8, R¹=F, R²=Cl, R⁴=R⁵=H, R¹⁸=*t*-butyl,

R ³	R ³	R ³	R ³	R ³
H	<i>c</i> -hexyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
ethyl	CHF ₂	CH ₃ S(O) ₂	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ C(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂

Compounds of Formula II wherein J=J-8, R¹=F, R⁴=R⁵=H, R¹⁸=*t*-butyl, X=O,

R ³	R ³	R ³	R ³	R ³
H	<i>c</i> -hexyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
ethyl	CHF ₂	CH ₃ S(O) ₂	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ C(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂

Table 9

Compounds of Formula I wherein J=J-9, R¹=F, R²=Cl, R⁴=R⁵=H, Q=O, R¹⁸=CH₂CH₂CH₂F,

R ³	R ³	R ³	R ³	R ³
H	c-hexyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
ethyl	CHF ₂	CH ₃ S(O) ₂	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ C(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂

Compounds of Formula II wherein J=J-9, R¹=F, R⁴=R⁵=H, Q=O, R¹⁸=CH₂CH₂CH₂F, X=O,

R ³	R ³	R ³	R ³	R ³
H	c-hexyl	CH ₃ OCH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂	ClCH ₂ CH ₂ CH ₂
ethyl	CHF ₂	CH ₃ S(O) ₂	propargyl	(CH ₃) ₃ COC(=O)
octyl	allyl	CH ₃ C(=O)	CH ₃ NHS(O) ₂	CH ₂ =CHCH ₂ CH ₂ CH ₂

Table 10

Compounds of Formula I wherein J=J-10, R¹=F, R²=Cl, R⁴=R⁵=H, Q=O, R⁹=R¹⁰=H, Z=CH₂, m=n=1, R¹⁹=R²⁰=H,

R ³	R ³	R ³	R ³	R ³
H	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	CHF ₂	CH ₃ S(O) ₂	PhCH ₂	octyl
ethyl	allyl	CH ₃ C(=O)	propargyl	(CH ₃) ₃ COC(=O)

Compounds of Formula II wherein J=J-10, R¹=F, R⁴=R⁵=H, Q=O, R⁹=R¹⁰=H, Z=CH₂, m=n=1, R¹⁹=R²⁰=H, X=O,

R ³	R ³	R ³	R ³	R ³
H	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	CHF ₂	CH ₃ S(O) ₂	PhCH ₂	octyl
ethyl	allyl	CH ₃ C(=O)	propargyl	(CH ₃) ₃ COC(=O)

Table 11

Compounds of Formula I wherein J=J-12, R¹=F, R²=Cl, R⁴=R⁵=H, Q=O, Q¹=S, R⁹=R¹⁰=H, Z=CH₂, m=n=1, only single bonds in left-hand ring,

R ³	R ³	R ³	R ³	R ³
H	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	CHF ₂	CH ₃ S(O) ₂	PhCH ₂	octyl
ethyl	allyl	CH ₃ C(=O)	propargyl	(CH ₃) ₃ COC(=O)

Compounds of Formula II wherein J=J-12, R¹=F, R⁴=R⁵=H, Q=O, Q¹=S, R⁹=R¹⁰=H, X=O, m=n=1, only single bonds in left-hand ring,

<u>Z=CH₂,</u>				
<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>
H	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	CHF ₂	CH ₃ S(O) ₂	PhCH ₂	octyl
ethyl	allyl	CH ₃ C(=O)	propargyl	(CH ₃) ₃ COC(=O)
<u>Z=CHF,</u>				
<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>
H	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	CHF ₂	CH ₃ S(O) ₂	PhCH ₂	octyl
ethyl	allyl	CH ₃ C(=O)	propargyl	(CH ₃) ₃ COC(=O)

Table 12

Compounds of Formula I wherein J=J-14, R¹=F, R²=Cl, R⁴=R⁵=H, Q¹=S, R⁹=R¹⁰=H, Z=CH₂, m=2, n=1,

<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>
H	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	CHF ₂	CH ₃ S(O) ₂	PhCH ₂	octyl
ethyl	allyl	CH ₃ C(=O)	propargyl	(CH ₃) ₃ COC(=O)

Compounds of Formula II wherein J=J-14, R¹=F, R⁴=R⁵=H, Q¹=S, R⁹=R¹⁰=H, Z=CH₂, m=2, n=1, X=O,

<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>
H	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	CHF ₂	CH ₃ S(O) ₂	PhCH ₂	octyl
ethyl	allyl	CH ₃ C(=O)	propargyl	(CH ₃) ₃ COC(=O)

Table 13

Compounds of Formula I wherein J=J-15, R¹=F, R²=Cl, R⁴=R⁵=H, R⁹=R¹⁰=H, Z=CH₂, m=2, n=1, R²³=Cl,

<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>
H	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	CHF ₂	CH ₃ S(O) ₂	PhCH ₂	octyl
ethyl	allyl	CH ₃ C(=O)	propargyl	(CH ₃) ₃ COC(=O)

Compounds of Formula II wherein J=J-15, R¹=F, R⁴=R⁵=H, R⁹=R¹⁰=H, Z=CH₂, m=2, n=1, R²³=Cl,

X=Direct Bond,

<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>
H	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)

CH ₃	CHF ₂	CH ₃ S(O) ₂	PhCH ₂	octyl
ethyl	allyl	CH ₃ C(=O)	propargyl	(CH ₃) ₃ COC(=O)
<u>X=O,</u>				
<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>
H	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	CHF ₂	CH ₃ S(O) ₂	PhCH ₂	octyl
ethyl	allyl	CH ₃ C(=O)	propargyl	(CH ₃) ₃ COC(=O)

Table 14

Compounds of Formula I wherein J=J-16, R¹=F, R²=Cl, R⁴=R⁵=H, R²³=Cl, R²⁴=CF₃, R²⁵=CH₃,

<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>
H	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	CHF ₂	CH ₃ S(O) ₂	PhCH ₂	octyl
ethyl	allyl	CH ₃ C(=O)	propargyl	(CH ₃) ₃ COC(=O)

Compounds of Formula II wherein J=J-16, R¹=F, R⁴=R⁵=H, R²³=Cl, R²⁴=CF₃, R²⁵=CH₃,

X=Direct Bond,

<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>
H	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	CHF ₂	CH ₃ S(O) ₂	PhCH ₂	octyl
ethyl	allyl	CH ₃ C(=O)	propargyl	(CH ₃) ₃ COC(=O)

X=O,

<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>
H	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	CHF ₂	CH ₃ S(O) ₂	PhCH ₂	octyl
ethyl	allyl	CH ₃ C(=O)	propargyl	(CH ₃) ₃ COC(=O)

Table 15

Compounds of Formula I wherein J=J-17, R¹=F, R²=Cl, R⁴=R⁵=H, R⁹=R¹⁰=H, m=n=1, R²⁶=CH₃,

Z=CH₂,

<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>
H	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	CHF ₂	CH ₃ S(O) ₂	PhCH ₂	octyl
ethyl	allyl	CH ₃ C(=O)	propargyl	(CH ₃) ₃ COC(=O)

Z=CHF,

<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>	<u>R³</u>
H	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	CHF ₂	CH ₃ S(O) ₂	PhCH ₂	octyl

ethyl allyl $\text{CH}_3\text{C}(=\text{O})$ propargyl $(\text{CH}_3)_3\text{COC}(=\text{O})$
 Compounds of Formula II wherein $\text{J}=\text{J}-17$, $\text{R}^1=\text{F}$, $\text{R}^4=\text{R}^5=\text{H}$, $\text{R}^9=\text{R}^{10}=\text{H}$, $m=n=1$, $\text{R}^{26}=\text{CH}_3$,

$\text{X}=\text{Direct Bond}$, $\text{Z}=\text{CH}_2$,

R^3	R^3	R^3	R^3	R^3
H	$\text{FCH}_2\text{CH}_2\text{CH}_2$	$\text{CH}_3\text{OCH}_2\text{CH}_2$	$\text{PhCH}_2\text{OC}(=\text{O})$	$\text{CH}_3\text{NHC}(=\text{O})$
CH_3	CHF_2	$\text{CH}_3\text{S}(\text{O})_2$	PhCH_2	octyl
ethyl	allyl	$\text{CH}_3\text{C}(=\text{O})$	propargyl	$(\text{CH}_3)_3\text{COC}(=\text{O})$

$\text{X}=\text{O}$, $\text{Z}=\text{CH}_2$,

R^3	R^3	R^3	R^3	R^3
H	$\text{FCH}_2\text{CH}_2\text{CH}_2$	$\text{CH}_3\text{OCH}_2\text{CH}_2$	$\text{PhCH}_2\text{OC}(=\text{O})$	$\text{CH}_3\text{NHC}(=\text{O})$
CH_3	CHF_2	$\text{CH}_3\text{S}(\text{O})_2$	PhCH_2	octyl
ethyl	allyl	$\text{CH}_3\text{C}(=\text{O})$	propargyl	$(\text{CH}_3)_3\text{COC}(=\text{O})$

$\text{X}=\text{O}$, $\text{Z}=\text{CHF}$,

R^3	R^3	R^3	R^3	R^3
H	$\text{FCH}_2\text{CH}_2\text{CH}_2$	$\text{CH}_3\text{OCH}_2\text{CH}_2$	$\text{PhCH}_2\text{OC}(=\text{O})$	$\text{CH}_3\text{NHC}(=\text{O})$
CH_3	CHF_2	$\text{CH}_3\text{S}(\text{O})_2$	PhCH_2	octyl
ethyl	allyl	$\text{CH}_3\text{C}(=\text{O})$	propargyl	$(\text{CH}_3)_3\text{COC}(=\text{O})$

Table 16

Compounds of Formula I wherein $\text{J}=\text{J}-18$, $\text{R}^1=\text{F}$, $\text{R}^2=\text{Cl}$, $\text{R}^4=\text{R}^5=\text{H}$, $\text{R}^9=\text{R}^{10}=\text{H}$, $m=2$, $n=1$, $\text{R}^{26}=\text{CH}_3$,

$\text{Z}=\text{CH}_2$,

R^3	R^3	R^3	R^3	R^3
H	$\text{FCH}_2\text{CH}_2\text{CH}_2$	$\text{CH}_3\text{OCH}_2\text{CH}_2$	$\text{PhCH}_2\text{OC}(=\text{O})$	$\text{CH}_3\text{NHC}(=\text{O})$
CH_3	CHF_2	$\text{CH}_3\text{S}(\text{O})_2$	PhCH_2	octyl
ethyl	allyl	$\text{CH}_3\text{C}(=\text{O})$	propargyl	$(\text{CH}_3)_3\text{COC}(=\text{O})$

$\text{Z}=\text{CHE}$,

R^3	R^3	R^3	R^3	R^3
H	$\text{FCH}_2\text{CH}_2\text{CH}_2$	$\text{CH}_3\text{OCH}_2\text{CH}_2$	$\text{PhCH}_2\text{OC}(=\text{O})$	$\text{CH}_3\text{NHC}(=\text{O})$
CH_3	CHF_2	$\text{CH}_3\text{S}(\text{O})_2$	PhCH_2	octyl
ethyl	allyl	$\text{CH}_3\text{C}(=\text{O})$	propargyl	$(\text{CH}_3)_3\text{COC}(=\text{O})$

Compounds of Formula II wherein $\text{J}=\text{J}-18$, $\text{R}^1=\text{F}$, $\text{R}^4=\text{R}^5=\text{H}$, $\text{R}^9=\text{R}^{10}=\text{H}$, $m=n=1$, $\text{R}^{26}=\text{CH}_3$,

$\text{X}=\text{O}$, $\text{Z}=\text{CH}_2$,

R^3	R^3	R^3	R^3	R^3
H	$\text{FCH}_2\text{CH}_2\text{CH}_2$	$\text{CH}_3\text{OCH}_2\text{CH}_2$	$\text{PhCH}_2\text{OC}(=\text{O})$	$\text{CH}_3\text{NHC}(=\text{O})$
CH_3	CHF_2	$\text{CH}_3\text{S}(\text{O})_2$	PhCH_2	octyl
ethyl	allyl	$\text{CH}_3\text{C}(=\text{O})$	propargyl	$(\text{CH}_3)_3\text{COC}(=\text{O})$

X=O, Z=CHF.

R^3	R^3	R^3	R^3	R^3
H	FCH ₂ CH ₂ CH ₂	CH ₃ OCH ₂ CH ₂	PhCH ₂ OC(=O)	CH ₃ NHC(=O)
CH ₃	CHF ₂	CH ₃ S(O) ₂	PhCH ₂	octyl
ethyl	allyl	CH ₃ C(=O)	propargyl	(CH ₃) ₃ COC(=O)

Table 17

Compounds of Formula I wherein J=J-1, R³=ethyl, R⁴=R⁵=H, R⁹=R¹⁰=H, m=2, n=1, Z=CH₂,

R^1	R^2	R^1	R^2	R^1	R^2	R^1	R^2	R^1	R^2
H	Cl	Cl	F	Cl	Cl	F	Br	F	NO ₂
H	F	Br	Cl	Br	F	F	CF ₃	F	CN

Compounds of Formula II wherein J=J-1, R³=ethyl, R⁴=R⁵=H, R⁹=R¹⁰=H, m=2, n=1, Z=CH₂,

R^1	X	R^1	X	R^1	X	R^1	X	R^1	X
H	O	Cl	O	Cl	S	F	O	F	direct bond
H	S	Br	O	Br	S	F	S	F	NCH ₃

Table 18

Compounds of Formula I wherein J=J-1, R¹=F, R²=Cl, R³=ethyl, R⁹=R¹⁰=H, m=2, n=1, Z=CH₂,


R^4	R^5	R^4	R^5	R^4	R^5	R^4	R^5	R^4	R^5
Cl	H	Br	H	CH ₃	CH ₃	ethyl	H	H	C(=O)CH ₃
F	H	CH ₃	H	H	CF ₃	H	S(O) ₂ CH ₃	spiro-cyclopropyl	

Compounds of Formula II wherein J=J-1, R¹=F, R³=ethyl, R⁹=R¹⁰=H, m=2, n=1, Z=CH₂, X=O,

R^4	R^5	R^4	R^5	R^4	R^5	R^4	R^5	R^4	R^5
Cl	H	Br	H	CH ₃	CH ₃	ethyl	H	H	C(=O)CH ₃
F	H	CH ₃	H	H	CF ₃	H	S(O) ₂ CH ₃	spiro-cyclopropyl	

Table 19

R³=ethyl, R¹=F, R⁴=R⁵=H,



Formula	R^2	Formula	X
I	Cl	II	Direct bond
I	F	II	NCH ₃
I	NO ₂	II	NH
I	CF ₃	II	NCH ₂ CH ₃

Formulation/Utility

Compounds of this invention will generally be used in formulation with an agriculturally suitable carrier comprising at least one of a liquid diluent, a solid diluent or a surfactant wherein the formulation is consistent with the physical properties of the active ingredient, mode of application and environmental factors such as soil type, moisture and temperature. Useful formulations include liquids such as solutions (including emulsifiable concentrates), suspensions, emulsions (including microemulsions and/or suspoemulsions) and the like which optionally can be thickened into gels. Useful formulations further include solids such as dusts, powders, granules, pellets, tablets, films, and the like which can be water-dispersible ("wettable") or water-soluble. Active ingredient can be (micro)encapsulated and further formed into a suspension or solid formulation; alternatively the entire formulation of active ingredient can be encapsulated (or "overcoated"). Encapsulation can control or delay release of the active ingredient. Sprayable formulations can be extended in suitable media and used at spray volumes from about one to several hundred liters per hectare. High-strength compositions are primarily used as intermediates for further formulation.

The formulations will typically contain effective amounts of active ingredient, diluent and surfactant within the following approximate ranges which add up to 100 percent by weight.

	Weight Percent		
	<u>Active Ingredient</u>	<u>Diluent</u>	<u>Surfactant</u>
Water-Dispersible and Water-soluble Granules, Tablets and Powders.	5-90	0-94	1-15
Suspensions, Emulsions, Solutions (including Emulsifiable Concentrates)	5-50	40-95	0-15
Dusts	1-25	70-99	0-5
Granules and Pellets	0.01-99	5-99.99	0-15
High Strength Compositions	90-99	0-10	0-2

Typical solid diluents are described in Watkins, et al., *Handbook of Insecticide Dust Diluents and Carriers*, 2nd Ed., Dorland Books, Caldwell, New Jersey. Typical liquid diluents are described in Marsden, *Solvents Guide*, 2nd Ed., Interscience, New York, 1950. *McCutcheon's Detergents and Emulsifiers Annual*, Allured Publ. Corp., Ridgewood, New Jersey, as well as Sisely and Wood, *Encyclopedia of Surface Active Agents*, Chemical Publ. Co., Inc., New York, 1964, list surfactants and recommended uses. All formulations can contain minor amounts of additives to reduce foam, caking, corrosion, microbiological growth and the like, or thickeners to increase viscosity.

Surfactants include, for example, polyethoxylated alcohols, polyethoxylated alkylphenols, polyethoxylated sorbitan fatty acid esters, dialkyl sulfosuccinates, alkyl sulfates, alkylbenzene sulfonates, organosilicones, *N,N*-dialkyltaurates, lignin sulfonates, naphthalene sulfonate formaldehyde condensates, polycarboxylates, and

5 polyoxyethylene/polyoxypropylene block copolymers. Solid diluents include, for example, clays such as bentonite, montmorillinite, attapulgite and kaolin, starch, sugar, silica, talc, diatomaceous earth, urea, calcium carbonate, sodium carbonate and bicarbonate, and sodium sulfate. Liquid diluents include, for example, water, *N,N*-dimethylformamide, dimethyl sulfoxide, *N*-alkylpyrrolidone, ethylene glycol,

10 polypropylene glycol, paraffins, alkylbenzenes, alkyl naphthalenes, oils of olive, castor, linseed, tung, sesame, corn, peanut, cotton-seed, soybean, rape-seed and coconut, fatty acid esters, ketones such as cyclohexanone, 2-heptanone, isophorone and 4-hydroxy-4-methyl-2-pentanone, and alcohols such as methanol, cyclohexanol, decanol and tetrahydrofurfuryl alcohol.

15 Solutions, including emulsifiable concentrates, can be prepared by simply mixing the ingredients. Dusts and powders can be prepared by blending and, usually, grinding as in a hammer mill or fluid-energy mill. Suspensions are usually prepared by wet-milling; see, for example, U.S. 3,060,084. Granules and pellets can be prepared by spraying the active material upon preformed granular carriers or by agglomeration techniques. See

20 Browning, "Agglomeration", *Chemical Engineering*, December 4, 1967, pp 147-48, *Perry's Chemical Engineer's Handbook*, 4th Ed., McGraw-Hill, New York, 1963, pages 8-57 and following, and WO 91/13546. Pellets can be prepared as described in U.S. 4,172,714. Water-dispersible and water-soluble granules can be prepared as taught in U.S. 4,144,050, U.S. 3,920,442 and DE 3,246,493. Tablets can be prepared as taught

25 in U.S. 5,180,587, U.S. 5,232,701 and U.S. 5,208,030. Films can be prepared as taught in GB 2,095,558 and U.S. 3,299,566.

For further information regarding the art of formulation, see U.S. 3,235,361, Col. 6, line 16 through Col. 7, line 19 and Examples 10-41; U.S. 3,309,192, Col. 5, line 43 through Col. 7, line 62 and Examples 8, 12, 15, 39, 41, 52, 53, 58, 132, 138-140,

30 162-164, 166, 167 and 169-182; U.S. 2,891,855, Col. 3, line 66 through Col. 5, line 17 and Examples 1-4; Klingman, *Weed Control as a Science*, John Wiley and Sons, Inc., New York, 1961, pp 81-96; and Hance et al., *Weed Control Handbook*, 8th Ed., Blackwell Scientific Publications, Oxford, 1989.

In the following Examples, all percentages are by weight and all formulations are

35 prepared in conventional ways. Compound numbers refer to compounds in Index Tables A-F.

Example AHigh Strength Concentrate

	Compound 8	98.5%
	silica aerogel	0.5%
5	synthetic amorphous fine silica	1.0%.

Example BWettable Powder

	Compound 18	65.0%
	dodecylphenol polyethylene glycol ether	2.0%
10	sodium ligninsulfonate	4.0%
	sodium silicoaluminate	6.0%
	montmorillonite (calcined)	23.0%.

Example CGranule

15	Compound 8	10.0%
	attapulgite granules (low volatile matter, 0.71/0.30 mm; U.S.S. No. 25-50 sieves)	90.0%.

Example D20 Extruded Pellet

	Compound 18	25.0%
	anhydrous sodium sulfate	10.0%
	crude calcium ligninsulfonate	5.0%
	sodium alkyl naphthalenesulfonate	1.0%
25	calcium/magnesium bentonite	59.0%.

Tests results indicate that the compounds of the present invention are highly active preemergent and/or postemergent herbicides and/or plant growth regulants. Many of them have utility for broad-spectrum pre- and/or postemergence weed control in areas where complete control of all vegetation is desired such as around fuel storage tanks, industrial storage areas, parking lots, drive-in theaters, around billboards and highway and railroad structures. Some of the compounds are useful for the control of selected grass and broadleaf weeds with tolerance to important agronomic crops which include but are not limited to barley, cotton, wheat, rape, sugarbeets, corn, soybeans, rice, and plantation crops including coffee, cocoa, oil palm, rubber, sugarcane, citrus, grapes, fruit trees, nut trees, banana, plantain, pineapple, conifers, e.g., loblolly pine, and turf species, e.g., Kentucky bluegrass, St. Augustine grass, Kentucky fescue and bermudagrass. Those skilled in the art will appreciate that not all compounds are equally effective

against all weeds. Alternatively, the subject compounds are useful to modify plant growth.

Compounds of this invention can be used alone or in combination with other commercial herbicides, insecticides or fungicides. A mixture of one or more of the following herbicides with a compound of this invention may be particularly useful for weed control: acetochlor, acifluorfen and its sodium salt, acrolein (2-propenal), alachlor, ametryn, amidosulfuron, amitrole, ammonium sulfamate, anilofos, asulam, atrazine, azimsulfuron, benazolin, benazolin-ethyl, benfluralin, benfuresate, bensulfuron-methyl, bensulide, bentazone, bifenox, bromacil, bromoxynil, bromoxynil octanoate, butachlor, butralin, butylate, chlomethoxyfen, chloramben, chlorbromuron, chloridazon, chlorimuron-ethyl, chlornitrofen, chlorotoluron, chlorpropham, chlorsulfuron, chlorthal-dimethyl, cinmethylin, cinosulfuron, clethodim, clomazone, clopyralid, clopyralid-olamine, cyanazine, cycloate, cyclosulfamuron, 2,4-D and its butotyl, butyl, isooctyl and isopropyl esters and its dimethylammonium, diolamine and trolamine salts, daimuron, dalapon, dalapon-sodium, dazomet, 2,4-DB and its dimethylammonium, potassium and sodium salts, desmedipham, desmetryn, dicamba and its dimethylammonium, potassium and sodium salts, dichlobenil, dichlorprop, diclofop-methyl, difenzoquat metilsulfate, diflufenican, dimepiperate, dimethylarsinic acid and its sodium salt, dinitramine, diphenamid, diquat dibromide, dithiopyr, diuron, DNOC, endothal, EPTC, esprocarb, ethalfluralin, ethametsulfuron-methyl, ethofumesate, ethyl α ,2-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1*H*-1,2,4-triazol-1-yl]-4-fluorobenzenepropanoate (F8426), fenoxaprop-ethyl, fenoxaprop-P-ethyl, fenuron, fenuron-TCA, flamprop-methyl, flamprop-M-isopropyl, flamprop-M-methyl, flazasulfuron, fluazifop-butyl, fluazifop-P-butyl, fluchloralin, flumetsulam, flumiclorac-pentyl, flumioxazin, fluometuron, fluoroglycofen-ethyl, flupoxam, fluridone, flurochloridone, fluroxypyr, fomesafen, fosamine-ammonium, glufosinate, glufosinate-ammonium, glyphosate, glyphosate-isopropylammonium, glyphosate-sesquisodium, glyphosate-trimesium, halosulfuron-methyl, haloxyfop-etotyl, haloxyfop-methyl, hexazinone, imazamethabenz-methyl, imazapyr, imazaquin, imazaquin-ammonium, imazethapyr, imazethapyr-ammonium, imazosulfuron, ioxynil, ioxynil octanoate, ioxynil-sodium, isoproturon, isouron, isoxaben, lactofen, lenacil, linuron, maleic hydrazide, MCPA and its dimethylammonium, potassium and sodium salts, MCPA-isooctyl, mecoprop, mecoprop-P, mefenacet, mefluidide, metam-sodium, methabenzthiazuron, methyl [[2-chloro-4-fluoro-5-[(tetrahydro-3-oxo-1*H*,3*H*-[1,3,4]thiadiazolo[3,4-*a*]pyridazin-1-ylidene)amino]phenyl]thioacetate (KIH 9201), methylarsonic acid and its calcium, monoammonium, monosodium and disodium salts, methyl [[[1-[5-[2-chloro-4-(trifluoromethyl)phenoxy]-2-nitrophenyl]-2-methoxyethylidene]amino]oxy]acetate (AKH-7088), methyl 5-[[[(4,6-dimethyl-2-

pyrimidinyl)amino]carbonyl]amino]sulfonyl]-1-(2-pyridinyl)-1*H*-pyrazole-4-carboxylate (NC-330), metobenzuron, metolachlor, metoxuron, metribuzin, metsulfuron-methyl, molinate, monolinuron, napropamide, naptalam, neburon, nicosulfuron, norflurazon, oryzalin, oxadiazon, oxyfluorfen, paraquat dichloride, pebulate, pendimethalin, 5 perfluidone, phenmedipham, picloram, picloram-potassium, pretilachlor, primisulfuron-methyl, prometon, prometryn, propachlor, propanil, propazine, propham, propyzamide, prosulfuron, pyrazolynate, pyrazosulfuron-ethyl, quinclorac, quizalofop-ethyl, quizalofop-P-ethyl, quizalofop-P-tefuryl, rimsulfuron, sethoxydim, siduron, simazine, sulcotrione (ICIA0051), sulfentrazone, sulfometuron-methyl, TCA, 10 TCA-sodium, tebuthiuron, terbacil, terbuthylazine, terbutryn, thenylchlor, thifensulfuron-methyl, thiobencarb, tralkoxydim, tri-allate, triasulfuron, tribenuron-methyl, triclopyr, triclopyr-butotyl, triclopyr-triethylammonium, tridiphane, trifluralin, triflusulfuron-methyl, and vernolate.


In certain instances, combinations with other herbicides having a similar spectrum 15 of control but a different mode of action will be particularly advantageous for preventing the development of resistant weeds.

Preferred are mixtures of a compound of Formula I with a compound selected from the group diuron, fluazifop-butyl, fluazifop-P-butyl, glufosinate, glufosinate-ammonium, glyphosate, glyphosate-isopropylammonium, glyphosate-sesquisodium, 20 glyphosate-trimesium, hexazinone, metribuzin, norflurazon, paraquat, quizalofop-ethyl, and quizalofop-P-ethyl. Specifically preferred are mixtures of a compound selected from the group 3-(7-chloro-1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl)-1-methyl-6-(trifluoromethyl)-2,4(1*H*,3*H*)-pyrimidinedione *S,S*-dioxide and 3-[(7-chloro-1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl)imino]tetrahydro-1*H*,3*H*- 25 [1,3,4]thiadiazolo[3,4-*a*]pyridazin-1-one *S,S*-dioxide with a compound selected from the group diuron, fluazifop-butyl, fluazifop-P-butyl, glufosinate, glufosinate-ammonium, glyphosate, glyphosate-isopropylammonium, glyphosate-sesquisodium, glyphosate-trimesium, hexazinone, metribuzin, norflurazon, paraquat, quizalofop-ethyl, and quizalofop-P-ethyl.

30 A herbicidally effective amount of the compounds of this invention is determined by a number of factors. These factors include: formulation selected, method of application, amount and type of vegetation present, growing conditions, etc. In general, a herbicidally effective amount of a compound(s) of this invention is applied at rates from about 0.001 to 20 kg/ha with a preferred rate range of 0.004 to 1.0 kg/ha. One skilled in 35 the art can easily determine application rates necessary for the desired level of weed control.

The following Tests demonstrate the control efficacy of the compounds of this invention against specific weeds. The weed control afforded by the compounds is not limited, however, to these species. See Index Tables A-F for compound descriptions.

Index Table A



5


Cmpd No.	R ¹	R ²	R ³	R ⁴	R ⁵	m.p. (°C)
1 (Ex. 1)	F	H	CH ₂ CH ₃	H	H	189-190
2 (Ex. 2)	F	Cl	CH ₂ CH ₃	H	H	190-192
3 (Ex. 13)	F	Cl	CH ₂ -(4-CH ₃ O-Ph)	H	H	171-174
4 (Ex. 13)	F	Cl	H	H	H	75-79
5 (Ex. 14)	F	Cl	C(=O)CH ₃	H	H	205-208
6 ^a	F	Cl	CH ₂ C≡CH	H	H	solid*

* See Index Table F for ¹H NMR data.

^a Compound contains 50% by weight of 2-[1,3-bis(2-propynyl)-7-chloro-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl]-4,5,6,7-tetrahydro-1*H*-isoindole-1,3(2*H*)-dione *S,S*-dioxide.


10

Index Table B




Cmpd No.	R ¹	R ²	R ³	R ⁴	R ⁵	R ¹³	R ¹⁴	m.p. (°C)
7 (Ex. 4)	F	Cl	CH ₂ CH ₃	H	H	H	CF ₃	>230
8 (Ex. 5)	F	Cl	CH ₂ CH ₃	H	H	CH ₃	CF ₃	147-149
9 (Ex. 6)	F	Cl	CH ₂ CH ₃	H	H	CH ₂ CH ₃	CF ₃	177-178
10 (Ex. 15)	F	Cl	CH ₂ -(4-CH ₃ O-Ph)	H	H	H	CF ₃	>230
11 (Ex. 16)	F	Cl	CH ₂ -(4-CH ₃ O-Ph)	H	H	CH ₃	CF ₃	180-182
12 (Ex. 17)	F	Cl	H	H	H	CH ₃	CF ₃	>230
13 (Ex. 18)	F	Cl	C(=O)CH ₃	H	H	CH ₃	CF ₃	213-215

60

Index Table C


<u>Cmpd No.</u>	<u>R¹</u>	<u>R²</u>	<u>R³</u>	<u>R⁴</u>	<u>R⁵</u>	<u>Z¹</u>	<u>m.p. (°C)</u>
14 (Ex. 8)	F	Cl	CH ₂ CH ₃	H	H	CH ₂	168-169
15 (Ex. 9)	F	Cl	CH ₂ CH ₃	H	H	O	200-202

Index Table D

5

<u>Cmpd No.</u>	<u>R¹</u>	<u>R²</u>	<u>R³</u>	<u>R⁴</u>	<u>R⁵</u>	<u>J</u>	<u>m.p. (°C)</u>
16 (Ex. 10)	F	Cl	CH ₂ CH ₃	H	H		oil*
17 (Ex. 11)	F	Cl	CH ₂ CH ₃	H	H		87-90
18 (Ex. 7)	F	Cl	CH ₂ CH ₃	H	H		143-146
19 (Ex. 12)	F	Cl	CH ₂ CH ₃	H	H		semisolid*

* See Index Table F for ¹H NMR data.

Index Table E

<u>Cmpd No.</u>	<u>R¹</u>	<u>R³</u>	<u>R⁴</u>	<u>R⁵</u>	<u>X</u>	<u>m.p. (°C)</u>
20 (Ex. 19)	F	CH ₂ Ph	H	H	O	150-152
21 (Ex. 3)	F	CH ₂ CH ₃	H	H	direct bond	oil*

5 * See Index Table F for ¹H NMR data.

Index Table F

<u>Cmpd No.</u>	<u>¹H NMR data (in CDCl₃ solution unless otherwise indicated)^a</u>
6	δ 7.26 (d,1H), 4.58 (s,2H), 4.48 (m,2H), 2.42 (br s,4H), 2.21 (t,1H), 1.82 (br s,2H).
16	δ 7.30 (d,1H,J=9.6 Hz), 5.99 (s,2H), 4.31 (s,2H), 4.21 (s,4H), 3.93 (dd,2H), 1.28 (t,3H).
19	δ 7.26 (d,1H), 7.04 (t,1H), 4.28 (s,2H), 3.96 (q,2H), 2.44 (s,3H), 1.26 (t,3H).
21	δ 7.08 (d,1H), 6.62 (d,2H), 4.32 (s,2H), 3.64 (q,2H), 2.42 (m,4H), 1.82 (m,4H), 1.38 (t,3H):

^a ¹H NMR data are in ppm downfield from tetramethylsilane. Couplings are designated by (s)-singlet, (d)-doublet, (t)-triplet, (q)-quartet, (m)-multiplet, (dd)-doublet of doublets, (br s)-broad singlet.

TEST A

Seeds of barley (*Hordeum vulgare*), barnyardgrass (*Echinochloa crus-galli*), bedstraw (*Galium aparine*), blackgrass (*Alopecurus myosuroides*), chickweed (*Stellaria media*), cocklebur (*Xanthium pensylvanicum*), corn (*Zea mays*), cotton (*Gossypium hirsutum*), crabgrass (*Digitaria sanguinalis*), downy brome (*Bromus tectorum*), giant foxtail (*Setaria faberii*), lambsquarters (*Chenopodium album*), morningglory (*Ipomoea hederacea*), rape (*Brassica napus*), rice (*Oryza sativa*), sorghum (*Sorghum bicolor*), soybean (*Glycine max*), sugar beet (*Beta vulgaris*), velvetleaf (*Abutilon theophrasti*), wheat (*Triticum aestivum*), wild buckwheat (*Polygonum convolvulus*), wild oat (*Avena fatua*) and purple nutsedge (*Cyperus rotundus*) tubers were planted and treated preemergence with test chemicals formulated in a non-phytotoxic solvent mixture which includes a surfactant.

At the same time, these crop and weed species were also treated with postemergence applications of test chemicals formulated in the same manner. Plants ranged in height from two to eighteen cm (one to four leaf stage) for postemergence treatments. Treated plants and controls were maintained in a greenhouse for twelve to sixteen days, after which all species were compared to controls and visually evaluated. Plant response ratings, summarized in Table A, are based on a scale of 0 to 10 where 0 is no effect and 10 is complete control. A dash (-) response means no test result.

TABLE A	COMPOUND	TABLE A	COMPOUND
Rate 2000 g/ha	7	Rate 2000 g/ha	7
POSTEMERGENCE		PREEMERGENCE	
Barley	0	Barley	0
Barnyardgrass	0	Barnyardgrass	0
Bedstraw	2	Bedstraw	0
Blackgrass	1	Blackgrass	0
Chickweed	0	Chickweed	0
Cocklebur	2	Cocklebur	0
Corn	1	Corn	0
Cotton	2	Cotton	0
Crabgrass	1	Crabgrass	0
Downy brome	0	Downy brome	0
Giant foxtail	1	Giant foxtail	0
Lambsquarter	1	Lambsquarter	0
Morningglory	2	Morningglory	3
Nutsedge	0	Nutsedge	0
Rape	2	Rape	0
Rice	0	Rice	0
Sorghum	0	Sorghum	0
Soybean	2	Soybean	2
Sugar beet	1	Sugar beet	0
Velvetleaf	1	Velvetleaf	0
Wheat	0	Wheat	0
Wild buckwheat	2	Wild buckwheat	0
Wild oat	0		

TABLE A	COMPOUND			
Rate 1000 g/ha	1	2	16	20
POSTEMERGENCE				
Barley	3	4	3	3
Barnyardgrass	3	9	6	6
Bedstraw	7	10	6	6
Blackgrass	4	5	3	2
Chickweed	3	8	5	2
Cocklebur	6	9	8	8
Corn	3	5	2	2
Cotton	9	10	10	9
Crabgrass	3	6	5	7
Downy brome	3	2	2	2
Giant foxtail	3	6	7	4
Lambsquarter	8	9	9	8
Morningglory	9	9	9	8
Nutsedge	0	2	2	2
Rape	7	10	10	7
Rice	5	6	7	5
Sorghum	3	8	6	2
Soybean	5	8	9	5
Sugar beet	9	10	10	4
Velvetleaf	8	10	10	6
Wheat	4	4	3	2
Wild buckwheat	10	10	9	6
Wild oat	3	4	2	2

TABLE A	COMPOUND			
Rate 1000 g/ha	1	2	16	20
PREEMERGENCE				
Barley	0	0	0	0
Barnyardgrass	6	9	8	0
Bedstraw	8	10	10	-
Blackgrass	3	3	2	2
Chickweed	3	9	2	-
Cocklebur	0	4	8	3
Corn	2	3	3	2
Cotton	0	10	10	0
Crabgrass	4	9	8	3
Downy brome	2	3	0	-
Giant foxtail	2	8	9	4
Lambsquarter	9	10	10	-
Morningglory	1	10	8	4
Nutsedge	0	5	-	-
Rape	6	10	10	-
Rice	0	4	3	0
Sorghum	0	2	7	0
Soybean	0	7	8	0
Sugar beet	8	10	10	-
Velvetleaf	8	10	10	2
Wheat	0	2	3	0
Wild buckwheat	0	10	9	-
Wild oat	2	4	2	0

TABLE A	COMPOUND
Rate 400 g/ha	7 12
POSTEMERGENCE	
Barley	0 10
Barnyardgrass	1 10
Bedstraw	1 10
Blackgrass	0 6
Chickweed	0 10
Cocklebur	1 10
Corn	0 10
Cotton	- 10
Crabgrass	0 8
Downy brome	0 8
Giant foxtail	0 9
Lambsquarter	0 10
Morningglory	- 10
Nutsedge	0 5
Rape	0 10
Rice	0 10
Sorghum	0 10
Soybean	1 10
Sugar beet	1 10
Velvetleaf	1 10
Wheat	0 10
Wild buckwheat	1 10
Wild oat	0 10

TABLE A	COMPOUND
Rate 400 g/ha	7 12
PREEMERGENCE	
Barley	0 0
Barnyardgrass	0 3
Bedstraw	0 10
Blackgrass	0 2
Chickweed	0 2
Cocklebur	0 10
Corn	0 0
Cotton	0 10
Crabgrass	0 4
Downy brome	0 3
Giant foxtail	0 5
Lambsquarter	0 9
Morningglory	1 10
Nutsedge	0 7
Rape	0 9
Rice	0 5
Sorghum	0 0
Soybean	0 9
Sugar beet	0 10
Velvetleaf	0 10
Wheat	0 0
Wild buckwheat	0 6
Wild oat	0 0

TABLE A	COMPOUND																				
	Rate 200 g/ha	1	2	3	4	5	6	8	9	10	11	13	14	15	16	17	18	20	21		
POSTEMERGENCE																					
Barley	3	3	5	2	3	5	8	6	0	4	6	2	1	3	5	9	3	2			
Barnyardgrass	2	7	4	10	10	7	10	10	0	6	10	3	7	2	9	9	6	5			
Bedstraw	5	10	4	8	9	9	10	10	0	10	10	5	6	5	10	10	5	6			
Blackgrass	3	3	2	2	6	3	7	9	0	3	2	3	2	2	3	6	1	1			
Chickweed	3	4	2	6	9	4	10	3	0	7	10	3	6	3	3	8	1	3			
Cocklebur	6	9	10	10	10	10	10	3	0	10	10	7	6	7	8	10	7	7			
Corn	2	3	3	6	2	3	9	9	0	2	9	1	4	1	3	5	2	2			
Cotton	9	10	-	10	10	10	10	9	0	10	10	10	10	9	10	10	10	9			
Crabgrass	2	4	5	6	4	4	9	2	0	2	6	2	2	2	5	9	4	5			
Downy brome	2	3	2	2	2	1	7	5	0	2	2	1	0	1	3	5	2	1			
Giant foxtail	2	4	6	5	5	4	9	6	0	2	6	2	3	2	6	9	3	3			
Lambsquarter	8	9	10	10	9	10	10	9	0	10	10	8	8	8	9	10	9	8			
Morningglory	8	9	10	10	10	10	10	8	0	10	10	9	9	9	10	10	9	9			
Nutsedge	0	2	1	4	2	2	5	-	0	0	3	2	-	1	8	-	1	3			
Rape	3	10	10	10	10	10	10	9	0	10	10	5	8	7	10	10	6	6			
Rice	3	6	4	5	6	4	10	3	0	3	10	4	4	3	5	7	3	5			
Sorghum	3	3	4	4	6	4	10	9	0	5	9	2	3	3	6	8	2	4			
Soybean	3	8	5	5	5	5	10	9	0	8	10	8	9	8	9	10	3	7			
Sugar beet	8	10	10	10	10	10	10	10	0	10	10	9	10	9	10	10	5	10			
Velvetleaf	6	10	10	10	10	10	10	10	0	10	10	10	10	10	10	10	7	9			
Wheat	3	4	3	3	4	3	10	9	0	3	6	4	3	2	2	9	2	1			
Wild buckwheat	8	10	10	10	10	10	10	10	0	10	10	7	10	8	10	10	3	8			
Wild oat	2	3	2	2	3	2	9	9	0	.3	3	4	4	1	2	7	2	1			

TABLE A	COMPOUND																				
	Rate 200 g/ha	1	2	3	4	5	6	8	9	10	11	13	14	15	16	17	18	20	21		
PREEMERGENCE																					
Barley	0	0	0	0	0	0	6	2	0	0	0	0	0	2	0	3	0	0			
Barnyardgrass	0	9	0	0	2	4	10	9	0	2	0	3	2	3	7	9	0	5			
Bedstraw	2	10	8	0	3	2	10	10	0	10	10	7	3	2	10	6	0	9			
Blackgrass	2	3	1	0	0	0	9	0	0	0	0	3	1	1	0	5	0	2			
Chickweed	0	7	0	0	0	0	10	8	0	0	0	3	4	0	0	3	0	0			
Cocklebur	0	3	7	4	2	7	10	1	0	3	10	3	3	6	4	6	2	0			
Corn	0	0	0	0	0	2	7	1	0	0	0	2	3	0	3	2	0	0			
Cotton	0	0	0	0	0	0	10	7	0	0	9	6	6	7	10	3	0	3			
Crabgrass	0	8	3	0	0	2	10	10	0	9	6	8	2	2	7	6	2	2			
Downy brome	3	2	1	0	0	0	10	0	0	2	0	0	0	0	2	4	0	1			
Giant foxtail	0	6	0	0	0	3	10	10	0	9	0	6	6	4	9	9	0	0			
Lambsquarter	5	10	10	10	10	10	10	10	0	10	9	10	10	10	10	10	-	10			
Morningglory	0	10	0	0	6	9	10	1	0	10	10	3	4	8	10	7	0	0			
Nutsedge	0	0	-	0	0	-	6	0	-	0	0	-	-	0	1	0	0	0			
Rape	3	9	0	0	7	0	10	9	0	2	8	4	7	9	10	7	0	8			
Rice	0	0	0	0	0	0	9	0	0	0	3	1	3	1	4	2	0	0			
Sorghum	0	0	0	0	0	0	10	8	0	0	0	2	1	0	7	2	0	0			
Soybean	0	5	2	0	0	0	10	2	0	0	10	1	3	7	7	7	0	0			
Sugar beet	0	10	8	2	7	8	10	10	0	10	10	10	10	7	10	10	2	10			
Velvetleaf	2	10	7	3	8	6	10	10	0	10	10	8	10	10	10	10	0	0			
Wheat	0	2	0	0	0	0	9	5	0	1	2	3	0	0	0	6	0	0			
Wild buckwheat	0	9	0	0	2	0	10	9	0	0	3	0	3	0	10	5	-	2			
Wild oat	2	2	0	0	0	0	8	1	0	1	3	3	0	2	3	6	0	0			

TABLE A	COMPOUND
Rate 100 g/ha	12
POSTEMERGENCE	
Barley	8
Barnyardgrass	10
Bedstraw	10
Blackgrass	3
Chickweed	10
Cocklebur	10
Corn	8
Cotton	10
Crabgrass	7
Downy brome	5
Giant foxtail	6
Lambsquarter	10
Morningglory	10
Nutsedge	3
Rape	10
Rice	10
Sorghum	10
Soybean	8
Sugar beet	10
Velvetleaf	10
Wheat	10
Wild buckwheat	10
Wild oat	9

TABLE A	COMPOUND
Rate 100 g/ha	12
PREEMERGENCE	
Barley	0
Barnyardgrass	0
Bedstraw	6
Blackgrass	0
Chickweed	0
Cocklebur	6
Corn	0
Cotton	0
Crabgrass	0
Downy brome	0
Giant foxtail	0
Lambsquarter	6
Morningglory	10
Nutsedge	3
Rape	6
Rice	2
Sorghum	0
Soybean	8
Sugar beet	8
Velvetleaf	3
Wheat	0
Wild buckwheat	0
Wild oat	0

TABLE A	COMPOUND																				
	Rate 50 g/ha	3	4	5	6	8	9	10	11	13	14	15	17	18	19	21					
POSTEMERGENCE																					
Barley	3	2	3	4	3	4	0	3	7	0	0	3	5	3	2						
Barnyardgrass	3	6	10	6	10	9	0	2	10	1	2	6	9	4	3						
Bedstraw	4	7	9	9	10	9	0	10	10	3	6	9	8	6	3						
Blackgrass	2	1	3	2	7	8	0	3	2	2	1	0	6	3	1						
Chickweed	2	4	6	2	10	3	0	7	10	2	1	2	8	8	2						
Cocklebur	10	9	10	10	10	2	0	10	10	7	3	7	10	7	5						
Corn	3	3	2	3	9	5	0	2	9	1	2	3	2	3	2						
Cotton	10	10	10	10	10	9	0	10	10	10	10	10	10	10	9						
Crabgrass	3	3	4	4	6	3	0	2	6	2	2	3	4	3	4						
Downy brome	1	1	2	1	6	4	0	1	3	1	1	2	5	3	1						
Giant foxtail	4	3	6	3	6	5	0	2	6	2	2	6	8	3	3						
Lambsquarter	9	8	9	9	10	8	0	10	10	8	9	9	9	10	6						
Morningglory	9	10	10	10	10	5	0	10	10	9	7	10	10	9	4						
Nutsedge	1	1	-	2	3	-	0	0	3	-	1	7	2	1	1						
Rape	9	10	10	10	10	9	0	10	10	3	5	9	9	3	3						
Rice	3	3	4	3	9	2	0	3	10	3	3	5	4	3	4						
Sorghum	3	3	6	4	6	4	0	2	10	2	2	4	7	5	4						
Soybean	2	3	4	2	10	9	0	8	9	8	7	9	7	3	5						
Sugar beet	10	10	10	10	10	6	0	10	10	7	9	10	10	10	9						
Velvetleaf	10	10	10	10	10	9	0	10	10	10	6	10	10	9	6						
Wheat	3	3	4	2	10	7	0	2	9	1	2	2	7	3	1						
Wild buckwheat	10	10	10	10	10	9	0	10	10	7	9	9	10	10	6						
Wild oat	1	2	3	2	5	9	0	2	6	3	3	1	6	2	1						

TABLE A	COMPOUND																		
	Rate 50 g/ha	3	4	5	6	8	9	10	11	13	14	15	17	18	19	21			
PREEMERGENCE																			
Barley	0	0	0	0	3	0	0	0	0	0	0	0	0	3	0	0			
Barnyardgrass	0	0	0	0	10	1	0	0	0	0	0	0	4	6	3	1			
Bedstraw	0	0	2	0	10	9	0	10	10	0	0	10	3	0	0				
Blackgrass	0	0	0	0	3	0	0	0	0	2	0	0	0	0	0				
Chickweed	0	0	0	0	10	1	0	0	0	0	0	0	0	0	0				
Cocklebur	3	0	0	0	9	0	0	2	0	2	0	4	2	0	0				
Corn	0	0	0	0	5	0	0	0	0	0	0	2	0	3	0				
Cotton	0	0	0	0	10	2	0	0	1	0	3	9	0	10	0				
Crabgrass	0	0	0	0	8	9	0	3	0	0	0	2	0	0	0				
Downy brome	0	0	0	0	7	0	0	0	0	2	0	0	3	1	2				
Giant foxtail	0	0	0	0	10	8	0	4	0	0	0	7	2	3	0				
Lambsquarter	3	5	9	8	10	9	0	10	7	10	9	8	9	7	9				
Morningglory	0	0	3	0	10	0	0	10	10	0	0	3	2	0	0				
Nutsedge	-	-	0	0	1	0	-	-	0	-	-	0	0	0	0				
Rape	0	0	0	0	10	4	0	-	2	0	0	10	1	0	0				
Rice	0	0	0	0	5	0	0	0	0	0	0	1	0	0	0				
Sorghum	0	0	0	0	4	0	0	0	0	0	0	3	0	0	0				
Soybean	0	0	0	0	9	0	0	0	0	0	0	7	0	3	0				
Sugar beet	0	2	3	7	10	5	0	8	7	7	9	9	9	9	9				
Velvetleaf	0	0	2	3	10	1	0	10	2	6	2	10	6	5	0				
Wheat	0	0	0	0	5	2	0	1	0	0	0	0	2	1	0				
Wild buckwheat	0	0	0	0	10	4	0	0	0	0	0	8	0	0	0				
Wild oat	0	0	0	0	3	2	0	0	0	0	0	0	3	1	0				

TABLE A	COMPOUND	TABLE A	COMPOUND
Rate	10 g/ha	Rate	10 g/ha
POSTEMERGENCE		PREEMERGENCE	
Barley	2	Barley	0
Barnyardgrass	3	Barnyardgrass	3
Bedstraw	3	Bedstraw	0
Blackgrass	1	Blackgrass	0
Chickweed	3	Chickweed	0
Cocklebur	4	Cocklebur	0
Corn	2	Corn	0
Cotton	-	Cotton	5
Crabgrass	3	Crabgrass	0
Downy brome	1	Downy brome	0
Giant foxtail	2	Giant foxtail	0
Lambsquarter	6	Lambsquarter	4
Morningglory	6	Morningglory	0
Nutsedge	1	Nutsedge	0
Rape	1	Rape	0
Rice	2	Rice	2
Sorghum	3	Sorghum	0
Soybean	3	Soybean	3
Sugar beet	10	Sugar beet	1
Velvetleaf	8	Velvetleaf	3
Wheat	3	Wheat	0
Wild buckwheat	10	Wild buckwheat	0
Wild oat	1	Wild oat	0

TEST B

The compounds evaluated in this test were formulated in a non-phytotoxic solvent mixture which includes a surfactant and applied to the soil surface before plant seedlings emerged (preemergence application), to water that covered the soil surface (flood application), and to plants that were in the one-to-four leaf stage (postemergence application). A sandy loam soil was used for the preemergence and postemergence tests, while a silt loam soil was used in the flood test. Water depth was approximately 2.5 cm for the flood test and was maintained at this level for the duration of the test.

Plant species in the preemergence and postemergence tests consisted of barnyardgrass (*Echinochloa crus-galli*), barley (*Hordeum vulgare*), bedstraw (*Galium aparine*), blackgrass (*Alopecurus myosuroides*), chickweed (*Stellaria media*), cocklebur (*Xanthium pensylvanicum*), corn (*Zea mays*), cotton (*Gossypium hirsutum*), crabgrass (*Digitaria sanguinalis*), downy brome (*Bromus tectorum*), giant foxtail (*Setaria faberii*), johnsongrass (*Sorghum halpense*), lambsquarters (*Chenopodium album*), morningglory (*Ipomoea hederacea*), pigweed (*Amaranthus retroflexus*), rape (*Brassica napus*), ryegrass (*Lolium multiflorum*), soybean (*Glycine max*), speedwell (*Veronica persica*),

sugar beet (*Beta vulgaris*), velvetleaf (*Abutilon theophrasti*), wheat (*Triticum aestivum*), wild buckwheat (*Polygonum convolvulus*), and wild oat (*Avena fatua*). All plant species were planted one day before application of the compound for the preemergence portion of this test. Plantings of these species were adjusted to produce plants of appropriate size
5 for the postemergence portion of the test. Plant species in the flood test consisted of rice (*Oryza sativa*), umbrella sedge (*Cyperus difformis*), duck salad (*Heteranthera limosa*), barnyardgrass (*Echinochloa crus-galli*) and late watergrass (*Echinochloa oryzicola*) grown to the 2 leaf stage for testing.

All plant species were grown using normal greenhouse practices. Visual
10 evaluations of injury expressed on treated plants, when compared to untreated controls, were recorded approximately fourteen to twenty one days after application of the test compound. Plant response this ratings, summarized in Table B, were recorded on a 0 to 100 scale where 0 is no effect and 100 is complete control. A dash (-) response means no test result.

TABLE B	COMPOUND	TABLE B	COMPOUND		
Rate 500 g/ha	2	Rate 250 g/ha	2	9	14
POSTEMERGENCE		POSTEMERGENCE			
Barley Igri	-	Barley Igri	35	65	30
Barnyardgr. (Flood)	70	Barnyardgr. (Flood)	35	95	25
Barnyardgrass	-	Barnyardgrass	45	100	40
Bedstraw	-	Bedstraw	85	100	35
Blackgrass	-	Blackgrass	10	90	40
Chickweed	-	Chickweed	30	70	70
Cocklebur	-	Cocklebur	90	75	80
Corn	-	Corn	25	90	40
Cotton	-	Cotton	100	100	100
Crabgrass	-	Crabgrass	25	70	55
Downy Brome	-	Downy Brome	0	35	40
Duck salad	25	Duck salad	20	0	95
Giant foxtail	-	Giant foxtail	40	90	50
Italn. Rygrass	-	Italn. Rygrass	0	30	35
Johnsongrass	-	Johnsongrass	50	90	75
Lambsquarter	-	Lambsquarter	100	100	95
Morningglory	-	Morningglory	100	90	100
Rape	-	Rape	100	100	90
Redroot Pigweed	-	Redroot Pigweed	-	100	90
Rice Japonica	60	Rice Japonica	45	75	70
Soybean	-	Soybean	-	60	60
Speedwell	-	Speedwell	100	-	100
Sugar beet	-	Sugar beet	100	100	100
Umbrella sedge	65	Umbrella sedge	45	60	100
Velvetleaf	-	Velvetleaf	100	100	100
Watergrass	75	Watergrass	35	80	30
Wheat	-	Wheat	30	60	30
Wild buckwheat	-	Wild buckwheat	65	100	100
Wild oat	-	Wild oat	35	90	40

TABLE B	COMPOUND		
	Rate 250 g/ha	2	9 14
PREEMERGENCE			
Barley Igri	10	10	0
Barnyardgrass	40	100	90
Bedstraw	20	100	85
Blackgrass	0	65	45
Chickweed	10	30	10
Cocklebur	0	10	85
Corn	10	40	25
Cotton	40	80	95
Crabgrass	75	95	85
Downy Brome	0	0	0
Giant foxtail	85	100	90
Italn. Rygrass	0	90	75
Johnsongrass	90	100	45
Lambsquarter	95	100	100
Morningglory	100	60	25
Rape	30	95	85
Redroot Pigweed	100	100	90
Soybean	10	50	30
Speedwell	-	100	100
Sugar beet	95	100	100
Velvetleaf	100	100	100
Wheat	0	35	10
Wild buckwheat	0	100	85
Wild oat	35	95	30

TABLE B	COMPOUND			
	Rate 125 g/ha	2	9 14	17
POSTEMERGENCE				
Barley Igri	35	35	30	25
Barnyardgr. (Flood)	10	65	0	75
Barnyardgrass	35	100	35	75
Bedstraw	60	100	35	100
Blackgrass	10	80	35	30
Chickweed	30	50	70	60
Cocklebur	90	50	80	95
Corn	15	90	30	30
Cotton	100	100	100	100
Crabgrass	20	60	45	30
Downy Brome	0	25	20	20
Duck salad	0	0	80	95
Giant foxtail	40	85	30	50
Italn. Rygrass	0	20	20	30
Johnsongrass	40	90	40	75
Lambsquarter	85	100	65	100
Morningglory	100	80	90	100
Rape	100	100	90	100
Redroot Pigweed	100	100	90	90
Rice Japonica	30	45	45	75
Soybean	100	40	50	85
Speedwell	100	-	100	100
Sugar beet	100	100	100	100
Umbrella sedge	15	0	95	85
Velvetleaf	100	100	100	100
Watergrass	10	35	25	70
Wheat	25	45	20	30
Wild buckwheat	65	100	80	100
Wild oat	35	90	35	35

TABLE B	COMPOUND			
	Rate 125 g/ha	2	9	14
PREEMERGENCE				
Barley Igri	0	10	0	20
Barnyardgrass	20	85	10	70
Bedstraw	0	100	0	100
Blackgrass	0	50	45	0
Chickweed	10	10	0	0
Cocklebur	0	0	35	70
Corn	0	25	0	20
Cotton	0	70	50	100
Crabgrass	20	75	20	65
Downy Brome	0	0	0	10
Giant foxtail	50	100	20	55
Italn. Rygrass	0	20	0	0
Johnsongrass	30	50	30	50
Lambsquarter	95	100	95	100
Morningglory	0	50	20	80
Rape	0	75	10	100
Redroot Pigweed	100	100	10	100
Soybean	0	30	10	50
Speedwell	-	100	0	100
Sugar beet	30	100	0	100
Velvetleaf	100	100	10	100
Wheat	0	20	0	0
Wild buckwheat	0	90	35	100
Wild oat	25	85	0	20

TABLE B	COMPOUND								
	Rate 62 g/ha	2	3	4	6	8	9	14	17
POSTEMERGENCE									
Barley Igri	35	35	30	30	35	30	30	25	
Barnyardgr. (Flood)	10	30	25	25	100	10	0	60	
Barnyardgrass	35	25	70	30	100	95	30	60	
Bedstraw	60	70	100	100	100	95	0	100	
Blackgrass	10	20	25	45	55	80	30	25	
Chickweed	30	40	75	70	100	50	55	60	
Cocklebur	90	90	100	100	100	40	70	90	
Corn	10	40	35	50	90	70	25	30	
Cotton	100	100	100	100	100	100	100	100	
Crabgrass	15	50	40	40	70	50	35	30	
Downy Brome	0	15	20	30	10	10	20	0	
Duck salad	0	0	0	0	0	0	0	80	
Giant foxtail	35	60	50	60	90	70	20	30	
Italn. Rygrass	0	0	15	20	75	0	10	25	
Johnsongrass	30	40	70	70	100	90	30	60	
Lambsquarter	85	95	85	100	100	95	65	100	
Morningglory	100	100	100	100	100	60	90	95	
Rape	60	100	100	-	100	100	90	100	
Redroot Pigweed	95	100	100	100	100	100	75	70	
Rice Japonica	20	30	25	25	80	35	0	45	
Soybean	70	50	50	50	-	40	-	80	
Speedwell	100	90	95	100	100	-	80	100	
Sugar beet	100	100	100	100	100	100	60	100	
Umbrella sedge	10	0	0	0	0	0	80	75	
Velvetleaf	100	100	100	100	100	100	100	100	
Watergrass	0	20	15	15	100	35	0	65	
Wheat	25	35	10	15	40	40	10	0	
Wild buckwheat	65	100	100	95	100	100	35	100	
Wild oat	35	25	20	25	30	65	20	30	

TABLE B	COMPOUND									
	Rate	62 g/ha	2	3	4	6	8	9	14	17
PREEMERGENCE										
Barley Igri	0	0	0	0	0	10	0	10		
Barnyardgrass	10	20	0	30	100	35	0	20		
Bedstraw	0	25	10	0	100	85	0	95		
Blackgrass	0	0	0	10	20	10	20	0		
Chickweed	-	10	20	10	100	10	0	0		
Cocklebur	0	0	0	55	50	0	20	50		
Corn	0	0	0	0	40	0	0	0		
Cotton	0	0	0	0	100	50	20	100		
Crabgrass	0	0	0	0	95	35	15	40		
Downy Brome	0	0	0	0	10	0	0	0		
Giant foxtail	20	0	0	0	100	90	-	30		
Italn. Rygrass	0	0	0	0	50	10	0	0		
Johnsongrass	10	0	0	20	95	30	0	30		
Lambsquarter	40	90	85	95	100	100	0	100		
Morningglory	0	15	10	35	100	50	10	70		
Rape	0	0	0	0	100	65	0	80		
Redroot Pigweed	100	0	0	10	100	100	-	80		
Soybean	0	10	10	0	100	20	0	35		
Speedwell	-	10	50	25	100	100	0	85		
Sugar beet	10	0	10	25	100	100	0	100		
Velvetleaf	100	60	10	85	100	30	0	100		
Wheat	0	0	0	0	10	10	0	0		
Wild buckwheat	0	25	10	0	100	60	0	95		
Wild oat	10	0	0	0	50	20	0	0		

TABLE B	COMPOUND								
	Rate 31 g/ha	2	3	4	6	8	9	14	17
POSTEMERGENCE									
Barley Igri	30	35	-	30	35	20	20	20	
Barnyardgr. (Flood)	-	30	20	25	75	0	0	20	
Barnyardgrass	30	20	60	25	100	95	-	30	
Bedstraw	50	70	90	100	100	90	0	65	
Blackgrass	10	20	25	40	40	60	20	20	
Chickweed	30	40	70	65	100	50	50	50	
Cocklebur	35	90	100	100	100	40	70	70	
Corn	10	30	30	30	90	55	-	20	
Cotton	100	100	100	100	100	100	90	100	
Crabgrass	10	40	30	40	50	35	35	20	
Downy Brome	0	15	20	30	0	0	0	0	
Duck salad	-	0	0	0	0	0	0	45	
Giant foxtail	30	35	40	50	85	40	20	30	
Italn. Rygrass	0	0	15	20	45	0	10	0	
Johnsongrass	20	30	40	-	90	80	-	30	
Lambsquarter	80	95	75	100	100	90	65	70	
Morningglory	80	100	100	100	100	60	60	90	
Rape	60	90	95	-	100	95	75	100	
Redroot Pigweed	95	90	90	90	100	100	70	70	
Rice Japonica	-	25	25	25	70	15	0	35	
Soybean	70	50	40	50	100	40	50	70	
Speedwell	100	70	-	95	100	-	80	85	
Sugar beet	80	100	100	100	100	95	40	100	
Umbrella sedge	-	0	0	0	0	0	35	45	
Velvetleaf	100	100	100	100	100	100	90	100	
Watergrass	-	20	15	10	60	20	0	20	
Wheat	25	30	10	15	40	35	0	0	
Wild buckwheat	65	95	100	95	100	100	35	95	
Wild oat	35	25	20	25	30	40	20	25	

TABLE B	COMPOUND								
	Rate 31 g/ha	2	3	4	6	8	9	14	17
PREEMERGENCE									
Barley Igri	0	0	0	0	0	0	0	0	0
Barnyardgrass	0	20	0	20	95	10	0	10	
Bedstraw	0	0	10	0	100	40	0	30	
Blackgrass	0	0	0	0	10	0	0	0	
Chickweed	-	0	20	10	85	0	0	0	
Cocklebur	0	0	0	20	30	0	10	40	
Corn	0	0	0	0	30	0	0	0	
Cotton	0	0	0	0	100	10	0	100	
Crabgrass	0	0	0	0	75	0	0	10	
Downy Brome	0	0	0	0	0	0	0	0	
Giant foxtail	0	0	0	0	100	15	0	0	
Italn. Rygrass	0	0	0	0	35	0	0	0	
Johnsongrass	0	0	0	10	35	10	0	0	
Lambsquarter	20	85	85	95	100	100	0	70	
Morningglory	0	0	10	30	30	10	0	35	
Rape	0	0	0	0	75	0	0	40	
Redroot Pigweed	90	0	0	10	100	100	0	70	
Soybean	0	10	0	0	50	20	0	20	
Speedwell	-	0	50	25	100	10	0	10	
Sugar beet	0	0	0	0	85	50	0	90	
Velvetleaf	0	0	10	30	100	0	0	100	
Wheat	0	0	0	0	0	0	0	0	
Wild buckwheat	0	0	0	0	100	0	0	90	
Wild oat	0	0	0	0	30	20	0	0	

TABLE B	COMPOUND				
	Rate 16 g/ha	3	4	6	8
POSTEMERGENCE					
Barley Igri	30	30	30	30	10
Barnyardgr. (Flood)	25	20	25	35	20
Barnyardgrass	20	50	25	95	10
Bedstraw	70	85	-	100	65
Blackgrass	20	25	40	40	0
Chickweed	40	60	45	95	45
Cocklebur	90	100	90	75	50
Corn	20	20	25	50	10
Cotton	100	100	100	100	100
Crabgrass	25	30	20	35	15
Downy Brome	15	15	20	0	0
Duck salad	0	0	0	0	30
Giant foxtail	25	30	50	45	25
Italn. Rygrass	0	0	0	35	0
Johnsongrass	20	40	55	85	25
Lambsquarter	40	65	90	100	60
Morningglory	100	100	100	100	70
Rape	80	90	100	100	95
Redroot Pigweed	90	90	-	95	70
Rice Japonica	20	20	15	45	30
Soybean	40	35	40	100	50
Speedwell	70	85	95	100	-
Sugar beet	100	95	100	100	100
Umbrella sedge	0	0	0	0	20
Velvetleaf	100	100	100	100	100
Watergrass	20	15	10	40	20
Wheat	20	0	10	35	0
Wild buckwheat	90	100	95	100	95
Wild oat	25	20	25	20	10

TABLE B	COMPOUND				
	Rate 16 g/ha	3	4	6	8
PREEMERGENCE					
Barley Igri	0	0	0	0	0
Barnyardgrass	20	0	10	50	0
Bedstraw	0	0	0	20	30
Blackgrass	0	0	0	10	0
Chickweed	0	0	10	50	0
Cocklebur	0	0	0	20	20
Corn	0	0	0	25	0
Cotton	0	0	0	90	100
Crabgrass	0	0	0	60	0
Downy Brome	0	0	0	0	0
Giant foxtail	0	0	0	90	0
Italn. Rygrass	0	0	0	0	0
Johnsongrass	0	0	-	30	0
Lambsquarter	10	0	95	100	15
Morningglory	0	0	20	30	35
Rape	0	0	0	20	0
Redroot Pigweed	0	0	0	100	35
Soybean	0	0	0	30	20
Speedwell	0	0	25	100	0
Sugar beet	0	0	0	35	45
Velvetleaf	0	0	20	100	70
Wheat	0	0	0	0	0
Wild buckwheat	0	0	0	85	85
Wild oat	0	0	0	10	0

TABLE B		COMPOUND				TABLE B		COMPOUND			
Rate	8 g/ha	3	4	6	8	Rate	8 g/ha	3	4	6	8
POSTEMERGENCE						PREEMERGENCE					
Barley Igri		30	20	30	30	Barley Igri		0	0	0	0
Barnyardgr. (Flood)		15	15	20	0	Barnyardgrass		0	0	10	40
Barnyardgrass		20	40	20	45	Bedstraw		0	0	0	15
Bedstraw		55	85	100	70	Blackgrass		0	0	0	0
Blackgrass		20	25	35	20	Chickweed		0	0	10	30
Chickweed		40	60	30	70	Cocklebur		0	0	0	15
Cocklebur		80	90	90	65	Corn		0	0	0	0
Corn		15	15	20	30	Cotton		0	0	0	30
Cotton		100	100	100	100	Crabgrass		0	0	0	35
Crabgrass		25	20	20	20	Downy Brome		0	0	0	0
Downy Brome		10	0	10	0	Giant foxtail		0	0	0	35
Duck salad		0	0	0	0	Italn. Rygrass		0	0	0	0
Giant foxtail		25	25	35	30	Johnsongrass		0	0	0	10
Italn. Rygrass		0	0	0	10	Lambsquarter		0	0	20	100
Johnsongrass		20	-	55	35	Morningglory		0	0	10	20
Lambsquarter		40	50	75	95	Rape		0	0	0	0
Morningglory		100	100	100	90	Redroot Pigweed		0	0	0	90
Rape		80	30	100	75	Soybean		0	0	0	0
Redroot Pigweed		90	80	90	95	Speedwell		0	-	10	100
Rice Japonica		20	15	0	30	Sugar beet		0	0	0	-
Soybean		40	35	40	85	Velvetleaf		0	0	0	100
Speedwell		70	85	90	100	Wheat		0	0	0	0
Sugar beet		100	-	100	100	Wild buckwheat		0	0	0	30
Umbrella sedge		0	0	0	0	Wild oat		0	0	0	0
Velvetleaf		100	100	100	100						
Watergrass		20	15	10	0						
Wheat		10	0	10	30						
Wild buckwheat		85	85	90	100						
Wild oat		20	20	20	20						

TEST C

Seeds of barnyardgrass (*Echinochloa crus-galli*), bindweed (*Convolvulus arvensis*), black nightshade (*Solanum ptycanthum dunal*), cassia (*Cassia obtusifolia*),
5 cocklebur (*Xanthium pensylvanicum*), common ragweed (*Ambrosia artemisiifolia*), corn (*Zea mays*), cotton (*Gossypium hirsutum*), crabgrass (*Digitaria* spp.), fall panicum (*Panicum dichotomiflorum*), giant foxtail (*Setaria faberii*), green foxtail (*Setaria viridis*), jimsonweed (*Datura stramonium*), johnsongrass (*Sorghum halepense*), lambsquarter (*Chenopodium album*), morningglory (*Ipomoea* spp.), prickly sida (*Sida spinosa*), shattercane (*Sorghum vulgare*), signalgrass (*Brachiaria platyphylla*),
10 smartweed (*Polygonum pensylvanicum*), soybean (*Glycine max*), sunflower (*Helianthus annuus*), velvetleaf (*Abutilon theophrasti*), wild proso (*Panicum miliaceum*), woolly

cupgrass (*Eriochloa villosa*), yellow foxtail (*Setaria lutescens*) and purple nutsedge (*Cyperus rotundus*) tubers were planted into a matapeake sandy loam soil. These crops and weeds were grown in the greenhouse until the plants ranged in height from two to eighteen cm (one to four leaf stage), then treated postemergence with the test chemicals
5 formulated in a non-phytotoxic solvent mixture which includes a surfactant. Pots receiving preemergence treatments were planted immediately prior to test chemical application. Pots treated in this fashion were placed in the greenhouse and maintained according to routine greenhouse procedures.

Treated plants and untreated controls were maintained in the greenhouse
10 approximately 14-21 days after application of the test compound. Visual evaluations of plant injury responses were then recorded. Plant response ratings, summarized in Table C, are reported on a 0 to 100 scale where 0 is no effect and 100 is complete control.

TABLE C	COMPOUND
Rate 70 g/ha	4
POSTEMERGENCE	
Barnyardgrass	95
Bindweed	100
Blk Nightshade	100
Cassia	25
Cocklebur	70
Corn	35
Corn IR	35
Corn IT	35
Cotton	100
Crabgrass	20
Fall Panicum	20
Giant Foxtail	20
Green Foxtail	25
Jimsonweed	70
Johnson Grass	25
Lambsquarter	35
Morningglory	100
Nutsedge	15
Prickly Sida	100
Ragweed	100
Shattercane	45
Signalgrass	15
Smartweed	30
Soybean	45
Soybean 4-4	55
Soybean W20	45
Sunflower	90
Velvetleaf	100
Wild Proso	30
Woolly cupgrass	30
Yellow Foxtail	20

TABLE C	COMPOUND
Rate 35 g/ha	4
POSTEMERGENCE	
Barnyardgrass	65
Bindweed	75
Blk Nightshade	85
Cassia	20
Cocklebur	50
Corn	25
Corn IR	30
Corn IT	30
Cotton	100
Crabgrass	0
Fall Panicum	15
Giant Foxtail	15
Green Foxtail	15
Jimsonweed	55
Johnson Grass	15
Lambsquarter	20
Morningglory	100
Nutsedge	0
Prickly Sida	35
Ragweed	85
Shattercane	35
Signalgrass	15
Smartweed	25
Soybean	40
Soybean 4-4	45
Soybean W20	35
Sunflower	85
Velvetleaf	100
Wild Proso	20
Woolly cupgrass	25
Yellow Foxtail	15

TABLE C		COMPOUND	TABLE C		COMPOUND
Rate	17 g/ha	4	Rate	8 g/ha	4 14
POSTEMERGENCE			POSTEMERGENCE		
Barnyardgrass		25	Barnyardgrass	10	15
Bindweed		45	Bindweed	25	-
Blk Nightshade		65	Blk Nightshade	25	-
Cassia		15	Cassia	15	-
Cocklebur		35	Cocklebur	30	-
Corn		20	Corn	15	20
Corn IR		20	Corn IR	20	30
Corn IT		20	Corn IT	15	20
Cotton		75	Cotton	70	100
Crabgrass		0	Crabgrass	0	25
Fall Panicum		10	Fall Panicum	0	15
Giant Foxtail		10	Giant Foxtail	10	20
Green Foxtail		15	Green Foxtail	10	25
Jimsonweed		30	Jimsonweed	25	-
Johnson Grass		15	Johnson Grass	10	25
Lambsquarter		15	Lambsquarter	10	-
Morningglory		100	Morningglory	75	-
Nutsedge		0	Nutsedge	0	-
Prickly Sida		30	Prickly Sida	25	-
Ragweed		60	Ragweed	25	-
Shattercane		30	Shattercane	25	25
Signalgrass		10	Signalgrass	0	-
Smartweed		20	Smartweed	15	-
Soybean		25	Soybean	20	45
Soybean 4-4		35	Soybean 4-4	35	50
Soybean W20		30	Soybean W20	25	65
Sunflower		65	Sunflower	40	30
Velvetleaf		90	Velvetleaf	20	-
Wild Proso		15	Wild Proso	15	35
Woolly cupgrass		15	Woolly cupgrass	15	30
Yellow Foxtail		15	Yellow Foxtail	10	35

TABLE C	COMPOUND	
Rate 4 g/ha	4	14
POSTEMERGENCE		
Barnyardgrass	10	10
Bindweed	15	-
Blk Nightshade	10	-
Cassia	10	-
Cocklebur	25	-
Corn	10	15
Corn IR	10	20
Corn IT	15	10
Cotton	10	85
Crabgrass	0	10
Fall Panicum	0	10
Giant Foxtail	0	15
Green Foxtail	10	15
Jimsonweed	10	-
Johnson Grass	0	20
Lambsquarter	0	-
Morningglory	25	-
Nutsedge	0	-
Prickly Sida	20	-
Ragweed	15	-
Shattercane	25	10
Signalgrass	0	-
Smartweed	10	-
Soybean	15	25
Soybean 4-4	20	30
Soybean W20	15	35
Sunflower	10	25
Velvetleaf	10	-
Wild Proso	10	25
Woolly cupgrass	10	15
Yellow Foxtail	10	25

TABLE C	COMPOUND	
Rate 2 g/ha	14	
POSTEMERGENCE		
Barnyardgrass	0	
Bindweed	-	
Blk Nightshade	-	
Cassia	-	
Cocklebur	-	
Corn	10	
Corn IR	10	
Corn IT	0	
Cotton	75	
Crabgrass	10	
Fall Panicum	0	
Giant Foxtail	10	
Green Foxtail	0	
Jimsonweed	-	
Johnson Grass	0	
Lambsquarter	-	
Morningglory	-	
Nutsedge	-	
Prickly Sida	-	
Ragweed	-	
Shattercane	0	
Signalgrass	-	
Smartweed	-	
Soybean	10	
Soybean 4-4	10	
Soybean W20	10	
Sunflower	15	
Soybean	10	
Soybean 4-4	10	
Soybean W20	10	
Sunflower	15	
Velvetleaf	-	
Wild Proso	10	
Woolly cupgrass	10	
Yellow Foxtail	10	

TEST D

Compounds evaluated in this test were formulated in a non-phytotoxic solvent mixture which includes a surfactant and applied to the soil surface before plant seedlings emerged (preemergence application) and to plants that were in the one-to-four leaf stage (postemergence application). A sandy loam soil was used for the preemergence test while a mixture of sandy loam soil and greenhouse potting mix in a 60:40 ratio was used

for the postemergence test. Test compounds were applied within approximately one day after planting seeds for the preemergence test.

Plantings of these crops and weed species were adjusted to produce plants of appropriate size for the postemergence test. All plant species were grown using normal greenhouse practices. Crop and weed species include winter barley (*Hordeum vulgare* cv. 'Igri'), blackgrass (*Alopecurus myosuroides*), chickweed (*Stellaria media*), downy brome (*Bromus tectorum*), galium (*Galium aparine*), green foxtail (*Setaria viridis*), kochia (*Kochia scoparia*), lambsquarters (*Chenopodium album*), speedwell (*Veronica persica*), rape (*Brassica napus*), ryegrass (*Lolium multiflorum*), sugar beet (*Beta vulgaris* cv. 'US1'), spring wheat (*Triticum aestivum* cv. 'ERA'), windgrass (*Apera spica-venti*), winter wheat (*Triticum aestivum* cv. 'Talent'), wild buckwheat (*Polygonum convolvulus*), wild mustard (*Sinapis arvensis*), and wild oat (*Avena fatua*).

Treated plants and untreated controls were maintained in a greenhouse for approximately 21 to 28 days, after which all treated plants were compared to untreated controls and visually evaluated. Plant response ratings, summarized in Table D, are based upon a 0 to 100 scale where 0 is no effect and 100 is complete control. A dash response (-) means no test result.


TABLE D	COMPOUND	TABLE D	COMPOUND
Rate 12 g/ha	8	Rate 8 g/ha	8
POSTEMERGENCE		POSTEMERGENCE	
Blackgrass	10	Blackgrass	10
Chickweed	75	Chickweed	55
Downy brome	10	Downy brome	10
Galium	100	Galium	100
Green foxtail	15	Green foxtail	40
Kochia	100	Kochia	100
Lambsquarters	95	Lambsquarters	80
Rape	100	Rape	100
Ryegrass	10	Ryegrass	10
Speedwell	100	Speedwell	90
Sugar beet	100	Sugar beet	100
Wheat (Spring)	30	Wheat (Spring)	25
Wheat (Winter)	25	Wheat (Winter)	20
Wild buckwheat	100	Wild buckwheat	100
Wild mustard	100	Wild mustard	100
Wild oat	20	Wild oat	15
Windgrass	10	Windgrass	15
Winter Barley	20	Winter Barley	20

TABLE D	COMPOUND
Rate 4 g/ha	8
POSTEMERGENCE	
Blackgrass	10
Chickweed	50
Downy brome	5
Galium	70
Green foxtail	50
Kochia	95
Lambsquarters	70
Rape	85
Ryegrass	5
Speedwell	65
Sugar beet	90
Wheat (Spring)	10
Wheat (Winter)	15
Wild buckwheat	85
Wild mustard	100
Wild oat	10
Windgrass	10
Winter Barley	15

TABLE D	COMPOUND
Rate 2 g/ha	8
POSTEMERGENCE	
Blackgrass	5
Chickweed	15
Downy brome	0
Galium	50
Green foxtail	30
Kochia	40
Lambsquarters	50
Rape	-
Ryegrass	5
Speedwell	55
Sugar beet	-
Wheat (Spring)	10
Wheat (Winter)	10
Wild buckwheat	100
Wild mustard	25
Wild oat	15
Windgrass	0
Winter Barley	10

CLAIMS

1. A compound selected from Formula I and Formula II, or an agriculturally-suitable salt thereof,



5

wherein

X is a direct bond; O; S; NH; N(C₁-C₃ alkyl); N(C₁-C₃ haloalkyl); or N(allyl);

R¹ is H; F; Cl; or Br;

R² is H; F; Cl; Br; CF₃; nitro; or cyano;

10 R³ is H; C₁-C₈ alkyl; C₃-C₈ cycloalkyl; C₃-C₈ alkenyl; C₃-C₈ alkynyl; C₁-C₈ haloalkyl; C₂-C₈ alkoxyalkyl; C₃-C₈ alkoxyalkoxyalkyl; C₃-C₈ haloalkynyl; C₃-C₈ haloalkenyl; C₁-C₈ alkylsulfonyl; C₁-C₈ haloalkylsulfonyl; C₃-C₈ alkoxyalkonylalkyl; S(O)₂NH(C₁-C₈ alkyl); C(O)R⁶; or benzyl optionally substituted on the phenyl ring with R⁷;

15 R⁴ is H; C₁-C₃ alkyl; or halogen;

R⁵ is H; C₁-C₆ alkyl; C₁-C₆ haloalkyl; halogen; S(O)₂(C₁-C₆ alkyl); or C(=O)R⁸;
or

R⁴ and R⁵ are taken together along with the carbon to which they are attached to form a spiro-cyclopropane ring;


20 R⁶ is C₁-C₆ alkyl; C₁-C₆ haloalkyl; C₁-C₆ alkoxy; NH(C₁-C₆ alkyl); phenyl optionally substituted with R⁷; benzyl; or C₂-C₈ dialkylamino;

R⁷ is C₁-C₆ alkyl; 1-2 halogen; C₁-C₆ alkoxy; or CF₃;


R⁸ is H; C₁-C₆ alkyl; C₁-C₆ alkoxy; or NH(C₁-C₆ alkyl);

J is


25



J-1




J-2




J-3


89




J-4




J-5




J-6




J-7




J-8




J-9




J-10




J-11




J-12




J-13




J-14




J-15



J-16



J-17



J-18

wherein the dashed line in J-5, J-6 and J-12 indicates that the left-hand ring contains only single bonds or one bond in the ring is a carbon-carbon double bond;

5 n and m are each independently 0; 1; 2; or 3; provided that $m + n$ is 2 or 3;

Z is CR⁹R¹⁰; O; S; S(O); S(O)₂; N(C₁-C₄ alkyl); or $\overset{\oplus}{\text{N}} \begin{array}{l} \text{R} \\ \diagdown \\ \ominus \end{array}$;

each R⁹ is independently H; C₁-C₃ alkyl; halogen; hydroxy; C₁-C₆ alkoxy; C₁-C₆ haloalkyl; C₁-C₆ haloalkoxy; C₂-C₆ alkylcarbonyloxy; or C₂-C₆ haloalkylcarbonyloxy;

5 each R¹⁰ is independently H; C₁-C₃ alkyl; hydroxy; or halogen;

R¹¹ and R¹² are each independently H; halogen; C₁-C₆ alkyl; C₃-C₆ alkenyl; or C₁-C₆ haloalkyl;

R¹³ is H; C₁-C₆ alkyl; C₁-C₆ haloalkyl; C₃-C₆ alkenyl; C₃-C₆ haloalkenyl; C₃-C₆ alkynyl; C₃-C₆ haloalkynyl; HC(=O); (C₁-C₄ alkyl)C(=O); or NH₂;

10 R¹⁴ is C₁-C₆ alkyl; C₁-C₆ alkylthio; C₁-C₆ haloalkyl; CF₃; or N(CH₃)₂;

W is N or CR¹⁵;

R¹⁵ is H; C₁-C₆ alkyl; halogen; or phenyl optionally substituted with C₁-C₆ alkyl, 1-2 halogen, C₁-C₆ alkoxy, or CF₃;

each Q is independently O or S;

15 Q¹ is O or S;

Z¹ is CR¹⁶R¹⁷; O; S; S(O); S(O)₂; N(C₁-C₄ alkyl); or $\overset{\oplus}{\text{N}} \begin{array}{l} \text{R} \\ \diagdown \\ \ominus \end{array}$;

each R¹⁶ is independently H; halogen; hydroxy; C₁-C₆ alkoxy; C₁-C₆ haloalkyl; C₁-C₆ haloalkoxy; C₂-C₆ alkylcarbonyloxy; or C₂-C₆ haloalkylcarbonyloxy;

each R¹⁷ is independently H; hydroxy; or halogen; or

20 when R¹⁶ and R¹⁷ are bonded to adjacent atoms they can be taken together with

the carbons to which they are attached to form $\begin{array}{c} \text{CH}_2 \\ / \quad \backslash \\ \text{HC} - \text{CH} - \end{array}$;

R¹⁸ is C₁-C₆ alkyl; halogen; or C₁-C₆ haloalkyl;

R¹⁹ and R²⁰ are each independently H; C₁-C₆ alkyl; or C₁-C₆ haloalkyl;

25 R²¹ and R²² are each independently C₁-C₆ alkyl; C₁-C₆ haloalkyl; C₃-C₆ alkenyl; C₃-C₆ haloalkenyl; C₃-C₆ alkynyl; or C₃-C₆ haloalkynyl;

R²³ is halogen or cyano;

R²⁴ is C₁-C₆ alkylsulfonyl; C₁-C₆ alkyl; C₁-C₆ haloalkyl; C₃-C₆ alkenyl; C₃-C₆ alkynyl; C₁-C₆ alkoxy; C₁-C₆ haloalkoxy; or halogen;

R²⁵ is C₁-C₆ alkyl; C₁-C₆ haloalkyl; C₃-C₆ alkenyl; or C₃-C₆ alkynyl; and

30 R²⁶ is C₁-C₆ alkyl; C₁-C₆ haloalkyl; or phenyl optionally substituted with C₁-C₆ alkyl, 1-2 halogen, 1-2 nitro, C₁-C₆ alkoxy, or CF₃;

provided that when X is O or S in Formula II, then J is other than J-6 or J-7.

2. A compound of Claim 1 wherein:

X is a direct bond or O;

R¹ is F or Cl;

R² is F; Cl; or Br;

5 R³ is H; C₁-C₆ alkyl; C₃-C₆ cycloalkyl; C₃-C₆ alkenyl; C₃-C₆ alkynyl; C₁-C₆ haloalkyl; C₂-C₆ alkoxyalkyl; C₃-C₆ haloalkenyl; C₁-C₆ alkylsufonyl; C₃-C₆ alkoxycarbonylalkyl; C(O)R⁶; or benzyl optionally substituted on the phenyl ring with R⁷;

R⁴ is H or halogen;

10 R⁵ is H;

J is J-1; J-2; J-3; J-4; J-5; J-6; J-8; J-12; J-15; or J-16;

Z is CR⁹R¹⁰; O; S; or N(C₁-C₄ alkyl);

each R⁹ is independently H; halogen; or C₁-C₆ haloalkoxy;

each R¹⁰ is independently H or halogen;

15 each Q is O;


Q¹ is S;

Z¹ is CR¹⁶R¹⁷; O; S; or N(C₁-C₄ alkyl);

each R¹⁶ is independently H; halogen; or haloalkoxy;

each R¹⁷ is independently H or halogen; or

20 when R¹⁶ and R¹⁷ are bonded to adjacent atoms they can be taken together with the carbons to which they are attached to form



R¹⁸ is *t*-butyl or CH₂CH₂CH₂F;

R²³ is Cl or cyano; and

25 R²⁴ is C₁-C₄ haloalkyl or C₁-C₆ haloalkoxy.

3. A compound of Claim 2 wherein:

R² is F or Cl;

30 R³ is C₁-C₆ alkyl; C₃-C₆ cycloalkyl; C₃-C₆ alkenyl; C₃-C₆ alkynyl; C₃-C₆ alkoxycarbonylalkyl; or benzyl optionally substituted on the phenyl ring with R⁷;

J is J-1; J-2; J-3; J-4; J-5; J-6; J-8; or J-12;

Z is CR⁹R¹⁰ or O;

each R⁹ is independently H or halogen;

R¹¹ is C₁-C₄ haloalkyl;

35 R¹² is C₁-C₄ alkyl;

R¹³ is CH₃;

R¹⁴ is CF₃;


W is CH;

Z¹ is CR¹⁶R¹⁷ or O;

each R¹⁶ is independently H or halogen;

each R¹⁷ is independently H or halogen; or

- 5 when R¹⁶ and R¹⁷ are bonded to adjacent atoms they can be taken together with the carbons to which they are attached to form



4. A compound of Claim 3 wherein:

R² is Cl;

- 10 R³ is C₁-C₆ alkyl; C₃-C₆ alkenyl; or C₃-C₆ alkynyl;

J is J-1; J-2; J-3; J-4; J-6; or J-12;

Z is CR⁹R¹⁰;

R⁹ is independently H or F;

R¹⁰ is independently H or F;

- 15 R¹¹ is CF₂H;

R¹² is CH₃;

Z¹ is CR¹⁶R¹⁷;

R¹⁶ is independently H or F; and

R¹⁷ is independently H or F.

- 20 5. A compound of Claim 1 of Formula I.

6. A compound of Claim 2 which is selected from the group:

3-[7-chloro-5-fluoro-1,3-dihydro-1-[(4-methoxyphenyl)methyl]-2,1-benzisothiazol-4-yl]-1-methyl-6-(trifluoromethyl)-2,4(1*H*,3*H*)-pyrimidinedione *S,S*-dioxide;

- 25 3-(7-chloro-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl)-1-methyl-6-(trifluoromethyl)-2,4(1*H*,3*H*)-pyrimidinedione *S,S*-dioxide;

3-(7-chloro-1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl)-1-methyl-6-(trifluoromethyl)-2,4(1*H*,3*H*)-pyrimidinedione *S,S*-dioxide; and



- 30 3-[(7-chloro-1-ethyl-5-fluoro-1,3-dihydro-2,1-benzisothiazol-4-yl)imino]tetrahydro-1*H*,3*H*-[1,3,4]thiadiazolo[3,4-*a*]pyridazin-1-one *S,S*-dioxide.

7. A herbicidal composition comprising an effective amount of a compound according to Claim 1 and at least one of a surfactant, a solid diluent or a liquid diluent.

8. A herbicidal composition comprising an effective amount of a compound according to Claim 2 and at least one of a surfactant, a solid diluent or a liquid diluent.
- 35

9. A method for controlling the growth of undesired vegetation comprising contacting the vegetation or its environment with a herbicidally effective amount of a composition of Claim 7.

10. A compound selected from the Formula 1 and 2:



5

wherein

X is a direct bond; O; S; NH; N(C₁-C₃ alkyl); N(C₁-C₃ haloalkyl); or N(allyl);

R¹ is H; F; Cl; or Br;

R² is H; F; Cl; Br; CF₃; nitro; or cyano;

10 R³ is H; C₁-C₈ alkyl; C₃-C₈ cycloalkyl; C₃-C₈ alkenyl; C₃-C₈ alkynyl; C₁-C₈ haloalkyl; C₂-C₈ alkoxyalkyl; C₃-C₈ alkoxyalkoxyalkyl; C₃-C₈ haloalkynyl; C₃-C₈ haloalkenyl; C₁-C₈ alkylsulfonyl; C₁-C₈ haloalkylsulfonyl; C₃-C₈ alkoxyalkonylalkyl; (C₁-C₈ alkyl)NHS(O)₂; C(O)R⁶; or benzyl optionally substituted on the phenyl ring with R⁷;

15 R⁴ is H; C₁-C₃ alkyl; or halogen;

R⁵ is H; C₁-C₆ alkyl; C₁-C₆ haloalkyl; halogen; S(O)₂(C₁-C₆ alkyl); or C(=O)R⁸;

or

R⁴ and R⁵ are taken together along with the carbon to which they are attached to form a spiro-cyclopropane ring;

20 R⁶ is C₁-C₆ alkyl; C₁-C₆ haloalkyl; C₁-C₆ alkoxy; NH(C₁-C₆ alkyl); phenyl optionally substituted with R⁷; benzyl; or C₂-C₈ dialkylamino;

R⁷ is C₁-C₆ alkyl; 1-2 halogen; C₁-C₆ alkoxy; or CF₃; and

R⁸ is H; C₁-C₆ alkyl; C₁-C₆ alkoxy; or NH(C₁-C₆ alkyl).

INTERNATIONAL SEARCH REPORT

Intern. al Application No
PCT/US 95/06706

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 6 C07D417/04 A01N43/80 C07D487/04 A01N43/824 C07D513/04
 C07D471/04 C07D275/06 C07D291/08 //(C07D471/04, 235:00,
 221:00), (C07D487/04, 249:00, 237:00), (C07D513/04, 285:00, 237:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US,A,4 906 281 (JUN H. CHANG) 6 March 1990 see the whole document ---	1,7-9
A	EP,A,0 468 924 (CIBA-GEIGY AG) 29 January 1992 see claims ---	1,7-9
A	WO,A,87 03782 (FMC CORPORATION) 2 July 1987 cited in the application see claims --- -/--	1,7-9

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

28 August 1995

Date of mailing of the international search report

- 5, 09. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+ 31-70) 340-3016

Authorized officer

Henry, J

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 95/06706

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CHEMICAL ABSTRACTS, vol. 110, no. 23, 5 June 1989 Columbus, Ohio, US; abstract no. 212837t, page 756; see abstract	1,7-9
A	& JP,A,63 222 167 (NISSAN CHEMICAL INDUSTRIES ,LTD) 16 September 1988 cited in the application -----	1,7-9

1

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 95/06706

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-4906281	06-03-90	NONE	
EP-A-468924	29-01-92	AU-B- 638854	08-07-93
		AU-A- 8126191	30-01-92
		JP-A- 4234360	24-08-92
		US-A- 5180418	19-01-93
WO-A-8703782	02-07-87	DE-A- 3688911	23-09-93
		DE-T- 3688911	09-12-93
		EP-A, B 0294375	14-12-88
		JP-B- 5033951	20-05-93
		JP-T- 62502896	19-11-87
		US-A- 5294595	15-03-94
		US-A- 5174809	29-12-92
		US-A- 5214154	25-05-93
		US-A- 4818275	04-04-89