



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁷ : C12N 15/82, 15/29, A01H 5/00</p>	A1	<p>(11) International Publication Number: WO 00/12733</p> <p>(43) International Publication Date: 9 March 2000 (09.03.00)</p>
<p>(21) International Application Number: PCT/US99/19604</p> <p>(22) International Filing Date: 25 August 1999 (25.08.99)</p> <p>(30) Priority Data: 60/098,230 28 August 1998 (28.08.98) US</p> <p>(71) Applicant: PIONEER HI-BRED INTERNATIONAL, INC. [US/US]; 800 Capital Square, 400 Locust Street, Des Moines, IA 50309 (US).</p> <p>(72) Inventors: MARTINO-CATT, Susan, J.; 499 N.W. 69th Place, Ankeny, IA 50021 (US). LAPPEGARD, Kathryn, K.; 1228 4th Street, Nevada, IA 50201 (US). OLSEN, Odd-Arne; Taarnveien 16, N-1430 Aas (NO). LINNESTAD, Casper; Ovre Linnestad, N-1540 Vestby (NO). ABBITT, Shane, E.; Apartment H, 919 S.E. Delaware Avenue, Ankeny, IA 50021 (US).</p> <p>(74) Agents: MICHEL, Marianne, H. et al.; Darwin Building, 7100 N.W. 62nd Avenue, Johnston, IA 50131-1000 (US).</p>		<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: SEED-PREFERRED PROMOTERS FROM <i>END</i> GENES</p>		
<p>(57) Abstract</p> <p>The present invention provides compositions and methods for regulating expression of heterologous nucleotide sequences in a plant. Compositions are novel nucleotide sequences for seed-preferred promoters isolated from genes for <i>end1</i> and <i>end2</i>. A method for expressing a heterologous nucleotide sequence in a plant using the promoter sequences disclosed herein is provided. The method comprises transforming a plant cell to comprise a heterologous nucleotide sequence operably linked to one of the constitutive promoters of the present invention and regenerating a stably transformed plant from the transformed plant cell.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

SEED-PREFERRED PROMOTERS FROM *END* GENES

FIELD OF THE INVENTION

The present invention relates to the field of plant molecular biology, more particularly to regulation of gene expression in plants.

BACKGROUND OF THE INVENTION

Expression of heterologous DNA sequences in a plant host is dependent upon the presence of an operably linked promoter that is functional within the plant host. Choice of the promoter sequence will determine when and where within the organism the heterologous DNA sequence is expressed. Where continuous expression is desired throughout the cells of a plant, constitutive promoters are utilized. In contrast, where gene expression in response to a stimulus is desired, inducible promoters are the regulatory element of choice. Where expression in specific tissues or organs are desired, tissue-specific promoters may be used. That is, they may drive expression in specific tissues or organs. Such tissue-specific promoters may be constitutive or inducible. In either case, additional regulatory sequences upstream and/or downstream from the core promoter sequence may be included in expression constructs of transformation vectors to bring about varying levels of expression of heterologous nucleotide sequences in a transgenic plant.

Frequently it is desirable to have constitutive or inducible expression of a DNA sequence in particular tissues or organs of a plant. For example, increased nutritional value of a plant might be accomplished by genetic manipulation of the plant's genome to comprise a seed-preferred promoter operably linked to a heterologous gene such that proteins with enhanced amino acid content are produced in the seed of the plant.

Alternatively, it might be desirable to inhibit expression of a native DNA sequence within a plant's tissues to achieve a desired phenotype. In this case, such inhibition might be accomplished with transformation of the plant to comprise a tissue-specific promoter operably linked to an antisense nucleotide sequence, such that constitutive expression of the antisense sequence produces an RNA transcript that interferes with translation of the mRNA of the native DNA sequence.

Seed development involves embryogenesis and maturation events as well as physiological adaptation processes that occur within the seed to insure progeny survival. Developing plant seeds accumulate and store carbohydrate, lipid, and protein that are subsequently used during germination. Expression of storage protein genes in seeds occurs primarily in the embryonic axis and cotyledons and in the endosperm of developing seeds but never in mature vegetative tissues. Generally, the expression patterns of seed proteins are highly regulated. This regulation includes spatial and temporal regulation during seed development. A variety of proteins accumulate and decay during embryogenesis and seed development and provide an excellent system for investigating different aspects of gene regulation as well as for providing regulatory sequences for use in genetic manipulation of plants.

Thus, isolation and characterization of seed-preferred promoters that can serve as regulatory regions for expression of heterologous nucleotide sequences of interest in a seed-preferred manner are needed for genetic manipulation of plants.

SUMMARY OF THE INVENTION

It is an object of the present invention to provide a novel nucleotide sequence for modulating gene expression in a plant.

It is a further object of the present invention to provide an isolated promoter capable of driving transcription in a seed-preferred manner.

It is a further object of the present invention to provide a method of improved control of an endogenous or exogenous product in the seed of a transformed plant.

It is a further object of the present invention to provide a method for providing useful changes in the phenotype of a seed of a transformed plant.

It is a further object of the present invention to provide a method for producing a novel product in the seed of a transformed plant.

It is a further object of the present invention to provide a method for producing a novel function in the seed of a transformed plant.

Therefore, in one aspect, the present invention relates to an isolated nucleic acid comprising an isolated promoter that is capable of driving transcription

in a seed-preferred manner, wherein said promoter comprises a nucleotide sequence selected from the group consisting of:

- a) sequences capable of driving expression of coding regions selected from the group consisting of coding regions for *end1* or *end2*;
- b) a sequence comprising at least 40 contiguous nucleotides of the sequence set forth in either of SEQ ID NOS:1 or 4;
- c) a sequence comprising a variant or fragment of the nucleotide sequence set forth in either of SEQ ID NOS: 1 or 4;
- d) the nucleotide sequences set forth in SEQ ID NOS: 1 or 4;
- e) a sequence that hybridizes to any one of SEQ ID NOS: 1 or 4 under low stringency conditions; and
- f) a nucleotide sequence complementary to a nucleotide sequence of (a) through (e).

In other aspects, the present invention relates to expression cassettes comprising the promoter operably linked to a nucleotide sequence, vectors containing the expression cassette, and plants stably transformed with at least one expression cassette.

In a further aspect, the present invention relates to a method for modulating expression in the seed of a stably transformed plant comprising the steps of (a) transforming a plant cell with an expression cassette comprising the promoter of the present invention operably linked to at least one nucleotide sequence; (b) growing the plant cell under plant growing conditions and (c) regenerating a stably transformed plant from the plant cell wherein expression of the nucleotide sequence alters the phenotype of the seed.

Compositions and methods for regulating expression of heterologous nucleotide sequences in a plant are provided. Compositions are novel nucleotide sequences for seed-preferred plant promoters, more particularly transcriptional initiation regions isolated from the plant genes *end1* and *end2*. A method for expressing a heterologous nucleotide sequence in a plant using the transcriptional initiation sequences disclosed herein is provided. The method comprises transforming a plant cell with a transformation vector that comprises a heterologous nucleotide sequence operably linked to one of the plant promoters of the present invention and regenerating a stably transformed plant from the transformed plant cell. In this manner, the promoter sequences are useful for

controlling the expression of endogenous as well as exogenous products in a seed-preferred manner.

Downstream from and under the transcriptional initiation regulation of the seed-specific region will be a sequence of interest which will provide for modification of the phenotype of the seed. Such modification includes modulating the production of an endogenous product, as to amount, relative distribution, or the like, or production of an exogenous expression product to provide for a novel function or product in the seed.

DETAILED DESCRIPTION OF THE INVENTION

In accordance with the invention nucleotide constructs are provided that allow initiation of transcription in seed. Constructs of the invention comprise regulated transcriptional initiation regions associated with seed formation and seed tissues. Thus, the compositions of the present invention comprise novel nucleotide sequences for plant promoters, particularly seed-preferred promoters, more particularly endosperm specific promoters, for the genes *end1* and *end2*. The *end1* promoter drives expression in transfer cells at an early stage in precursor cells and continues expression into mature cells. The *end2* promoter drives expression in aleurone cells.

The promoters for these genes may be isolated from the 5' untranslated region flanking their respective transcription initiation sites. Methods for isolation of promoter regions are well known in the art.

The term "isolated" refers to material, such as a nucleic acid, which is: (1) substantially or essentially free from components which normally accompany or interact with it as found in its naturally occurring environment; the isolated material optionally comprises material not found with the material in its natural environment; or (2) if the material is in its natural environment, the material has been synthetically (non-naturally) altered or produced by deliberate human intervention to a composition and/or placed at a locus in the cell (e.g., genome or subcellular organelle) not native to a material found in that environment. The alteration to yield the synthetic material can be performed on the material within or removed from its natural state.

Methods are readily available in the art for the hybridization of nucleic acid sequences. Promoter sequences from other plants may be isolated according to

well-known techniques based on their sequence homology to the promoter sequences set forth herein. In these techniques, all or part of the known promoter sequence is used as a probe which selectively hybridizes to other sequences present in a population of cloned genomic DNA fragments (i.e. genomic libraries) from a chosen organism.

For example, the entire promoter sequence or portions thereof may be used as probes capable of specifically hybridizing to corresponding promoter sequences. To achieve specific hybridization under a variety of conditions, such probes include sequences that are unique and are preferably at least about 10 nucleotides in length, and most preferably at least about 20 nucleotides in length. Such probes may be used to amplify corresponding promoter sequences from a chosen organism by the well-known process of polymerase chain reaction (PCR). This technique may be used to isolate additional promoter sequences from a desired organism or as a diagnostic assay to determine the presence of the promoter sequence in an organism.

Such techniques include hybridization screening of plated DNA libraries (either plaques or colonies; see e.g. Innis *et al.* (1990) *PCR Protocols, A Guide to Methods and Applications*, eds., Academic Press).

The terms "stringent conditions" or "stringent hybridization conditions" includes reference to conditions under which a probe will hybridize to its target sequence, to a detectably greater degree than other sequences (e.g., at least 2-fold over background). Stringent conditions are sequence-dependent and will be different in different circumstances. By controlling the stringency of the hybridization and/or washing conditions, target sequences can be identified which are 100% complementary to the probe (homologous probing). Alternatively, stringency conditions can be adjusted to allow some mismatching in sequences so that lower degrees of similarity are detected (heterologous probing). Generally, a probe is less than about 1000 nucleotides in length, preferably less than 500 nucleotides in length.

Typically, stringent conditions will be those in which the salt concentration is less than about 1.5 M Na ion, typically about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short probes (e.g., 10 to 50 nucleotides) and at least about 60°C for long probes

(e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide.

For purposes of defining the invention preferably low stringency conditions are employed including hybridization with a buffer solution of 30 to 35% formamide, 1 M NaCl, 1% SDS (sodium dodecyl sulphate) at 37°C, and a wash in 1X to 2X SSC (20X SSC = 3.0 M NaCl/0.3 M trisodium citrate) at 50 to 55°C. More preferably moderate stringency conditions are employed including hybridization in 40 to 45% formamide, 1 M NaCl, 1% SDS at 37°C, and a wash in 0.5X to 1X SSC at 55 to 60°C. Most preferably high stringency conditions are employed including hybridization in 50% formamide, 1 M NaCl, 1% SDS at 37°C, and a wash in 0.1X SSC at 60 to 65°C. Hybridization times are not critical and can range from about four hours to about sixteen hours.

An extensive guide to the hybridization of nucleic acids is found in Tijssen, *Laboratory Techniques in Biochemistry and Molecular Biology--Hybridization with Nucleic Acid Probes*, Part I, Chapter 2 "Overview of principles of hybridization and the strategy of nucleic acid probe assays", Elsevier, New York (1993); and *Current Protocols in Molecular Biology*, Chapter 2, Ausubel, *et al.*, Eds., Greene Publishing and Wiley-Interscience, New York (1995). See Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (2nd ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.) In general, sequences that correspond to promoter sequences of the invention and hybridize to the promoter sequence disclosed herein will be at least 50% homologous, preferably 60%, 70%, 80%, 85%, 90%, and even 95% homologous or more with the disclosed sequences.

The promoter regions of the invention may be isolated from any plant, including, but not limited to corn (*Zea mays*), (*Brassica napus*, *Brassica rapa ssp.*), alfalfa (*Medicago sativa*), rice (*Oryza sativa*), rye (*Secale cereale*), sorghum (*Sorghum bicolor*, *Sorghum vulgare*), sunflower (*Helianthus annuus*), wheat (*Triticum aestivum*), soybean (*Glycine max*), tobacco (*Nicotiana tabacum*), potato (*Solanum tuberosum*), peanuts (*Arachis hypogaea*), cotton (*Gossypium hirsutum*), sweet potato (*Ipomoea batatas*), cassava (*Manihot esculenta*), coffee (*Cofea spp.*), coconut (*Cocos nucifera*), pineapple (*Ananas comosus*), citrus trees (*Citrus spp.*), cocoa (*Theobroma cacao*), tea (*Camellia sinensis*), banana (*Musa spp.*), avocado (*Persea americana*), fig (*Ficus casica*), guava (*Psidium guajava*), mango (*Mangifera indica*), olive (*Olea europaea*), oats, barley, vegetables, ornamentals,

and conifers. Preferably plants include corn, soybean, sunflower, safflower, oil seed *Brassica*, wheat, rice, barley, rye, alfalfa, and sorghum.

The coding sequence expressed by the promoters of the invention may be used for varying the phenotype of the seeds. Various changes in phenotype are of interest including modifying the fatty acid composition in seeds, altering the starch or carbohydrate profile, altering the amino acid content of the seed, and the like. These results can be achieved by providing expression of heterologous or increased expression of endogenous products in seeds. Alternatively, the results can be achieved by providing for a reduction of expression of one or more endogenous products, particularly enzymes or cofactors in the seed. These changes result in a change in phenotype of the transformed seed.

Genes of interest include, generally, those involved in oil, starch, protein, carbohydrate or nutrient metabolism as well as those affecting kernel size, sucrose loading, and the like. In particular, *end1* may find use in regulating the influx of nutrients. Both promoters are useful in disease resistance and in regulating expression of phytate genes particularly to lower phytate levels in the seed.

General categories of genes of interest for the purpose of present invention include for example, those genes involved in information, such as Zinc fingers, those involved in communication, such as kinases, and those involved in housekeeping, such as heat shock proteins. More specific categories of transgenes, for example, include genes encoding important traits for agronomics, insect resistance, disease resistance, herbicide resistance, and grain characteristics. It is recognized that any gene of interest can be operably linked to the promoter of the invention and expressed in the seed.

Important traits such as oil, starch and protein content can be genetically altered in addition to using traditional breeding methods. Modifications include altering the content of oleic acid, saturated and unsaturated oils, increasing levels of lysine and sulfur-containing amino acids and providing other essential amino acids, and also modification of starch. Hordothionin protein modifications are described in WO94/16078; WO96/38562; WO96/08220; and U.S. Pat. No. 5,703,409 issued December 30, 1997, the disclosures of which are incorporated herein in their entirety by reference. Another example is lysine and/or sulfur rich seed protein encoded by the soybean 2S albumin described in WO97/35023, and

the chymotrypsin inhibitor from barley, Williamson *et al.* (1987) *Eur. J. Biochem.* 165:99-106, the disclosures of each are incorporated by reference. Derivatives of the following genes can be made by site directed mutagenesis to increase the level of preselected amino acids in the encoded polypeptide. For example, the gene encoding the barley high lysine polypeptide (BHL), is derived from barley chymotrypsin inhibitor, WO98/20133, incorporated herein by reference. Other proteins include methionine-rich plant proteins such as from sunflower seed (Lilley *et al.* (1989) *Proceedings of the World Congress on Vegetable Protein Utilization in Human Foods and Animal Feedstuffs*; Applewhite, H. (ed.); American Oil Chemists Soc., Champaign, IL:497-502, incorporated herein in its entirety by reference), corn (Pedersen *et al.* (1986) *J. Biol. Chem.* 261:6279; Kirihara *et al.* (1988) *Gene* 71:359, both incorporated herein in its entirety by reference) and rice (Musumura *et al.* (1989) *Plant Mol. Biol.* 12:123, incorporated herein in its entirety by reference). Other agronomically important genes encode latex, Flourey 2, growth factors, seed storage factors and transcription factors.

The quality of grain is reflected in traits such as levels and types of oils, saturated and unsaturated, quality and quantity of essential amino acids, and levels of cellulose. In corn, modified hordeothionin proteins, described in WO94/16078; WO96/38562; WO96/08220; and U.S. Pat. No. 5,703,409; provide descriptions of modifications of proteins for desired purposes.

Commercial traits can also be encoded on a gene(s) which could alter or increase for example, starch for the production of paper, textiles, and ethanol, or provide expression of proteins with other commercial uses. Another important commercial use of transformed plants is the production of polymers and bioplastics such as described in U.S. Patent No. 5,602,321 issued February 11, 1997. Genes such as B-ketothiolase, PHBase (polyhydroxybutyrate synthase) and acetoacetyl-CoA reductase (see Schubert *et al.* (1988) *J. Bacteriol* 170(12):5837-5847) facilitate expression of polyhydroxyalkanoates (PHAs).

Exogenous products include plant enzymes and products as well as those from other sources including prokaryotes and other eukaryotes. Such products include enzymes, cofactors, hormones, and the like. The level of seed proteins, particularly modified seed proteins having improved amino acid distribution to improve the nutrient value of the seed can be increased. This is achieved by the expression of such proteins having enhanced amino acid content.

Insect resistance genes may encode resistance to pests that have great yield drag such as rootworm, cutworm, European Corn Borer, and the like. Such genes include, for example, *Bacillus thuringiensis* endotoxin genes (U.S. Pat. Nos. 5,366,892; 5,747,450; 5,737,514; 5,723,756; 5,593,881; Geiser *et al.* (1986) *Gene* 48:109); lectins (Van Damme *et al.* (1994) *Plant Mol. Biol.* 24:825); and the like.

Genes encoding disease resistance traits may include detoxification genes, such as against fumonisin (U.S. Patent Application No. 08/484,815 filed June 7, 1995); avirulence (*avr*) and disease resistance (R) genes (Jones *et al.* (1994) *Science* 266:789; Martin *et al.* (1993) *Science* 262:1432; Mindrinis *et al.* (1994) *Cell* 78:1089; and the like.

Agronomic traits in seeds can be improved by altering expression of genes that affect the response of seed growth and development during environmental stress, Cheikh-N *et al.* (1994) *Plant Physiol.* 106(1):45-51) and genes controlling carbohydrate metabolism to reduce kernel abortion in maize, Zinselmeier *et al.* (1995) *Plant Physiol.* 107(2):385-391.

As noted, the heterologous nucleotide sequence operably linked to one of the promoters disclosed herein may be an antisense sequence for a targeted gene. By "antisense DNA nucleotide sequence" is intended a sequence that is in inverse orientation to the 5'-to-3' normal orientation of that nucleotide sequence. When delivered into a plant cell, expression of the antisense DNA sequence prevents normal expression of the DNA nucleotide sequence for the targeted gene. The antisense nucleotide sequence encodes an RNA transcript that is complementary to and capable of hybridizing to the endogenous messenger RNA (mRNA) produced by transcription of the DNA nucleotide sequence for the targeted gene. In this case, production of the native protein encoded by the targeted gene is inhibited to achieve a desired phenotypic response. Thus the promoter sequences disclosed herein may be operably linked to antisense DNA sequences to reduce or inhibit expression of a native protein in the plant seed.

By "promoter" or "transcriptional initiation region" is intended a regulatory region of DNA usually comprising a TATA box capable of directing RNA polymerase II to initiate RNA synthesis at the appropriate transcription initiation site for a particular coding sequence. A promoter may additionally comprise other recognition sequences generally positioned upstream or 5' to the TATA box, referred to as upstream promoter elements, which influence the transcription

initiation rate. It is recognized that having identified the nucleotide sequences for the promoter regions disclosed herein, it is within the state of the art to isolate and identify further regulatory elements in the 5' untranslated region upstream from the particular promoter regions identified herein. Thus the promoter regions disclosed herein are generally further defined by comprising upstream regulatory elements such as those responsible for tissue and temporal expression of the coding sequence, enhancers and the like. In the same manner, the promoter elements which enable expression in the desired tissue such as the seed can be identified, isolated, and used with other core promoters to confirm seed-preferred expression.

The regulatory sequences of the present invention, when operably linked to a heterologous nucleotide sequence of interest and inserted into a transformation vector, enable seed-preferred expression of the heterologous nucleotide sequence in the seeds of a plant stably transformed with this vector.

By "seed-preferred" is intended expression in the seed, including at least one of embryo, kernel, pericarp, endosperm, nucellus, aleurone, pedicel, and the like.

By "heterologous nucleotide sequence" is intended a sequence that is not naturally occurring with the promoter sequence. While this nucleotide sequence is heterologous to the promoter sequence, it may be homologous, or native, or heterologous, or foreign, to the plant host.

It is recognized that the promoters may be used with their native coding sequences to increase or decrease expression resulting in a change in phenotype in the transformed seed.

The isolated promoter sequences of the present invention can be modified to provide for a range of expression levels of the heterologous nucleotide sequence. Thus, less than the entire promoter regions may be utilized and the ability to drive seed-preferred expression retained. However, it is recognized that expression levels of mRNA may be decreased with deletions of portions of the promoter sequences. Generally, at least about 20 nucleotides of an isolated promoter sequence will be used to drive expression of a nucleotide sequence.

It is recognized that to increase transcription levels enhancers may be utilized in combination with the promoter regions of the invention. Enhancers are nucleotide sequences that act to increase the expression of a promoter region.

Enhancers are known in the art and include the SV40 enhancer region, the 35S enhancer element, and the like.

Modifications of the isolated promoter sequences of the present invention can provide for a range of expression of the heterologous nucleotide sequence. Thus, they may be modified to be weak promoters or strong promoters. Generally, by "weak promoter" is intended a promoter that drives expression of a coding sequence at a low level. By "low level" is intended at levels of about 1/10,000 transcripts to about 1/100,000 transcripts to about 1/500,000 transcripts. Conversely, a strong promoter drives expression of a coding sequence at a high level, or at about 1/10 transcripts to about 1/00 transcripts to about 1/1,000 transcripts.

The nucleotide sequences for the promoters of the present invention may be the naturally occurring sequences or sequences having substantial homology. By "substantial homology" is intended a sequence exhibiting substantial functional and structural equivalence with the naturally occurring sequence. Any structural differences between substantially homologous sequences do not effect the ability of the sequence to function as a promoter as disclosed in the present invention. Thus, sequences having substantial sequence homology with the sequence of a particular seed-preferred promoter of the present invention will direct seed-preferred expression of an operably linked heterologous nucleotide sequence. Two promoter nucleotide sequences are considered substantially homologous when they have at least about 70%, preferably at least about 80%, more preferably at least about 90%, still more preferably at least about 95% sequence homology. Substantially homologous sequences of the present invention include variants of the disclosed sequences such as those that result from site-directed mutagenesis, as well as synthetically derived sequences.

Substantially homologous sequences of the present invention also refer to those fragments of a particular promoter nucleotide sequences disclosed herein that operate to promote the seed-preferred expression of an operably linked heterologous nucleotide sequence. These fragments will comprise at least about 20 contiguous nucleotides, preferably at least about 50 contiguous nucleotides, more preferably at least about 75 contiguous nucleotides, even more preferably at least about 100 contiguous nucleotides of the particular promoter nucleotide sequence disclosed herein. The nucleotides of such fragments will usually

comprise the TATA recognition sequence of the particular promoter sequence. Such fragments may be obtained by use of restriction enzymes to cleave the naturally occurring promoter nucleotide sequences disclosed herein; by synthesizing a nucleotide sequence from the naturally occurring promoter DNA sequence; or may be obtained through the use of PCR technology. See particularly, Mullis *et al.* (1987) *Methods Enzymol.* 155:335-350, and Erlich, ed. (1989) *PCR Technology* (Stockton Press, New York). Again, variants of these promoter fragments, such as those resulting from site-directed mutagenesis, are encompassed by the compositions of the present invention.

Nucleotide sequences comprising at least about 40 contiguous sequences of the sequences set forth in SEQ ID NOS: 1 and 4 are encompassed. These sequences may be isolated by hybridization, PCR, and the like. Such sequences encompass fragments capable of driving seed-preferred expression, fragments useful as probes to identify similar sequences, as well as elements responsible for temporal or tissue specificity. Biologically active variants of the promoter sequences are also encompassed by the method of the present invention. Such variants should retain promoter activity, particularly the ability to drive expression in seed or seed tissues. Biologically active variants include, for example, the native promoter sequences of the invention having one or more nucleotide substitutions, deletions or insertions. Promoter activity may be measured by Northern blot analysis, reporter activity measurements when using transcriptional fusions, and the like. See, for example, Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (2nd ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.), herein incorporated by reference.

The following terms are used to describe the sequence relationships between two or more nucleic acids or polynucleotides: (a) "reference sequence", (b) "comparison window", (c) "percentage of sequence identity", and (d) "substantial identity".

(a) As used herein, "reference sequence" is a defined sequence used as a basis for sequence comparison. A reference sequence may be a subset or the entirety of a specified sequence; for example, as a segment of a full-length promoter sequence, or the complete promoter sequence.

(b) As used herein, "comparison window" makes reference to a contiguous and specified segment of a polynucleotide sequence, wherein the

polynucleotide sequence may be compared to a reference sequence and wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. Generally, the comparison window is at least 20 contiguous nucleotides in length and optionally can be 30, 40, 50, 100, or more contiguous nucleotides in length. Those of skill in the art understand that to avoid a high similarity to a reference sequence due to inclusion of gaps in the polynucleotide sequence a gap penalty is typically introduced and is subtracted from the number of matches.

Methods of alignment of sequences for comparison are well-known in the art. Optimal alignment of sequences for comparison may be conducted by the homology alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443; by computerized implementations of these algorithms, including, but not limited to: GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG) (575 Science Drive, Madison, Wisconsin). An example of the BLAST family of programs, which can be used to search database sequence similarity for the purposes of this invention, includes BLASTN program for nucleotide query sequences against nucleotide sequence dataset. See, Ausubel *et al.*, eds. (1995) *Current Protocols in Molecular Biology*, Chapter 19 (Greene Publishing and Wiley-Interscience, New York).

The BLAST homology alignment algorithm is useful for comparing fragments of the reference nucleotide or amino acid sequence to sequences from public databases. It is then necessary to apply a method of aligning the complete reference sequence against the complete public sequence to establish a % identity (in the case of polynucleotides) or % similarity (in the case of polypeptides). The GAP algorithm is such a method.

GAP uses the algorithm of Needleman and Wunsch (*J. Mol. Biol.* 48: 443-453, 1970) to find the alignment of two complete sequences that maximizes the number of matches and minimizes the number of gaps. GAP considers all possible alignments and gap positions and creates the alignment with the largest number of matched bases and the fewest gaps. It allows for the provision of a gap creation penalty and a gap extension penalty in units of matched bases. GAP

must make a profit of gap creation penalty number of matches for each gap it inserts. If a gap extension penalty greater than zero is chosen, GAP must, in addition, make a profit for each gap inserted of the length of the gap times the gap extension penalty. Default gap creation penalty values and gap extension penalty values in Version 10 of the Wisconsin Genetics Software Package for protein sequences are 8 and 2, respectively. For nucleotide sequences the default gap creation penalty is 50 while the default gap extension penalty is 3. The gap creation and gap extension penalties can be expressed as an integer selected from the group of integers consisting of from 0 to 200. Thus, for example, the gap creation and gap extension penalties can be 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65 or greater. Unless otherwise stated, for purposes of the invention, the preferred method of determining percent sequence identity is by the GAP version 10 algorithm using default parameters.

GAP presents one member of the family of best alignments. There may be many members of this family, but no other member has a better quality. GAP displays four figures of merit for alignments: Quality, Ratio, Identity, and Similarity. The Quality is the metric maximized in order to align the sequences. Ratio is the quality divided by the number of bases in the shorter segment. Percent Identity is the percent of the symbols that actually match. Percent Similarity is the percent of the symbols that are similar. Symbols that are across from gaps are ignored. A similarity is scored when the scoring matrix value for a pair of symbols is greater than or equal to 0.50, the similarity threshold. The scoring matrix used in Version 10 of the Wisconsin Genetics Software Package is BLOSUM62 (see Henikoff & Henikoff (1989) Proc. Natl. Acad. Sci. USA 89:10915).

(c) As used herein, "percentage of sequence identity" means the value determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of

comparison and multiplying the result by 100 to yield the percentage of sequence identity.

(d) The term "substantial identity" of polynucleotide sequences means that a polynucleotide comprises a sequence that has at least 70% sequence identity, preferably at least 80%, more preferably at least 90% and most preferably at least 95%, compared to a reference sequence using one of the alignment programs described using standard parameters.

Another indication that nucleotide sequences are substantially identical is if two nucleic acid molecules hybridize to each other under stringent conditions. Generally, stringent temperature conditions are selected to be about 5°C to about 2°C lower than the melting point (T_m) for the specific sequence at a defined ionic strength and pH. The denaturation or melting of DNA occurs over a narrow temperature range and represents the disruption of the double helix into its complementary single strands. The process usually is characterized by the temperature of the midpoint of transition, T_m , which is sometimes described as the melting temperature. Formulas are available in the art for the determination of melting temperatures. Typically, stringent wash conditions are those in which the salt concentration is about 0.02 molar at pH 7 and the temperature is at 50, 55, or 60°C.

The nucleotide sequences for the seed-preferred promoters disclosed in the present invention, as well as variants and fragments thereof, are useful in the genetic manipulation of any plant when operably linked with a heterologous nucleotide sequence whose expression is to be controlled to achieve a desired phenotypic response. By "operably linked" is intended the transcription or translation of the heterologous nucleotide sequence is under the influence of the promoter sequence. In this manner, the nucleotide sequences for the promoters of the invention may be provided in expression cassettes along with heterologous nucleotide sequences for expression in the plant of interest, more particularly in the seed of the plant.

Such expression cassettes will comprise a transcriptional initiation region comprising one of the promoter nucleotide sequences of the present invention, or variants or fragments thereof, operably linked to the heterologous nucleotide sequence whose expression is to be controlled by the seed-preferred promoters disclosed herein. Such an expression cassette is provided with a plurality of

restriction sites for insertion of the nucleotide sequence to be under the transcriptional regulation of the regulatory regions. The expression cassette may additionally contain selectable marker genes.

The expression cassette will include in the 5'-to-3' direction of transcription, a transcriptional and translational initiation region, a heterologous nucleotide sequence of interest, and a transcriptional and translational termination region functional in plants. The termination region may be native with the transcriptional initiation region comprising one of the promoter nucleotide sequences of the present invention, may be native with the DNA sequence of interest, or may be derived from another source. Convenient termination regions are available from the Ti-plasmid of *A. tumefaciens*, such as the octopine synthase and nopaline synthase termination regions. See also, Guerineau *et al.* (1991) *Mol. Gen. Genet.* 262:141-144; Proudfoot (1991) *Cell* 64:671-674; Sanfacon *et al.* (1991) *Genes Dev.* 5:141-149; Mogen *et al.* (1990) *Plant Cell* 2:1261-1272; Munroe *et al.* (1990) *Gene* 91:151-158; Ballas *et al.* (1989) *Nucleic Acids Res.* 17:7891-7903; Joshi *et al.* (1987) *Nucleic Acid Res.* 15:9627-9639.

The expression cassette comprising the promoter sequence of the present invention operably linked to a heterologous nucleotide sequence may also contain at least one additional nucleotide sequence for a gene to be cotransformed into the organism. Alternatively, the additional sequence(s) can be provided on another expression cassette.

Where appropriate, the heterologous nucleotide sequence whose expression is to be under the control of the promoter sequence of the present invention and any additional nucleotide sequence(s) may be optimized for increased expression in the transformed plant. That is, these nucleotide sequences can be synthesized using plant preferred codons for improved expression. Methods are available in the art for synthesizing plant-preferred nucleotide sequences. See, for example, U.S. Patent Nos. 5,380,831 and 5,436,391, and Murray *et al.* (1989) *Nucleic Acids Res.* 17:477-498, herein incorporated by reference.

Additional sequence modifications are known to enhance gene expression in a cellular host. These include elimination of sequences encoding spurious polyadenylation signals, exon-intron splice site signals, transposon-like repeats, and other such well-characterized sequences that may be deleterious to gene

expression. The G-C content of the heterologous nucleotide sequence may be adjusted to levels average for a given cellular host, as calculated by reference to known genes expressed in the host cell. When possible, the sequence is modified to avoid predicted hairpin secondary mRNA structures.

The expression cassettes may additionally contain 5' leader sequences in the expression cassette construct. Such leader sequences can act to enhance translation. Translation leaders are known in the art and include: picornavirus leaders, for example, EMCV leader (Encephalomyocarditis 5' noncoding region) (Elroy-Stein *et al.* (1989) *Proc. Nat. Acad. Sci. USA* 86:6126-6130); potyvirus leaders, for example, TEV leader (Tobacco Etch Virus) (Allison *et al.* (1986)); MDMV leader (Maize Dwarf Mosaic Virus) (*Virology* 154:9-20); human immunoglobulin heavy-chain binding protein (BiP) (Macejak *et al.* (1991) *Nature* 353:90-94); untranslated leader from the coat protein mRNA of alfalfa mosaic virus (AMV RNA 4) (Jobling *et al.* (1987) *Nature* 325:622-625); tobacco mosaic virus leader (TMV) (Gallie *et al.* (1989) *Molecular Biology of RNA*, pages 237-256); and maize chlorotic mottle virus leader (MCMV) (Lommel *et al.* (1991) *Virology* 81:382-385). See also Della-Cioppa *et al.* (1987) *Plant Physiology* 84:965-968. Other methods known to enhance translation and/or mRNA stability can also be utilized, for example, introns, and the like.

In those instances where it is desirable to have the expressed product of the heterologous nucleotide sequence directed to a particular organelle, particularly the plastid, amyloplast or vacuole, or to the endoplasmic reticulum, or secreted at the cell's surface or extracellularly, the expression cassette may further comprise a coding sequence for a transit peptide. Such transit peptides are well known in the art and include, but are not limited to, the transit peptide for the acyl carrier protein, the small subunit of RUBISCO, plant EPSP synthase, and the like.

In preparing the expression cassette, the various DNA fragments may be manipulated, so as to provide for the DNA sequences in the proper orientation and, as appropriate, in the proper reading frame. Toward this end, adapters or linkers may be employed to join the DNA fragments or other manipulations may be involved to provide for convenient restriction sites, removal of superfluous DNA, removal of restriction sites, or the like. For this purpose, *in vitro* mutagenesis,

primer repair, restriction, annealing, resubstitutions, for example, transitions and transversions, may be involved.

Reporter genes or selectable marker genes may be included in the expression cassette. Examples of suitable reporter genes known in the art can be found in, for example, Jefferson *et al.* (1991) in *Plant Molecular Biology Manual*, ed. Gelvin *et al.* (Kluwer Academic Publishers), pp. 1-33; DeWet *et al.* (1987) *Mol. Cell. Biol.* 7:725-737; Goff *et al.* (1990) *EMBO J.* 9:2517-2522; Kain *et al.* (1995) *BioTechniques* 19:650-655; and Chiu *et al.* (1996) *Current Biology* 6:325-330.

Selectable marker genes for selection of transformed cells or tissues can include genes that confer antibiotic resistance or resistance to herbicides. Examples of suitable selectable marker genes include, but are not limited to, genes encoding resistance to chloramphenicol (Herrera Estrella *et al.* (1983) *EMBO J.* 2:987-992); methotrexate (Herrera Estrella *et al.* (1983) *Nature* 303:209-213; Meijer *et al.* (1991) *Plant Mol. Biol.* 16:807-820); hygromycin (Waldron *et al.* (1985) *Plant Mol. Biol.* 5:103-108; Zhijian *et al.* (1995) *Plant Science* 108:219-227); streptomycin (Jones *et al.* (1987) *Mol. Gen. Genet.* 210:86-91); spectinomycin (Bretagne-Sagnard *et al.* (1996) *Transgenic Res.* 5:131-137); bleomycin (Hille *et al.* (1990) *Plant Mol. Biol.* 7:171-176); sulfonamide (Guerineau *et al.* 1990) *Plant Mol. Biol.* 15:127-136); bromoxynil (Stalker *et al.* (1988) *Science* 242:419-423); glyphosate (Shaw *et al.* (1986) *Science* 233:478-481); phosphinothricin (DeBlock *et al.* (1987) *EMBO J.* 6:2513-2518).

Other genes that could serve utility in the recovery of transgenic events but might not be required in the final product would include, but are not limited to, examples such as GUS (b-glucuronidase; Jefferson (1987) *Plant Mol. Biol. Rep.* 5:387), GFP (green fluorescence protein; Chalfie *et al.* (1994) *Science* 263:802), luciferase (Riggs *et al.* (1987) *Nucleic Acids Res.* 15(19):8115 and Luehrsen *et al.* (1992) *Methods Enzymol.* 216:397-414) and the maize genes encoding for anthocyanin production (Ludwig *et al.* (1990) *Science* 247:449).

The expression cassette comprising the particular promoter sequence of the present invention operably linked to a heterologous nucleotide sequence of interest can be used to transform any plant. In this manner, genetically modified plants, plant cells, plant tissue, seed, and the like can be obtained.

Transformation protocols as well as protocols for introducing nucleotide sequences into plants may vary depending on the type of plant or plant cell, i.e.,

monocot or dicot, targeted for transformation. Suitable methods of introducing nucleotide sequences into plant cells and subsequent insertion into the plant genome include microinjection (Crossway *et al.* (1986) *Biotechniques* 4:320-334), electroporation (Riggs *et al.* (1986) *Proc. Natl. Acad. Sci. USA* 83:5602-5606), Agrobacterium-mediated transformation (Townsend *et al.*, U.S. Pat No. 5,563,055; Zhao *et al.* WO US98/01268), direct gene transfer (Paszkowski *et al.* (1984) *EMBO J.* 3:2717-2722), and ballistic particle acceleration (see, for example, Sanford *et al.*, U.S. Patent No. 4,945,050; Tomes *et al.* (1995) "Direct DNA Transfer into Intact Plant Cells via Microprojectile Bombardment," in *Plant Cell, Tissue, and Organ Culture: Fundamental Methods*, ed. Gamborg and Phillips (Springer-Verlag, Berlin); and McCabe *et al.* (1988) *Biotechnology* 6:923-926). Also see Weissinger *et al.* (1988) *Ann. Rev. Genet.* 22:421-477; Sanford *et al.* (1987) *Particulate Science and Technology* 5:27-37 (onion); Christou *et al.* (1988) *Plant Physiol.* 87:671-674 (soybean); McCabe *et al.* (1988) *Bio/Technology* 6:923-926 (soybean); Finer and McMullen (1991) *In Vitro Cell Dev. Biol.* 27P:175-182 (soybean); Singh *et al.* (1998) *Theor. Appl. Genet.* 96:319-324 (soybean); Datta *et al.* (1990) *Biotechnology* 8:736-740 (rice); Klein *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:4305-4309 (maize); Klein *et al.* (1988) *Biotechnology* 6:559-563 (maize); Tomes, U.S. Patent No. 5,240,855; Busing *et al.*, U.S. Patent Nos. 5,322,783 and 5,324,646; Tomes *et al.* (1995) "Direct DNA Transfer into Intact Plant Cells via Microprojectile Bombardment," in *Plant Cell, Tissue, and Organ Culture: Fundamental Methods*, ed. Gamborg (Springer-Verlag, Berlin) (maize); Klein *et al.* (1988) *Plant Physiol.* 91:440-444 (maize); Fromm *et al.* (1990) *Biotechnology* 8:833-839 (maize); Hooykaas-Van Slogteren *et al.* (1984) *Nature* (London) 311:763-764; Bowen *et al.*, U.S. Patent No. 5,736,369 (cereals); Bytebier *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:5345-5349 (Liliaceae); De Wet *et al.* (1985) in *The Experimental Manipulation of Ovule Tissues*, ed. Chapman *et al.* (Longman, New York), pp. 197-209 (pollen); Kaeppler *et al.* (1990) *Plant Cell Reports* 9:415-418 and Kaeppler *et al.* (1992) *Theor. Appl. Genet.* 84:560-566 (whisker-mediated transformation); D'Halluin *et al.* (1992) *Plant Cell* 4:1495-1505 (electroporation); Li *et al.* (1993) *Plant Cell Reports* 12:250-255 and Christou and Ford (1995) *Annals of Botany* 75:407-413 (rice); Osjoda *et al.* (1996) *Nature Biotechnology* 14:745-750 (maize via *Agrobacterium tumefaciens*); all of which are herein incorporated by reference.

The cells that have been transformed may be grown into plants in accordance with conventional ways. See, for example, McCormick *et al.* (1986) *Plant Cell Reports* 5:81-84. These plants may then be grown, and either pollinated with the same transformed strain or different strains, and the resulting hybrid having constitutive expression of the desired phenotypic characteristic identified. Two or more generations may be grown to ensure that constitutive expression of the desired phenotypic characteristic is stably maintained and inherited and then seeds harvested to ensure constitutive expression of the desired phenotypic characteristic has been achieved.

The following examples are offered by way of illustration and not by way of limitation.

EXPERIMENTAL

Promoter regions for the maize genes *end1* and *end2* were isolated from maize plants and cloned. These genes were selected as sources of seed-preferred promoters based on the spatial expression of their gene products. The method for their isolation is described below.

Example 1: Isolation of Promoter Sequences

The procedure for promoter isolation is described in the User Manual for the Genome Walker kit sold by Clontech Laboratories, Inc., Palo Alto, California. Genomic DNA from maize line V3-4 A63 was prepared by grinding 10-day-old seedling leaves in liquid nitrogen, and the DNA prepared as described by Chen and Dellaporta (1994) in *The Maize Handbook*, ed. Freeling and Walbot (Springer-Verlag, Berlin) with a few minor modifications. Precipitated DNA was recovered using an inoculation loop and transferred to a 1.5 ml eppendorf tube containing 500 μ l of TE (10 mM Tris pH 8.0, 1 mM EDTA). The DNA was allowed to dissolve at room temperature for 15 minutes, phenol extracted and 2-propanol precipitated in 700 μ l. The precipitate was recovered and washed with 70% ethanol. The DNA was then placed in a clean 1.5 ml eppendorf tube to air dry and resuspended in 200 μ l of TE. RNase A was added to 10 μ g/ml and the mixture was incubated at 37°C for several hours. The DNA was then extracted once with phenol-chloroform, then chloroform, then ethanol precipitated and resuspended in TE. The DNA was then used exactly as described in the Genome Walker User Manual

(Clontech PT3042-1 version PR68687). Briefly, the DNA was digested separately with restriction enzymes *DraI*, *EcoRV*, *PvuII*, *Scal*, and *StuI*, all blunt-end cutters. The DNA was extracted with phenol, then chloroform, then ethanol precipitated. The Genome Walker adapters were ligated onto the ends of the restricted DNA. The resulting DNA is referred to as DL1-DL5, respectively.

For isolation of specific promoter regions, two nonoverlapping gene-specific primers (27-30 bp in length) were designed from the 5' end of the maize genes identified from sequence databases. The primers were designed to amplify the region upstream of the coding sequence, i.e. the 5' untranslated region and promoter of the chosen gene. The sequence of the primers are given below for each promoter described. The first round of PCR was performed on each DNA sample (DL1-5) with Clontech primer AP1 (sequence 5'-gtaatcgcactcactatagggc-3') and the gene-specific primer (gsp)1 with the sequences shown in SEQ ID NOS: 2 and 5.

PCR was performed in a model PTC-100 thermal cycler with HotBonnet from MJ Research (Watertown, MA) using reagents supplied with the Genome Walker kit. The following cycle parameters were used: 7 cycles of 94°C for 2 seconds, then 72°C for 3 minutes, followed by 32 cycles of 94°C for 2 seconds and 67°C for 3 minutes. Finally, the samples were held at 67°C for 4 minutes and then at 4°C until further analysis.

As described in the User Manual, the DNA from the first round of PCR was then diluted and used as a template in a second round of PCR using the Clontech AP2 primer (sequence 5'-actatagggcgcgcgtggt-3') and gene-specific primer (gsp)2 with the sequences shown in SEQ ID NOS: 3 and 6.

The cycle parameters for the second round were: 5 cycles of 94°C for 2 seconds, then 72°C for 3 minutes. Finally, the samples were held at 67°C for 4 minutes and then held at 4°C. Approximately 10¹ of each reaction were run on a 0.8% agarose gel, and bands (usually 500bp or larger) were excised, purified with the Sephaglas BandPrep kit (Pharmacia, Piscataway, NJ) and cloned into the TA vector pCR2.1 (Invitrogen, San Diego, California). Clones were sequenced for verification.

Example 2: Expression Data Using Promoter Sequences

Three promoter::GUS fusion constructs were prepared by the methods described below. All vectors were constructed using standard molecular biology techniques (Sambrook *et al.*, *Supra*). A reporter gene and a selectable marker gene for gene expression and selection was inserted between the multiple cloning sites of the pBluescript cloning vector (Stratagene Inc., 11011 N. Torrey Pines Rd., La Jolla, CA; www.stratagene.com). The reporter gene was the - glucuronidase (GUS) gene (Jefferson, R.A. *et al.*, 1986, *Proc. Natl. Acad. Sci. (USA)* 83:8447-8451) into whose coding region was inserted the second intron from the potato ST-LS1 gene (Vancanneyt *et al.*, *Mol. Gen. Genet.* 220:245-250, 1990), to produce intron-GUS, in order to prevent expression of the gene in *Agrobacterium* (see Ohta, S. *et al.*, 1990, *Plant Cell Physiol.* 31(6):805-813). The respective promoter regions were ligated in frame to sites 5' to the GUS gene. A fragment containing bases 2 to 310 from the terminator of the potato proteinase inhibitor (pinII) gene (An *et al.*, *Plant Cell* 1:115-122, 1989) was blunt-end ligated downstream of the GUS coding sequence, to create the GUS expression cassette. The 3' end of the terminator carried a NotI restriction site.

The promoter fusion end1::GUS::pinII was constructed using the GUS::pinII plasmid digested with BamHI, filled in with Klenow, and digested with NcoI. The promoter was isolated from the TOPOTA vector by digestion with Aval, filled in with Klenow, and then digestion with NcoI. The fragment was ligated into the digested expression cassette, and successful subcloning was confirmed by restriction digest and sequencing.

The *Agrobacterium* transformation plasmids were constructed by inserting the GUS expression cassette as a HindIII/NotI fragment and the BAR expression cassette as a NotI/SacI fragment between the right and left T-DNA borders in pSB11 at HindIII and SacI sites. The GUS cassette was inserted proximal to the right T-DNA border. The plasmid pSB11 was obtained from Japan Tobacco Inc. (Tokyo, Japan). The construction of pSB11 from pSB21 and the construction of pSB21 from starting vectors is described by Komari *et al.* (1996, *Plant J.* 10:165-174). The T-DNA of the plasmids were integrated into the superbinary plasmid pSB1 (Saito *et al.*, EP 672 752 A1) by homologous recombination between the two plasmids. The plasmid pSB1 was also obtained from Japan Tobacco Inc. *E. coli* strain HB101 containing the expression cassettes was mated with

Agrobacterium strain LBA4404 harboring pSB1 to create the cointegrate plasmid in Agrobacterium using the method of Ditta *et al.*, (*Proc. Natl. Acad. Sci. USA* 77:7347-7351, 1980). Successful recombination was verified by a Sall restriction digest of the plasmid.

Example 3: Transformation and Regeneration of Maize Callus via Agrobacterium

Preparation of Agrobacterium suspension:

Agrobacterium is streaked out from a -80° frozen aliquot onto a plate containing PHI-L medium and cultured at 28°C in the dark for 3 days. PHI-L media comprises 25 ml/l Stock Solution A, 25 ml/l Stock Solution B, 450.9 ml/l Stock Solution C and spectinomycin (Sigma Chemicals) added to a concentration of 50 mg/l in sterile ddH₂O (stock solution A: K₂HPO₄ 60.0 g/l, NaH₂PO₄ 20.0 g/l, adjust pH to 7.0 w/KOH and autoclave; stock solution B: NH₄Cl 20.0 g/l, MgSO₄·7H₂O 6.0 g/l, KCl 3.0 g/l, CaCl₂ 0.20 g/l, FeSO₄·7H₂O 50.0 mg/l, autoclave; stock solution C: glucose 5.56g/l, agar 16.67 g/l (#A-7049, Sigma Chemicals, St. Louis, MO) and autoclave).

The plate can be stored at 4°C and used usually for about 1 month. A single colony is picked from the master plate and streaked onto a plate containing PHI-M medium [yeast extract (Difco) 5.0 g/l; peptone (Difco) 10.0 g/l; NaCl 5.0 g/l; agar (Difco) 15.0 g/l; pH 6.8, containing 50 mg/L spectinomycin] and incubated at 28°C in the dark for 2 days.

Five ml of either PHI-A, [CHU(N6) basal salts (Sigma C-1416) 4.0 g/l, Eriksson's vitamin mix (1000X, Sigma-1511) 1.0 ml/l; thiamine.HCl 0.5 mg/l (Sigma); 2,4-dichlorophenoxyacetic acid (2,4-D, Sigma) 1.5 mg/l; L-proline (Sigma) 0.69 g/l; sucrose (Mallinckrodt) 68.5 g/l; glucose (Mallinckrodt) 36.0 g/l; pH 5.2] for the PHI basic medium system, or PHI-I [MS salts (GIBCO BRL) 4.3 g/l; nicotinic acid (Sigma) 0.5 mg/l; pyridoxine.HCl (Sigma) 0.5 mg/l; thiamine.HCl 1.0 mg/l; myo-inositol (Sigma) 0.10 g/l; vitamin assay casamino acids (Difco Lab) 1.0 g/l; 2, 4-D 1.5 mg/l; sucrose 68.50 g/l; glucose 36.0 g/l; adjust pH to 5.2 w/KOH and filter-sterilize] for the PHI combined medium system and 5 ml of 100 mM (3'-5'-Dimethoxy-4'-hydroxyacetophenone, Aldrich chemicals) are added to a 14 ml Falcon tube in a hood. About 3 full loops (5 mm loop size) Agrobacterium is collected from the plate and suspended in the tube, then the tube is vortexed to

make an even suspension. One ml of the suspension is transferred to a spectrophotometer tube and the OD of the suspension adjusted to 0.72 at 550 nm by adding either more Agrobacterium or more of the same suspension medium, for an Agrobacterium concentration of approximately 0.5×10^9 cfu/ml to 1×10^9 cfu/ml. The final Agrobacterium suspension is aliquoted into 2 ml microcentrifuge tubes, each containing 1 ml of the suspension. The suspensions are then used as soon as possible.

Embryo isolation, infection and co-cultivation:

About 2 ml of the same medium (here PHI-A or PHI-I) used for the Agrobacterium suspension are added into a 2 ml microcentrifuge tube. Immature embryos are isolated from a sterilized ear with a sterile spatula (Baxter Scientific Products S1565) and dropped directly into the medium in the tube. A total of about 100 embryos are placed in the tube. The optimal size of the embryos is about 1.0-1.2 mm. The cap is then closed on the tube and the tube vortexed with a Vortex Mixer (Baxter Scientific Products S8223-1) for 5 sec. at maximum speed. The medium is removed and 2 ml of fresh medium are added and the vortexing repeated. All of the medium is drawn off and 1 ml of Agrobacterium suspension is added to the embryos and the tube vortexed for 30 sec. The tube is allowed to stand for 5 min. in the hood. The suspension of Agrobacterium and embryos was poured into a Petri plate containing either PHI-B medium [CHU(N6) basal salts (Sigma C-1416) 4.0 g/l; Eriksson's vitamin mix (1000X, Sigma-1511) 1.0 ml/l; thiamine.HCl 0.5 mg/l; 2,4-D 1.5 mg/l; L-proline 0.69 g/l; silver nitrate 0.85 mg/l; gelrite (Sigma) 3.0 g/l; sucrose 30.0 g/l; acetosyringone 100 mM; pH 5.8], for the PHI basic medium system, or PHI-J medium [MS Salts 4.3 g/l; nicotinic acid 0.50 mg/l; pyridoxine HCl 0.50 mg/l; thiamine.HCl 1.0 mg/l; myo-inositol 100.0 mg/l; 2,4-D 1.5 mg/l; sucrose 20.0 g/l; glucose 10.0 g/l; L-proline 0.70 g/l; MES (Sigma) 0.50 g/l; 8.0 g/l agar (Sigma A-7049, purified) and 100 mM acetosyringone with a final pH of 5.8 for the PHI combined medium system. Any embryos left in the tube are transferred to the plate using a sterile spatula. The Agrobacterium suspension is drawn off and the embryos placed axis side down on the media. The plate is sealed with Parafilm tape or Pylon Vegetative Combine Tape (product named "E.G.CUT" and is available in 18 mm x 50 m sections; Kyowa Ltd., Japan) and incubated in the dark at 23-25°C for about 3 days of co-cultivation.

Resting, selection and regeneration steps:

For the resting step, all of the embryos are transferred to a new plate containing PHI-C medium [CHU(N6) basal salts (Sigma C-1416) 4.0 g/l; Eriksson's vitamin mix (1000X Sigma-1511) 1.0 ml/l; thiamine.HCl 0.5 mg/l; 2,4-D 1.5 mg/l; L-proline 0.69 g/l; sucrose 30.0 g/l; MES buffer (Sigma) 0.5 g/l; agar (Sigma A-7049, purified) 8.0 g/l; silver nitrate 0.85 mg/l; carbenicillin 100 mg/l; pH 5.8]. The plate is sealed with Parafilm or Pylon tape and incubated in the dark at 28°C for 3-5 days.

Longer co-cultivation periods may compensate for the absence of a resting step since the resting step, like the co-cultivation step, provides a period of time for the embryo to be cultured in the absence of a selective agent. Those of ordinary skill in the art can readily test combinations of co-cultivation and resting times to optimize or improve the transformation frequency without undue experimentation.

For selection, all of the embryos are then transferred from the PHI-C medium to new plates containing PHI-D medium, as a selection medium, [CHU(N6) basal salts (SIGMA C-1416) 4.0 g/l; Eriksson's vitamin mix (1000X, Sigma-1511) 1.0 ml/l; thiamine.HCl 0.5 mg/l; 2,4-D 1.5 mg/l; L-proline 0.69 g/l; sucrose 30.0 g/l; MES buffer 0.5 g/l; agar (Sigma A-7049, purified) 8.0 g/l; silver nitrate 0.85 mg/l; carbenicillin (ICN, Costa Mesa, CA) 100 mg/l; bialaphos (Meiji Seika K.K., Tokyo, Japan) 1.5 mg/l for the first two weeks followed by 3 mg/l for the remainder of the time.; pH 5.8] putting about 20 embryos onto each plate. The plates are sealed as described above and incubated in the dark at 28°C for the first two weeks of selection. The embryos are transferred to fresh selection medium at two-week intervals. The tissue is subcultured by transferring to fresh selection medium for a total of about 2 months. The herbicide-resistant calli are then "bulked up" by growing on the same medium for another two weeks until the diameter of the calli is about 1.5-2 cm.

For regeneration, the calli are then cultured on PHI-E medium [MS salts 4.3 g/l; myo-inositol 0.1 g/l; nicotinic acid 0.5 mg/l, thiamine.HCl 0.1 mg/l, Pyridoxine.HCl 0.5 mg/l, Glycine 2.0 mg/l, Zeatin 0.5 mg/l, sucrose 60.0 g/l, Agar (Sigma, A-7049) 8.0 g/l, Indoleacetic acid (IAA, Sigma) 1.0 mg/l, Abscisic acid (ABA, Sigma) 0.1 mM, Bialaphos 3 mg/l, carbenicillin 100 mg/l adjusted to pH 5.6] in the dark at 28°C for 1-3 weeks to allow somatic embryos to mature. The calli

are then cultured on PHI-F medium (MS salts 4.3 g/l; myo-inositol 0.1 g/l; Thiamine.HCl 0.1 mg/l, Pyridoxine.HCl 0.5 mg/l, Glycine 2.0 mg/l, nicotinic acid 0.5 mg/l; sucrose 40.0 g/l; gelrite 1.5 g/l; pH 5.6] at 25°C under a daylight schedule of 16 hrs. light (270 $\mu\text{E m}^{-2}\text{sec}^{-1}$) and 8 hrs. dark until shoots and roots develop. Each small plantlet is then transferred to a 25x150 mm tube containing PHI-F medium and grown under the same conditions for approximately another week. The plants are transplanted to pots with soil mixture in a greenhouse. GUS+ events are determined at the callus stage or regenerated plant stage.

For Hi-II a preferred optimized protocol was 0.5×10^9 cfu/ml *Agrobacterium* a 3-5 day resting step and no AgNO_3 in the infection medium (PHI-A medium).

Example 4: : *In Situ* Localization of *End2* mRNA in Developing Maize Kernel

In situ hybridization was performed using the protocol of Jackson, D.P. (1991) *In situ* Hybridization in Plants, *Molecular Plant Pathology: A Practical Approach*, D.J. Bowles, S. J. Gurr, and M. McPherson, eds. Oxford University Press, England, pp.63-74. Both a sense and antisense probe corresponding to a protein of the *end2* cDNA were used. Probes were labelled non-isotopically with digoxigenin and incubated with various sections of 5 DAP and 9 DAP (days after pollination) kernels of maize line CJ27 which had been fixed and embedded. Following extensive washing to remove unbound probe, sections were incubated with anti-digoxigenin alkaline phosphatase to detect areas of probe hybridization. For *end2*, mRNA was detected specifically with the antisense probe and restricted to the aleurone layer of endosperm tissue. The sense control probe did not hybridize.

Example 5: Northern Analysis of Gene Expression in Vegetative Tissue and Developing Kernels

Total RNA (10 μg) was size fractionated on a 1% formaldehyde agarose gel and transferred to a nitrocellular membrane. Membranes were hybridized under stringent conditions with ^{32}P -labelled probes representing cDNA fragments of the various genes. After extensive washing to remove unbound probe, membranes were exposed on X-ray film. RNA samples were obtained from vegetative tissues as well as developing maize kernels.

The *end1* expression pattern showed no expression in vegetative tissues. *End1* was predominantly expressed in early to mid-development whole kernel. Expression was also seen in isolated 7 DAP (days after pollination) embryo, 10 DAP endosperm and 10 DAP pericarp.

The expression pattern of *end2* likewise showed no expression in vegetative tissue. *End2* showed expression in 9 – 14 DAP whole kernel. *End2* also showed expression in isolated 7 DAP embryo, 10 DAP endosperm, and 10 DAP pericarp.

All publications and patent applications mentioned in the specification are indicative of the level of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.

WHAT IS CLAIMED IS:

1. An isolated promoter that is capable of driving transcription in a seed-preferred manner, wherein said promoter comprises a nucleotide sequence selected from the group consisting of:
 - a) sequences capable of driving expression of coding regions selected from the group consisting of coding regions for *end1* or *end2*;
 - b) a sequence comprising at least 40 contiguous nucleotides of the sequence set forth in either of SEQ ID NOS:1 or 4;
 - c) a sequence comprising a variant or fragment of the nucleotide sequence set forth in either of SEQ ID NOS: 1 or 4;
 - d) the nucleotide sequences set forth in SEQ ID NOS: 1 or 4;
 - e) a sequence that hybridizes to any one of SEQ ID NOS: 1 or 4 under low stringency conditions; and
 - f) a nucleotide sequence complementary to a nucleotide sequence of (a) through (e).

2. An expression cassette comprising a promoter and a nucleotide sequence operably linked to said promoter, wherein said promoter is capable of initiating seed-preferred transcription of said nucleotide sequence in a plant cell, wherein said promoter comprises a nucleotide sequence selected from the group consisting of:
 - a) sequences capable of driving expression of coding regions selected from the group consisting of coding regions for *end1* or *end2*;
 - b) a sequence comprising at least 40 contiguous nucleotides of the sequence set forth in either of SEQ ID NOS:1 or 4;
 - c) a sequence comprising a variant or fragment of the nucleotide sequence set forth in either of SEQ ID NOS: 1 or 4;
 - d) the nucleotide sequences set forth in SEQ ID NOS: 1 or 4;
 - e) a sequence that hybridizes to any one of SEQ ID NOS:

- 1 or 4 under low stringency conditions; wherein expression of said sequence changes the phenotype of seed of a plant transformed with said expression cassette; and
- f) a nucleotide sequence complementary to a nucleotide sequence of (a) through (e).
3. A transformation vector comprising an expression cassette, said expression cassette comprising a maize promoter and a nucleotide sequence operably linked to said promoter, wherein said promoter is capable of initiating seed-preferred transcription of said nucleotide sequence in a plant cell, wherein said promoter comprises a nucleotide sequence selected from the group consisting of:
- a) sequences capable of driving expression of coding regions selected from the group consisting of coding regions for *end1* or *end2*;
 - b) sequence comprising at least 40 contiguous nucleotides of the sequence set forth in either of SEQ ID NOS:1 or 4;
 - c) a sequence comprising a variant or fragment of the nucleotide sequence set forth in either of SEQ ID NOS: 1 or 4;
 - d) the nucleotide sequences set forth in SEQ ID NOS: 1 or 4;
 - e) a sequence that hybridizes to any one of SEQ ID NOS: 1 or 4 under low stringency conditions; and
 - f) a nucleotide sequence complementary to a nucleotide sequence of (a) through (e).
4. A plant stably transformed with an expression cassette comprising a maize promoter and a nucleotide sequence operably linked to said promoter, wherein said promoter is capable of initiating constitutive transcription of said nucleotide sequence in a plant cell, wherein said promoter comprises a nucleotide sequence selected from the group consisting of:
- a) sequences capable of driving expression of coding regions selected from the group consisting of coding regions for *end1* or *end2*;

- b) a sequence comprising at least 40 contiguous nucleotides of the sequence set forth in either of SEQ ID NOS:1 or 4;
 - c) a sequence comprising a variant or fragment of the nucleotide sequence set forth in either of SEQ ID NOS: 1 or 4;
 - d) the nucleotide sequences set forth in SEQ ID NOS: 1 or 4;
 - e) a sequence that hybridizes to any one of SEQ ID NOS: 1 or 4 under low stringency conditions; wherein expression of the nucleotide sequences alters the phenotype of seed of said transformed plant; and
 - f) a nucleotide sequence complementary to a nucleotide sequence of (a) through (e).
5. The plant of claim 4, wherein said plant is a monocot.
6. The plant of claim 5, wherein said monocot is maize, wheat, rice, barley, sorghum, or rye.
7. The plant of claim 4, wherein said plant is a dicot.
8. Seed of the plant of claims 4.
9. A method for selectively expressing a nucleotide sequence in a plant seed, said method comprising transforming a plant cell with a transformation vector comprising an expression cassette, said expression cassette comprising a maize promoter and a nucleotide sequence operably linked to said promoter, wherein said promoter is capable of initiating seed-preferred transcription of said nucleotide sequence in a plant cell, wherein said promoter comprises a nucleotide sequence selected from the group consisting of:
- a) sequences capable of driving expression of coding regions selected from the group consisting of coding regions for *end1* or *end2*;
 - b) a sequence comprising at least 40 contiguous nucleotides of the sequence set forth in either of SEQ ID NOS:1 or 4;

- c) a sequence comprising a variant or fragment of the nucleotide sequence set forth in either of SEQ ID NOS: 1 or 4;
 - d) the nucleotide sequences set forth in SEQ ID NOS: 1 or 4;
 - e) a sequence that hybridizes to any one of SEQ ID NOS: 1 or 4 under low stringency conditions; and
 - f) a nucleotide sequence complementary to a nucleotide sequence of (a) through (e).
10. The method of claim 9 further comprising regenerating a stably transformed plant from said transformed plant cell; wherein expression of said nucleotide sequences alters the phenotype of said plant seed.
11. The method of claim 9, wherein said heterologous nucleotide sequence encodes a gene involved in fatty acid synthesis.
12. The method of claim 9, wherein said heterologous nucleotide sequence encodes a gene having enhanced amino acid content.
13. A plant cell stably transformed with an expression cassette comprising a maize promoter and a nucleotide sequence operably linked to said promoter, wherein said promoter is capable of initiating constitutive transcription of said nucleotide sequence in a plant cell, wherein said promoter comprises a nucleotide sequence selected from the group consisting of:
- a) sequences capable of driving expression of coding regions selected from the group consisting of coding regions for *end1* or *end2*;
 - b) a sequence comprising at least 40 contiguous nucleotides of the sequence set forth in either of SEQ ID NOS:1 or 4;
 - c) a sequence comprising a variant or fragment of the nucleotide sequence set forth in either of SEQ ID NOS: 1 or 4;
 - d) the nucleotide sequences set forth in SEQ ID NOS: 1 or 4;
 - e) a sequence that hybridizes to any one of SEQ ID NOS:

1 or 4 under low stringency conditions; wherein expression of the nucleotide sequences alters the phenotype of seed of said transformed plant cell; and

- f) a nucleotide sequence complementary to a nucleotide sequence of (a) through (e).

14. The plant cell of claim 13, wherein said plant cell is from a monocotyledonous plant.
15. The plant cell of claim 14, wherein said plant cell is from maize, wheat, rice, barley, sorghum, or rye.
16. The plant cell of claim 13, wherein said plant cell is from a dicotyledonous plant.

SEQUENCE LISTING

<110> Pioneer Hi-Bred International, Inc.

<120> Seed-Preferred Promoters from End Genes

<130> 0933-PCT

<150> US 60/098,230

<151> 1998-08-28

<160> 6

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 713

<212> DNA

<213> Zea mays

<400> 1

ggctggtaaa	aaccattatt	aactttaaca	tcgaatcaaa	actgacaaat	tttatacttt	60
cacagagcag	cagaaattta	tacaatatga	ttgaatacaa	gatgtaggac	ccgatggaga	120
gaatTTTTTT	gtctcctata	tgcttgaata	cccaacataa	tatcttcgca	gcatactatc	180
tatctaatag	aaaaattata	atatagttaa	atacttaagt	agtatctagt	ggatagaatt	240
caatatctca	tacatgcatg	aggagtaata	tctactagac	atgcaacata	tttttatcta	300
tctaatagaa	tatatataat	aaagttaa	attatatgca	tcacctacta	tatataattt	360
gatatctttt	agatgtataa	gggactaaga	ataatatctc	tagcacacac	gcaatgcatt	420
atctatctaa	atatattata	taatagttaa	atattaatta	tacgtagtct	aaacctacat	480
ataagcctac	ccatccccac	ttagagatct	cagtgtcaca	catagaccat	acatctcact	540
tcgccaagaa	aatttcgtca	acagttgaag	ttatacccat	ggcaaaaacta	ctcttgggtt	600
tgctccttgc	ccttgctatt	ctagggacaa	catcggtg	tggttggtgta	caagaagggc	660
gaattctgca	gatatccatc	acactggcgg	ccgctcgagc	atgcatctag	agg	713

<210> 2

<211> 27

<212> DNA

<213> Zea mays

<400> 2

tggtgggtat	tcaagcatat	aggagac				27
------------	------------	---------	--	--	--	----

<210> 3

<211> 29

<212> DNA

<213> Zea mays

<400> 3

ccatcgggtc	ctacatcttg	tattcaatc				29
------------	------------	-----------	--	--	--	----

<210> 4

<211> 1224

<212> DNA

<213> Zea mays

<400> 4

tactataggg	cagcgtggt	cgacggccc	ggctggtaaa	aagtaattga	acccaaaata	60
tcatgggatg	tttgggtgaag	acagtgatca	gtgatttttt	tatatctata	tatatatcaa	120
agatacttga	ttttctagaa	ggttcttttt	gttgttttcc	cttatgtttt	tacgcatgat	180

gcaattcttt	ttgagagggt	tccgatgcat	tgatgttatt	gtattatctc	ctatatatag	240
gtcgacgtac	attatgtatt	gcaataacca	gttaactgga	tccagcttcg	cttagttttt	300
agtttttggc	agaaaaaatg	atcaatgttt	cacaaaccaa	atatttttat	aacttttgat	360
gaaagaagat	caccacggtc	atatctaggg	gtggtaacaa	attgcatct	aaatgtttct	420
tcataaaaaa	taaggcttct	taataaattt	tagttcaaaa	taaatacgaa	taaagtctga	480
ttctaactctg	attcgatcct	taaattttat	aatgcaaaat	ttagagctca	ttaccacctc	540
tagtcatatg	tctagtctga	ggtatatcca	aaaagccctt	tctctaaatt	ccacacccaa	600
ctcagatggt	tgcaaataaa	tactcogact	ccaaaatgta	ggtgaagtgc	aactttctcc	660
atthtatatc	aacatttggt	atthtttggt	taacatttca	cactcaaac	taattaataa	720
aatacgtggt	tgttgaacgt	gcgcacatgt	ctcccttaca	ttatgttttt	ttatthtatgt	780
attattgttg	ttttcctccg	aacaacttgt	caacatatca	tcattggtct	ttaatattta	840
tgaatatgga	agcctagtta	tttacacttg	gctacacact	agttgtagtt	ttgccacttg	900
tctaacatgc	aactctagta	gttttgccac	ttgcctggca	cgcgactcta	gtattgacac	960
ttgtatagca	aataatgcca	atacgacacc	tggccttaca	tgaaacatta	tttttgacac	1020
ttgtatacca	tgcaacatta	ccattgacat	ttgtccatac	acattatatac	aaatatattg	1080
agcgcacgtc	acaaaactcga	tacaaagctg	gatgaccctc	cctcaccaca	tctataaaaa	1140
cccagcgcct	actgtaaatac	actcacaaca	caacacatat	cttttagtaa	cttttcaata	1200
ggcgtcccc	aagaactagt	aaac				1224

<210> 5
 <211> 30
 <212> DNA
 <213> Zea mays

<400> 5
 acttatcagg ctttgagggt cattctcaca 30

<210> 6
 <211> 32
 <212> DNA
 <213> Zea mays

<400> 6
 tccatcagca tgagagagcc tatggcaaac at 32

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/19604

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C12N15/82 C12N15/29 A01H5/00		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) IPC 7 C12N A01H		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 93 09237 A (SANDOZ AG ;SANDOZ AG (DE); SANDOZ LTD (CH)) 13 May 1993 (1993-05-13) page 30 -page 36	1
P, X	WO 98 53086 A (CHAO WUN S ;GRIMES HOWARD D (US); UNIV WASHINGTON (US)) 26 November 1998 (1998-11-26) see SEQ ID NO:6 and 7	1
A	US 5 633 363 A (COLBERT JAMES T ET AL) 27 May 1997 (1997-05-27) see sequence ID NO:1	1
	-/-	
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
° Special categories of cited documents :		
"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "G" document member of the same patent family		
Date of the actual completion of the international search 21 December 1999		Date of mailing of the international search report 11/01/2000
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer Maddox, A

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/19604

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>PUIGDOMENECH, P., ET AL.: "Zea mays tandem genes for alpha-tubulin and alpha2-tubulin" EMBL ACCESSION NO:X15704, 28 December 1989 (1989-12-28), XP002126401 the whole document</p> <p>& MONTOLIU, L., ET AL.: "A tandem of alpha-tubulin genes preferentially expressed in radicular tissues of Zea mays" PLANT MOLECULAR BIOLOGY, vol. 14, 1989, pages 1-15,</p>	1
A	<p>WO 97 30162 A (SOUTHERTON SIMON GEORGE ; TEASDALE ROBERT DIXON (AU); MOURADOV AIDY) 21 August 1997 (1997-08-21) figure 31; example 7</p>	1
A	<p>WO 98 30698 A (CALGENE INC ; PEAR JULIE R (US); STALKER DAVID M (US)) 16 July 1998 (1998-07-16) figures 1A-D</p>	1
P,A	<p>WALBOT, V.: "605094A04.x2 605 - Endosperm cDNA library from Schmidt lab Zea mays cDNA, mRNA sequence" EMBL ACCESSION NO:AI833693, 14 July 1999 (1999-07-14), XP002126402 the whole document</p>	1
P,A	<p>WALBOT, V., ET AL.: "605086G12.y1 605 - Endosperm cDNA library from Schmidt lab Zea mays cDNA, mRNA sequence" EMBL ACCESSION NO:AI812206, 8 July 1999 (1999-07-08), XP002126403 the whole document</p>	1
P,A	<p>WO 98 56921 A (DOW AGROSCIENCES LLC) 17 December 1998 (1998-12-17) see SEQ ID NOS:15 and 16</p>	1
A	<p>WO 97 28247 A (BIOCEM ; LUDEVID DOLORES (ES); TORRENT MARGARITA (ES); ALVAREZ INAK) 7 August 1997 (1997-08-07) the whole document</p>	1-16
A	<p>WO 98 26064 A (DEKALB GENETICS CORP) 18 June 1998 (1998-06-18)</p>	1-16
A	<p>WO 98 08961 A (DOAN DANNY N P ; OLSEN ODD ARNE (NO); LINNESTAD CASPER (NO)) 5 March 1998 (1998-03-05) the whole document</p>	1-16
	-/-	

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/19604

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 677 474 A (ROGERS JOHN C) 14 October 1997 (1997-10-14) the whole document	1-16
A	US 5 420 034 A (KNAUF VIC C ET AL) 30 May 1995 (1995-05-30) the whole document	11

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/19604

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

see additional sheet

1. As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- The additional search fees were accompanied by the applicant's protest.
- No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-16 all partially

Seed-preferred promoter as represented by SEQ ID NO:1 and its uses

2. Claims: 1-16 all partially

Seed-preferred promoter as represented by SEQ ID NO:4 and its uses

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/US 99/19604

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9309237 A	13-05-1993	AU 670417 B	18-07-1996
		AU 2920392 A	07-06-1993
		EP 0611395 A	24-08-1994
		HU 66207 A	28-10-1994
		JP 7500965 T	02-02-1995
		ZA 9208538 A	05-05-1994
WO 9853086 A	26-11-1998	AU 7500998 A	11-12-1998
US 5633363 A	27-05-1997	NONE	
WO 9730162 A	21-08-1997	AU 1713297 A	02-09-1997
		CA 2259456 A	21-08-1997
		CN 1216066 A	05-05-1999
		EP 0882133 A	09-12-1998
		NO 983775 A	15-10-1998
WO 9830698 A	16-07-1998	AU 5732298 A	03-08-1998
WO 9856921 A	17-12-1998	AU 7830198 A	30-12-1998
WO 9728247 A	07-08-1997	FR 2744134 A	01-08-1997
		AU 1550297 A	22-08-1997
		CA 2242903 A	07-08-1997
		EP 0877754 A	18-11-1998
WO 9826064 A	18-06-1998	AU 5795898 A	03-07-1998
		EP 0942971 A	22-09-1999
WO 9808961 A	05-03-1998	AU 3781897 A	19-03-1998
US 5677474 A	14-10-1997	AU 638409 B	01-07-1993
		AU 4037289 A	05-03-1990
		EP 0428572 A	29-05-1991
		JP 4500153 T	16-01-1992
		WO 9001551 A	22-02-1990
US 5420034 A	30-05-1995	US 5753475 A	19-05-1998
		US 5608152 A	04-03-1997
		US 5750385 A	12-05-1998
		US 5981839 A	09-11-1999
		AT 104352 T	15-04-1994
		AU 612326 B	11-07-1991
		AU 7630287 A	04-02-1988
		DE 3789582 D	19-05-1994
		EP 0255378 A	03-02-1988
		ES 2054676 T	16-08-1994
		IN 170125 A	15-02-1992
		NZ 221259 A	28-05-1990
		AU 609724 B	09-05-1991
		AU 7630387 A	04-02-1988
		EP 0255377 A	03-02-1988
		FI 873252 A	01-02-1988
		FI 873253 A	01-02-1988
		JP 63119680 A	24-05-1988
		JP 63112987 A	18-05-1988
		NO 873190 A	01-02-1988
		NO 873191 A	01-02-1988

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/19604

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5420034 A		NZ 221258 A	28-05-1990
		US 5110728 A	05-05-1992
		US 5315001 A	24-05-1994