

**(12) STANDARD PATENT**  
**(19) AUSTRALIAN PATENT OFFICE**

(11) Application No. **AU 2009328318 B2**

(54) Title  
**Anti-cMet antibody**

(51) International Patent Classification(s)  
**C07K 16/28** (2006.01)                      **C07K 16/00** (2006.01)  
**A61K 39/395** (2006.01)                      **C07K 16/46** (2006.01)  
**A61P 35/00** (2006.01)

(21) Application No: **2009328318**                      (22) Date of Filing: **2009.12.02**

(87) WIPO No: **WO10/069765**

(30) Priority Data

(31) Number	(32) Date	(33) Country
<b>PCT/IB2008/055663</b>	<b>2008.12.02</b>	<b>IB</b>
<b>61/184,502</b>	<b>2009.06.05</b>	<b>US</b>

(43) Publication Date: **2010.06.24**

(44) Accepted Journal Date: **2015.11.05**

(71) Applicant(s)  
**Pierre Fabre Medicament**

(72) Inventor(s)  
**Goetsch, Liliane;Wurch, Thierry;Bes, Cedric**

(74) Agent / Attorney  
**Phillips Ormonde Fitzpatrick, L 16 333 Collins St, Melbourne, VIC, 3000**

(56) Related Art  
**WO 2009/007427 A2 (PIERRE FABRE MEDICAMENT) 15 January 2009**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
24 June 2010 (24.06.2010)

PCT

(10) International Publication Number  
**WO 2010/069765 A1**

(51) International Patent Classification:

C07K 16/28 (2006.01) A61K 39/395 (2006.01)  
C07K 16/00 (2006.01) A61P 35/00 (2006.01)  
C07K 16/46 (2006.01)

(21) International Application Number:

PCT/EP2009/066201

(22) International Filing Date:

2 December 2009 (02.12.2009)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

PCT/IB2008/055663  
2 December 2008 (02.12.2008) IB  
61/184,502 5 June 2009 (05.06.2009) US

(71) Applicant (for all designated States except US):  
**PIERRE FABRE MEDICAMENT** [FR/FR]; 45, place  
Abel Gance, F-92100 Boulogne-Billancourt (FR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GOETSCH, Liliane**  
[FR/FR]; 15 route de Cluses, F-74130 Ayze (FR).  
**WURCH, Thierry** [FR/FR]; 7 route des Etoles, Le Clos  
Bellevue, F-74140 Machilly (FR). **BES, Cédric** [FR/FR];  
12 boulevard des Acanthes, F-30250 Villevicille (FR).

(74) Agent: **WARCOIN, Jacques**; Cabinet Regimbeau, 20,  
rue de Chazelles, F-75847 Paris Cedex 17 (FR).

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ,  
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO,

DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,  
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,  
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,  
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,  
NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD,  
SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT,  
TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,  
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ,  
TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,  
MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM,  
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,  
ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted  
a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of  
the earlier application (Rule 4.17(iii))
- of inventorship (Rule 4.17(iv))

Published:

- with international search report (Art. 21(3))
- with (an) indication(s) in relation to deposited biological  
material furnished under Rule 13bis separately from the  
description (Rules 13bis.4(d)(i) and 48.2(a)(viii))
- with sequence listing part of description (Rule 5.2(a))



WO 2010/069765 A1

(54) Title: ANTI-cMET ANTIBODY

(57) Abstract: Antibody capable of binding specifically to the human c-Met receptor and/or capable of specifically inhibiting the tyrosine kinase activity of said receptor, with an improved antagonistic activity, said antibody comprising a modified hinge region. A composition comprising such an antibody antagonist to c-Met and its use as a medicament for treating cancer.

*ANTI-cMET ANTIBODY*

The present invention relates to a novel divalent antibody capable of binding specifically to the human c-Met receptor and/or capable of specifically inhibiting the tyrosine kinase activity of said receptor, as well as the amino acid and nucleic acid sequences coding for said antibody. More particularly, the antibody according to the invention is capable of inhibiting the c-Met dimerization. The invention likewise comprises the use of said antibody as a medicament for the prophylactic and/or therapeutic treatment of cancers or any pathology connected with the overexpression of said receptor as well as in processes or kits for diagnosis of illnesses connected with the over-expression of c-Met. The invention finally comprises products and/or compositions comprising such an antibody in combination with other antibodies and/or chemical compounds directed against other growth factors involved in tumor progression or metastasis and/or compounds and/or anti-cancer agents or agents conjugated with toxins and their use for the prevention and/or the treatment of certain cancers.

Receptor tyrosine kinase (RTK) targeted agents such as trastuzumab, cetuximab, bevacizumab, imatinib and gefitinib inhibitors have illustrated the interest of targeting this protein class for treatment of selected cancers.

c-Met, is the prototypic member of a sub-family of RTKs which also includes RON and SEA. The c-Met RTK family is structurally different from other RTK families and is the only known high-affinity receptor for hepatocyte growth factor (HGF), also called scatter factor (SF) [D.P. Bottaro et al., Science 1991, 251:802-804; L. Naldini et al., Eur. Mol. Biol. Org. J. 1991, 10:2867-2878]. c-Met and HGF are widely expressed in a variety of tissue and their expression is normally restricted to cells of epithelial and mesenchymal origin respectively [M.F. Di Renzo et al., Oncogene 1991, 6:1997-2003; E. Sonnenberg et al., J. Cell. Biol. 1993, 123:223-235]. They are both required for normal mammalian development and have been shown to be particularly important in cell migration, morphogenic differentiation, and organization of the three-dimensional tubular structures as well as growth and angiogenesis [F. Baldt et al., Nature 1995, 376:768-771; C. Schmidt et al., Nature. 1995:373:699-702; Tsarfaty et al., Science 1994, 263:98-101]. While the controlled regulation of c-Met and HGF have been shown

to be important in mammalian development, tissue maintenance and repair [Nagayama T., Nagayama M., Kohara S., Kamiguchi H., Shibuya M., Katoh Y., Itoh J., Shinohara Y., Brain Res. 2004, 5;999(2):155-66; Tahara Y., Ido A., Yamamoto S., Miyata Y., Uto H., Hori T., Hayashi K., Tsubouchi H., J Pharmacol Exp Ther. 2003, 307(1):146-51],  
5 their dysregulation is implicated in the progression of cancers.

Aberrant signalling driven by inappropriate activation of c-Met is one of the most frequent alteration observed in human cancers and plays a crucial role in tumorigenesis and metastasis [Birchmeier et al., Nat. Rev. Mol. Cell Biol. 2003, 4:915-925; L. Trusolino and Comoglio P. M., Nat Rev. Cancer. 2002, 2(4):289-300].

10 Inappropriate c-Met activation can arise by ligand-dependent and independent mechanisms, which include overexpression of c-Met, and/or paracrine or autocrine activation, or through gain in function mutation [J.G. Christensen, Burrows J. and Salgia R., Cancer Letters. 2005, 226:1-26]. However an oligomerization of c-Met receptor, in presence or in absence of the ligand, is required to regulate the binding  
15 affinity and binding kinetics of the kinase toward ATP and tyrosine-containing peptide substrates [Hays JL, Watowich SJ, Biochemistry, 2004 Aug 17, 43:10570-8]. Activated c-Met recruits signalling effectors to its multidocking site located in the cytoplasm domain, resulting in the activation of several key signalling pathways, including Ras-MAPK, PI3K, Src and Stat3 [Gao CF, Vande Woude GF, Cell Res. 2005, 15(1):49-51;  
20 Furge KA, Zhang YW, Vande Woude GF, Oncogene. 2000, 19(49):5582-9]. These pathways are essential for tumour cell proliferation, invasion and angiogenesis and for evading apoptosis [Furge KA, Zhang YW, Vande Woude GF, Oncogene, 2000, 19(49):5582-9; Gu H., Neel BG, Trends Cell Biol. 2003 Mar, 13(3):122-30; Fan S., Ma YX, Wang JA, Yuan RQ, Meng Q., Cao Y., Laterra JJ, Goldberg ID, Rosen EM,  
25 Oncogene. 2000 Apr 27, 19(18):2212-23]. In addition, a unique facet of the c-Met signalling relative to other RTK is its reported interaction with focal adhesion complexes and non kinase binding partners such as  $\alpha 6\beta 4$  integrins [Trusolino L., Bertotti A., Comoglio PM, Cell. 2001, 107:643-54], CD44v6 [Van der Voort R., Taher TE, Wielenga VJ, Spaargaren M., Prevo R., Smit L., David G., Hartmann G., Gherardi E., Pals ST, J. Biol. Chem. 1999, 274(10):6499-506], Plexin B1 or semaphorins [Giordano S., Corso S., Conrotto P., Artigiani S., Gilestro G., Barberis D., Tamagnone L., Comoglio PM, Nat Cell Biol. 2002, 4(9):720-4; Conrotto P., Valdembri D., Corso

S., Serini G., Tamagnone L., Comoglio PM, Bussolino F., Giordano S., Blood. 2005, 105(11):4321-9; Conrotto P., Corso S., Gamberini S., Comoglio PM, Giordano S., Oncogene. 2004, 23:5131-7] which may further add to the complexity of regulation of cell function by this receptor. Finally recent data demonstrate that c-Met could be  
5 involved in tumor resistance to gefitinib or erlotinib suggesting that combination of compound targeting both EGFR and c-Met might be of significant interest [Engelman JA et al., Science, 2007, 316:1039-43].

In the past few years, many different strategies have been developed to attenuate c-Met signalling in cancer cell lines. These strategies include i) neutralizing antibodies  
10 against c-Met or HGF/SF [Cao B., Su Y., Oskarsson M., Zhao P., Kort EJ, Fisher RJ, Wang LM, Vande Woude GF, Proc Natl Acad Sci U S A. 2001, 98(13):7443-8; Martens T., Schmidt NO, Eckerich C., Fillbrandt R., Merchant M., Schwall R., Westphal M., Lamszus K., Clin Cancer Res. 2006, 12(20):6144-52] or the use of HGF/SF antagonist NK4 to prevent ligand binding to c-Met [Kuba K., Matsumoto K.,  
15 Date K., Shimura H., Tanaka M., Nakamura T., Cancer Res., 2000, 60:6737-43], ii) small ATP binding site inhibitors to c-Met that block kinase activity [Christensen JG, Schreck R., Burrows J., Kuruganti P., Chan E, Le P., Chen J., Wang X., Ruslim L., Blake R., Lipson KE, Ramphal J., Do S., Cui JJ, Cherrington JM, Mendel DB, Cancer Res. 2003, 63:7345-55], iii) engineered SH2 domain polypeptide that interferes with  
20 access to the multidocking site and RNAi or ribozyme that reduce receptor or ligand expression. Most of these approaches display a selective inhibition of c-Met resulting in tumor inhibition and showing that c-Met could be of interest for therapeutic intervention in cancer.

Within the molecules generated for c-Met targeting, some are antibodies. The  
25 most extensively described is the anti-c-Met 5D5 antibody generated by Genentech [WO 96/38557] which behaves as a potent agonist when added alone in various models and as an antagonist when used as a Fab fragment. A monovalent engineered form of this antibody described as one armed 5D5 (OA5D5) and produced as a recombinant protein in *E. Coli* is also the subject of a patent application [WO 2006/015371] by  
30 Genentech. However, this molecule that could not be considered as an antibody because of its particular scaffold, displays also mutations that could be immunogenic in humans. In terms of activity, this unglycosylated molecule is devoided of effector functions and

finally, no clear data demonstrate that OA5D5 inhibits dimerization of c-Met. Moreover, when tested in the G55 *in vivo* model, a glioblastoma cell line that expresses c-Met but not HGF mRNA and protein and that grows independently of the ligand, the one armed anti-c-Met had no significant effect on G55 tumor growth suggesting that  
5 OA5D5 acts primarily by blocking HGF binding and is not able to target tumors activated independently of HGF [Martens T. et al, Clin. Cancer Res., 2006, 12(20):6144-6152].

Another antibody targeting c-Met is described by Pfizer as an antibody acting “predominantly as c-Met antagonist, and in some instance as a c-Met agonist” [WO  
10 2005/016382]. No data showing any effect of Pfizer antibodies on c-Met dimerization is described in this application.

One of the innovative aspects of the present invention is to generate a chimeric and/or humanized monoclonal antibody without intrinsic agonist activity and inhibiting c-Met dimerization. More particularly, an innovative aspect of the present invention is  
15 to generate a chimeric and/or humanized monoclonal antibody with antagonist activity and inhibiting c-Met dimerization.

In addition of targeting ligand-dependent tumors, this approach will also impair ligand-independent activations of c-Met due to its overexpression or mutations of the intra cellular domains which remained dependent to oligomerization for signalling.  
20 Another aspect of the activity of this antibody could be a steric hindrance for c-Met interaction with its partners that will result in impairment of c-Met functions. This antibody is humanized and engineered preferentially, but not limited, as human IgG1 to get effector functions such as ADCC and CDC in addition to functions linked to the specific blockade of the c-Met receptor.

25 Surprisingly, for the first time, inventors have managed to generate a chimeric and/or humanized monoclonal antagonist antibody capable of binding to c-Met but also capable of inhibiting the c-Met dimerization, said monoclonal antibody being divalent contrary to existing antagonist antibodies directed against c-Met. If it is true that, in the prior art, it is sometimes suggested that an antibody capable of inhibiting the  
30 dimerization of c-Met with its partners could be an interesting one, it has never been disclosed, or clearly suggested, an antibody capable of doing so. Moreover, regarding antibody specificity, it was not evident at all to succeed in the generation of such an

active divalent antibody.

As it was explained before, the inhibition of the c-Met dimerization is a capital aspect of the invention as such antibodies will present a real interest for a larger population of patients. Not only ligand-dependent activated c-Met cancer, as it was the case until the present invention, but also ligand-independent activated c-Met cancer could be treated with antibodies generated by the process of the present invention.

Antibodies were evaluated by BRET analysis on cells expressing both c-Met-RLuc/c-Met-YFP and selected on their ability to inhibit at least 40 %, preferably 45 %, 50 %, 55 % and most preferably 60 % of the BRET signal.

The BRET technology is known as being representative of the protein dimerization [Angers et al., PNAS, 2000, 97:3684-89].

The BRET technology is well known by the man skilled in the art and will be detailed in the following examples. More particularly, BRET (Bioluminescence Resonance Energy Transfer) is a non-radiative energy transfer occurring between a bioluminescent donor (Renilla Luciferase (Rluc)) and a fluorescent acceptor, a mutant of GFP (Green Fluorescent Protein) or YFP (Yellow fluorescent protein). In the present case EYFP (Enhanced Yellow Fluorescent Protein) was used. The efficacy of transfer depends on the orientation and the distance between the donor and the acceptor. Then, the energy transfer can occur only if the two molecules are in close proximity (1-10 nm). This property is used to generate protein-protein interaction assays. Indeed, in order to study the interaction between two partners, the first one is genetically fused to the Renilla Luciferase and the second one to the yellow mutant of the GFP. Fusion proteins are generally, but not obligatory, expressed in mammalian cells. In presence of its membrane permeable substrate (coelenterazine), Rluc emits blue light. If the GFP mutant is closer than 10 nm from the Rluc, an energy transfer can occur and an additional yellow signal can be detected. The BRET signal is measured as the ratio between the light emitted by the acceptor and the light emitted by the donor. So the BRET signal will increase as the two fusion proteins are brought into proximity or if a conformational change brings Rluc and GFP mutant closer.

If the BRET analysis consists in a preferred embodiment, any method known by the man skilled in the art can be used to measure c-Met dimerization. Without limitation, the following technologies can be mentioned: FRET (Fluorescence

Resonance Energy Transfer), HTRF (Homogenous Time resolved Fluorescence), FLIM (Fluorescence Lifetime Imaging Microscopy) or SW-FCCS single wavelength fluorescence cross-correlation spectroscopy).

Other classical technologies could also be used, such as Co-immunoprecipitation, Alpha screen, Chemical cross-linking, Double-Hybrid, Affinity Chromatography, ELISA or Far western blot.

The terms “antibody”, “antibodies” or “immunoglobulin” are used interchangeably in the broadest sense and include monoclonal antibodies (e.g., full length or intact monoclonal antibodies), polyclonal antibodies, multivalent antibodies or multispecific antibodies (e.g., bispecific antibodies so long as they exhibit the desired biological activity).

More particularly, such molecule consists in a glycoprotein comprising at least two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds. Each heavy chain is comprised of a heavy chain variable region (or domain) (abbreviated herein as HCVR or VH) and a heavy chain constant region. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein as LCVR or VL) and a light chain constant region. The light chain constant region is comprised of one domain, CL. The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR). Each VH and VL is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. The variable regions of the heavy and light chains contain a binding domain that interacts with an antigen. The constant regions of the antibodies may mediate the binding of the immunoglobulin to host tissues or factors, including various cells of the immune system (e.g. effector cells) and the first component (C1q) of the classical complement system.

The heavy chains of immunoglobulins can be divided into three functional regions: the Fd region, the hinge region, and the Fc region (fragment crystallizable). The Fd region comprises the VH and CH1 domains and, in combination with the light chain, forms Fab - the antigen-binding fragment. The Fc fragment is responsible for the immunoglobulin effector functions, which includes, for example, complement fixation



and binding to cognate Fc receptors of effector cells. The hinge region, found in IgG, IgA, and IgD immunoglobulin classes, acts as a flexible spacer that allows the Fab portion to move freely in space relative to the Fc region. The hinge domains are structurally diverse, varying in both sequence and length among immunoglobulin classes and subclasses.

According to crystallographic studies, the immunoglobulin hinge region can be further subdivided structurally and functionally into three regions: the upper hinge, the core, and the lower hinge (Shin et al., Immunological Reviews 130:87, 1992). The upper hinge includes amino acids from the carboxyl end of CH1 to the first residue in the hinge that restricts motion, generally the first cysteine residue that forms an interchain disulfide bond between the two heavy chains. The length of the upper hinge region correlates with the segmental flexibility of the antibody. The core hinge region contains the inter-heavy chain disulfide bridges. The lower hinge region joins the amino terminal end of, and includes residues in the CH2 domain. The core hinge region of human IgG1 contains the sequence Cys-Pro-Pro-Cys that, when dimerized by disulfide bond formation, results in a cyclic octapeptide believed to act as a pivot, thus conferring flexibility. Conformational changes permitted by the structure and flexibility of the immunoglobulin hinge region polypeptide sequence may affect the effector functions of the Fc portion of the antibody.

The term « Monoclonal Antibody » is used in accordance with its ordinary meaning to denote an antibody obtained from a population of substantially homogeneous antibodies, i.e. the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. In other words, a monoclonal antibody consists in a homogenous antibody resulting from the proliferation of a single clone of cells (e.g., hybridoma cells, eukaryotic host cells transfected with DNA encoding the homogenous antibody, prokaryotic host cells transformed with DNA encoding the homogenous antibody, etc.), and which is generally characterized by heavy chains of a single class and subclass, and light chains of a single type. Monoclonal antibodies are highly specific, being directed against a single antigen. Furthermore, in contrast to polyclonal antibodies preparations that typically include different antibodies directed against different determinants, or

epitope, each monoclonal antibody is directed against a single determinant on the antigen.

In the present description, the terms polypeptides, polypeptide sequences, amino acid sequences, peptides and proteins attached to antibody compounds or to their sequence are interchangeable.

5 The invention relates to a monoclonal antibody, or a divalent functional fragment or derivative thereof, capable to inhibit the c-Met dimerization, said antibody comprising a heavy chain comprising CDR-H1, CDR-H2 and CDR-H3 with respectively the amino acid sequences SEQ ID Nos. 1, 2 and 3; and a light chain comprising CDR-L1, CDR-L2 and  
10 CDR-L3 with respectively the amino acid sequences SEQ ID Nos. 5, 6 and 7, said antibody being further characterized in that it also comprises a hinge region comprising the amino acid sequence selected from the group consisting of SEQ ID Nos. 22 to 28 and SEQ ID Nos. 58 to 72.

The invention further relates to a monoclonal antibody, or a divalent functional fragment or derivative thereof, capable to inhibit the c-Met dimerization and comprising a  
15 heavy chain comprising CDR-H1, CDR-H2 and CDR-H3 with respectively the amino acid sequences SEQ ID Nos. 1, 2 and 3 or a sequence having at least 80 % identity after optimum alignment with sequences SEQ ID Nos. 1, 2 and 3; and a light chain comprising CDR-L1, CDR-L2 and CDR-L3 with respectively the amino acid sequences SEQ ID Nos. 5, 6 and 7 or a  
20 sequence having at least 80% identity after optimum alignment with sequences SEQ ID Nos. 5, 6 or 7, said antibody being further characterized in that it also comprises a hinge region comprising the amino acid sequence SEQ ID No. 56.

More particularly, the invention relates to a monoclonal antibody, or a divalent functional fragment or derivative thereof, as above described characterized in that it also comprises a hinge region comprising the amino acid sequence SEQ ID No. 57.

25 In other words, the invention relates to a monoclonal antibody, or a divalent functional fragment or derivative thereof, capable to inhibit the c-Met dimerization and comprising a heavy chain comprising CDR-H1, CDR-H2 and CDR-H3 with respectively the amino acid sequences SEQ ID Nos. 1, 2 and 3 or a sequence having at least 80% identity after optimum alignment with sequences SEQ ID Nos. 1, 2 and 3; and a light chain comprising CDR-L1,

CDR-L2 and CDR-L3 with respectively the amino acid sequences SEQ ID Nos. 5, 6 and 7 or a sequence having at least 80% identity after optimum alignment with sequences SEQ ID Nos. 5, 6 or 7, said antibody being further characterized in that it also comprises a hinge region comprising the amino acid sequence SEQ ID No. 57.

5 More particularly, the invention relates to a monoclonal antibody, or a divalent functional fragment or derivative thereof, as above described characterized in that it also comprises a hinge region comprising the amino acid sequence SEQ ID No. 21.

In other words, the invention also relates to a monoclonal antibody, or a divalent functional fragment or derivative thereof, capable to inhibit the c-Met dimerization and comprising a heavy chain comprising CDR-H1, CDR-H2 and CDR-H3 with respectively the amino acid sequences SEQ ID Nos. 1, 2 and 3 or a sequence having at least 80% identity after optimum alignment with sequences SEQ ID Nos. 1, 2 and 3; and a light chain comprising CDR-L1, CDR-L2 and CDR-L3 with respectively the amino acid sequences SEQ ID Nos. 5, 6 and 7 or a sequence having at least 80% identity after optimum alignment with sequences SEQ ID Nos. 5, 6 or 7, said antibody being further characterized in that it also comprises a hinge region comprising the amino acid sequence SEQ ID No. 21.

As it will be apparent for the man skilled in the art, the consensus sequences SEQ ID Nos. 57 and 21 are comprised in the consensus sequence SEQ ID No. 56.

Table 1

	#01	#02	#03	#04	#05	#06	#07	#08	#09	#10	#11	#12	#13	#14
SEQ ID NO 56	x1	x2	x3	C	x5	x6	x7	x8	x9	C	x11	x12	C	x14
SEQ ID NO 57	x1	x2	x3	C	x5	x6	x7	x8	x9	C	P	P	C	P
SEQ ID NO 21	x1	x2	x3	C	x5	-	C	x8	x9	C	x11	x12	C	x14

15

For SEQ ID No. 56:

X1: P, R, C, -      X5: D, C, G, -      X8: H, V, K, -      X12: P, -  
 X2: K, C, R, -      X6: K, C, -      X9: T, C, E, P, -      X14: P, T  
 X3: S, C, D, -      X7: T, C, -      X11: P, I

20

The expression “functional fragments and derivatives” will be defined in details later in the present specification.

By CDR regions or CDR(s), it is intended to indicate the hypervariable regions of the heavy and light chains of the immunoglobulins as defined by IMGT.

25

The IMGT unique numbering has been defined to compare the variable domains whatever the antigen receptor, the chain type, or the species [Lefranc M.-P., Immunology Today 18, 509 (1997); Lefranc M.-P., The Immunologist, 7, 132-136 (1999); Lefranc, M.-P., Pommié, C., Ruiz, M., Giudicelli, V., Foulquier, E., Truong, L., Thouvenin-Contet, V. and Lefranc, Dev. Comp. Immunol., 27, 55-77 (2003)]. In the

IMGT unique numbering, the conserved amino acids always have the same position, for instance cysteine 23 (1st-CYS), tryptophan 41 (CONSERVED-TRP), hydrophobic amino acid 89, cysteine 104 (2nd-CYS), phenylalanine or tryptophan 118 (J-PHE or J-TRP). The IMGT unique numbering provides a standardized delimitation of the framework regions (FR1-IMGT: positions 1 to 26, FR2-IMGT: 39 to 55, FR3-IMGT: 66 to 104 and FR4-IMGT: 118 to 128) and of the complementarity determining regions: CDR1-IMGT: 27 to 38, CDR2-IMGT: 56 to 65 and CDR3-IMGT: 105 to 117. As gaps represent unoccupied positions, the CDR-IMGT lengths (shown between brackets and separated by dots, e.g. [8.8.13]) become crucial information. The IMGT unique numbering is used in 2D graphical representations, designated as IMGT Colliers de Perles [Ruiz, M. and Lefranc, M.-P., *Immunogenetics*, 53, 857-883 (2002); Kaas, Q. and Lefranc, M.-P., *Current Bioinformatics*, 2, 21-30 (2007)], and in 3D structures in IMGT/3Dstructure-DB [Kaas, Q., Ruiz, M. and Lefranc, M.-P., T cell receptor and MHC structural data. *Nucl. Acids. Res.*, 32, D208-D210 (2004)].

Three heavy chain CDRs and 3 light chain CDRs exist. The term CDR or CDRs is used here in order to indicate, according to the case, one of these regions or several, or even the whole, of these regions which contain the majority of the amino acid residues responsible for the binding by affinity of the antibody for the antigen or the epitope which it recognizes.

By "percentage of identity" between two nucleic acid or amino acid sequences in the sense of the present invention, it is intended to indicate a percentage of nucleotides or of identical amino acid residues between the two sequences to be compared, obtained after the best alignment (optimum alignment), this percentage being purely statistical and the differences between the two sequences being distributed randomly and over their entire length. The comparisons of sequences between two nucleic acid or amino acid sequences are traditionally carried out by comparing these sequences after having aligned them in an optimum manner, said comparison being able to be carried out by segment or by "comparison window". The optimum alignment of the sequences for the comparison can be carried out, in addition to manually, by means of the local homology algorithm of Smith and Waterman (1981) [Ad. App. Math. 2:482], by means of the local homology algorithm of Needleman and Wunsch (1970) [J. Mol. Biol. 48: 443], by means of the similarity search method of Pearson and Lipman

(1988) [Proc. Natl. Acad. Sci. USA 85:2444), by means of computer software using these algorithms (GAP, BESTFIT, FASTA and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI, or else by BLAST N or BLAST P comparison software).

5           The percentage of identity between two nucleic acid or amino acid sequences is determined by comparing these two sequences aligned in an optimum manner and in which the nucleic acid or amino acid sequence to be compared can comprise additions or deletions with respect to the reference sequence for an optimum alignment between these two sequences. The percentage of identity is calculated by determining the  
10           number of identical positions for which the nucleotide or the amino acid residue is identical between the two sequences, by dividing this number of identical positions by the total number of positions in the comparison window and by multiplying the result obtained by 100 in order to obtain the percentage of identity between these two sequences.

15           For example, it is possible to use the BLAST program, "BLAST 2 sequences" (Tatusova et al., "Blast 2 sequences - a new tool for comparing protein and nucleotide sequences", FEMS Microbiol Lett. 174:247-250) available on the site <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>, the parameters used being those given by default (in particular for the parameters "open gap penalty": 5, and "extension gap  
20           penalty": 2; the matrix chosen being, for example, the matrix "BLOSUM 62" proposed by the program), the percentage of identity between the two sequences to be compared being calculated directly by the program.

          By amino acid sequence having at least 80 %, preferably 85 %, 90 %, 95 % and  
25           98 % identity with a reference amino acid sequence, those having, with respect to the reference sequence, certain modifications, in particular a deletion, addition or substitution of at least one amino acid, a truncation or an elongation are preferred. In the case of a substitution of one or more consecutive or nonconsecutive amino acid(s), the substitutions are preferred in which the substituted amino acids are replaced by "equivalent" amino acids. The expression "equivalent amino acids" is aimed here at  
30           indicating any amino acid capable of being substituted with one of the amino acids of the base structure without, however, essentially modifying the biological activities of the corresponding antibodies and such as will be defined later, especially in the

examples. These equivalent amino acids can be determined either by relying on their structural homology with the amino acids which they replace, or on results of comparative trials of biological activity between the different antibodies capable of being carried out.

- 5 By way of example, mention is made of the possibilities of substitution capable of being carried out without resulting in a profound modification of the biological activity of the corresponding modified antibody.

As non limitative example, the following table 2 is giving substitution possibilities conceivable with a conservation of the biological activity of the modified  
10 antibody. The reverse substitutions are also, of course, possible in the same conditions.

**Table 2**

<b>Original residue</b>	<b>Substitution(s)</b>
Ala (A)	Val, Gly, Pro
Arg (R)	Lys, His
Asn (N)	Gln
Asp (D)	Glu
Cys (C)	Ser
Gln (Q)	Asn
Glu (G)	Asp
Gly (G)	Ala
His (H)	Arg
Ile (I)	Leu
Leu (L)	Ile, Val, Met
Lys (K)	Arg
Met (M)	Leu
Phe (F)	Tyr
Pro (P)	Ala
Ser (S)	Thr, Cys
Thr (T)	Ser
Trp (W)	Tyr
Tyr (Y)	Phe, Trp
Val (V)	Leu, Ala

15 It must be understood here that the invention does not relate to the antibodies in natural form, that is to say they are not in their natural environment but that they have been able to be isolated or obtained by purification from natural sources, or else obtained by genetic recombination, or by chemical synthesis, and that they can then contain unnatural amino acids as will be described further on.

It must also be understood, as previously mentioned, that the invention concerns more particularly a chimeric and/or a humanized divalent antibody, or any divalent functional fragment or derivative, with an antagonistic activity. Divalent antibodies of the prior art are agonists or partial agonists. The monoclonal antibody of the invention, including a modified hinge as previously described, i.e. including a hinge region comprising the amino acid sequence SEQ ID No. 56, 57 or 21, is novel and presents the particularity to have an improved antagonistic activity compared to the chimeric or humanized antibody 224G11 without such a modified hinge as it will appear from the following examples.

Contrary to the prior art, inventors have obtained an improved antagonistic activity without modifying the format of the antibody. Actually, in the closest prior art represented by the antibody 5D5, it has been necessary to develop a monovalent fragment of the antibody to generate an antagonistic activity. In the present application, by the use of the hinge of the invention, it is possible for the first time to obtain a full divalent antibody with increased antagonistic activity, and this contrary to the general knowledge.

In a preferred embodiment, the antibody of the invention comprises a hinge region comprising an amino acid sequence selected from the group consisting of SEQ ID Nos. 22 to 28 and 58 to 72, or a sequence having at least 80% identity after optimum alignment with sequences SEQ ID Nos. 22 to 28 and 58 to 72.

For more clarity, the following tables 3 and 4 regroup the amino acids and nucleotides sequences of the different preferred hinges of the invention.

**Table 3**

25

SEQ ID No.	Amino acids	SEQ ID No.	Nucleotides
22	RKCCVECP	29	AGGAAGTGCTGTGTGGAGTGCCCCCCTGCCCA
23	PRDCGCKPCICT	30	CCCCGGGACTGTGGGTGCAAGCCTTGCATTTGTACC
24	PKSCGCKPCICT	31	CCCAAGAGCTGTGGGTGCAAGCCTTGCATTTGTACC
25	PKSCGCKPCICP	32	CCAAAGAGCTGCGGCTGCAAGCCTTGTATCTGTCCC
26	PRDCGCKPCPPCP	33	CCACGGGACTGTGGCTGCAAGCCTGCCCTCCGTGTCCA
27	PRDCGCHTCPPCP	34	CCCAGAGACTGTGGGTGTACACCTGCCCTCCTGTCCCT
28	PKSCDCHCPCPPCP	35	CCCAAAGCTGCGATTGCCACTGTCCCTCCATGTCCA



Table 4

SEQ ID No.	Amino acids	SEQ ID No.	Nucleotides
58	CKSCDKTHTCPPCP	73	TGCAAGAGCTGCGACAAGACCCACACCTGTCCCCCCTGCCCT
59	PCSCDKTHTCPPCP	74	CCCTGCAGCTGCGACAAGACCCACACCTGTCCCCCCTGCCCT
60	PKCCDKTHTCPPCP	75	CCCAAGTGCTGCGACAAGACCCACACCTGTCCCCCCTGCCCT
61	PKSCCKTHTCPPCP	76	CCTAAGAGCTGTTGCAAGACCCACACCTGTCCCCCCTGCCCT
62	PKSCDCTHTCPPCP	77	CCCAAGAGCTGCGACTGCACCCACACCTGTCCCCCCTGCCCT
63	PKSCDKCHTCPPCP	78	CCCAAGAGCTGCGACAAGTGCCACACCTGTCCCCCCTGCCCT
64	PKSCDKTHCCPPCP	79	CCCAAGAGCTGCGACAAGACCCACTGCTGTCCCCCCTGCCCT
65	KCDKTHTCPPCP	80	AAGTGCAGACAAGACCCACACCTGTCCCCCCTGCCCT
66	PKSCDCHTCPPCP	81	CCCAAGAGCTGCGACTGCCACACCTGTCCCCCCTGCCCT
67	PKSCDCTHCCPPCP	82	CCCAAGAGCTGCGACTGCACCCACTGTCCCCCCTGCCCT
68	PCSKHTCPPCP	83	CCCTGCAGCTGCAAGCACACCTGTCCCCCCTGCCCT
69	PSCCTHTCPPCP	84	CCTAGCTGCTGCACCCACACCTGTCCCCCCTGCCCT
70	PSCDKHCCPPCP	85	CCCAGCTGCGACAAGCACTGCTGTCCCCCCTGCCCT
71	PKSCTCPPCP	86	CCCAAGAGCTGCACCTGTCCCCCTTGTCCT
72	PKSCDKCVECPCP	87	CCCAAGAGCTGCGATAAGTGCGTGGAGTGCCCCCCTTGTCCT

According a first approach, the antibody will be defined by its heavy chain  
5 sequence. More particularly, the antibody of the invention, or one of its functional  
fragments or derivatives, is characterized in that it comprises a heavy chain comprising  
at least one CDR chosen from CDRs comprising the amino acid sequences SEQ ID  
Nos. 1 to 3.

The mentioned sequences are the following ones:

- 10 SEQ ID No. 1: GYIFTAYT  
SEQ ID No. 2: IKPNNGLA  
SEQ ID No. 3: ARSEITTEFDY

According to a preferred aspect, the antibody of the invention, or one of its  
15 functional fragments or derivatives, comprises a heavy chain comprising at least one,  
preferably two, and most preferably three, CDR(s) chosen from CDR-H1, CDR-H2 and  
CDR-H3, wherein:

- CDR-H1 comprises the amino acid sequence SEQ ID No. 1,  
- CDR-H2 comprises the amino acid sequence SEQ ID No. 2,  
20 - CDR-H3 comprises the amino acid sequence SEQ ID No. 3.

In a second approach, the antibody will be now defined by its light chain  
sequence. More particularly, according to a second particular aspect of the invention,

the antibody, or one of its functional fragments or derivatives, is characterized in that it comprises a light chain comprising at least one CDR chosen from CDRs comprising the amino acid sequence SEQ ID Nos. 5 to 7.

The mentioned sequences are the following ones:

- 5        SEQ ID No. 5:     ESVDSYANSF  
          SEQ ID No. 6:     RAS  
          SEQ ID No. 7:     QQSKEDPLT

10        According to another preferred aspect, the antibody of the invention, or one of its functional fragments or derivatives, comprises a light chain comprising at least one, preferably two, and most preferably three, CDR(s) chosen from CDR-L1, CDR-L2 and CDR-L3, wherein:

- CDR-L1 comprises the amino acid sequence SEQ ID No. 5,
- CDR-L2 comprises the amino acid sequence SEQ ID No. 6,
- 15        - CDR-L3 comprises the amino acid sequence SEQ ID No. 7.

The murine hybridoma capable of secreting monoclonal antibodies according to the present invention, especially hybridoma of murine origin, was deposited at the CNCM (Institut Pasteur, Paris, France) on 03/14/2007 under the number CNCM I-3731.

20        In the present application, IgG1 are preferred to get effector functions, and most preferably ADCC and CDC.

The skilled artisan will recognize that effector functions include, for example, C1q binding; complement dependent cytotoxicity (CDC); Fc receptor binding; antibody-dependent cell-mediated cytotoxicity (ADCC); phagocytosis; and down regulation of cell surface receptors (e.g. B cell receptor; BCR).

25        The antibodies according to the present invention, are preferably specific monoclonal antibodies, especially of murine, chimeric or humanized origin, which can be obtained according to the standard methods well known to the person skilled in the art.

30        In general, for the preparation of monoclonal antibodies or their functional fragments or derivatives, especially of murine origin, it is possible to refer to techniques which are described in particular in the manual "Antibodies" (Harlow and Lane, Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor

NY, pp. 726, 1988) or to the technique of preparation from hybridomas described by Kohler and Milstein (Nature, 256:495-497, 1975).

The monoclonal antibodies according to the invention can be obtained, for example, from an animal cell immunized against the c-Met, or one of its fragments  
5 containing the epitope specifically recognized by said monoclonal antibodies according to the invention. Said c-Met, or one of its said fragments, can especially be produced according to the usual working methods, by genetic recombination starting with a nucleic acid sequence contained in the cDNA sequence coding for the c-Met or by peptide synthesis starting from a sequence of amino acids comprised in the peptide  
10 sequence of the c-Met.

The monoclonal antibodies according to the invention can, for example, be purified on an affinity column on which the c-Met or one of its fragments containing the epitope specifically recognized by said monoclonal antibodies according to the invention has previously been immobilized. More particularly, said monoclonal  
15 antibodies can be purified by chromatography on protein A and/or G, followed or not followed by ion-exchange chromatography aimed at eliminating the residual protein contaminants as well as the DNA and the LPS, in itself followed or not followed by exclusion chromatography on Sepharose™ gel in order to eliminate the potential aggregates due to the presence of dimers or of other multimers. In an even more  
20 preferred manner, the whole of these techniques can be used simultaneously or successively.

The antibody of the invention, or a divalent functional fragment or derivative thereof, consists preferably of a chimeric antibody.

By chimeric antibody, it is intended to indicate an antibody which contains a  
25 natural variable (light chain and heavy chain) region derived from an antibody of a given species in combination with the light chain and heavy chain constant regions of an antibody of a species heterologous to said given species (e.g. mouse, horse, rabbit, dog, cow, chicken, etc.).

The antibodies or their fragments of chimeric type according to the invention  
30 can be prepared by using the techniques of genetic recombination. For example, the chimeric antibody can be produced by cloning a recombinant DNA containing a promoter and a sequence coding for the variable region of a non-human, especially

murine, monoclonal antibody according to the invention and a sequence coding for the constant region of human antibody. A chimeric antibody of the invention encoded by such a recombinant gene will be, for example, a mouse-man chimera, the specificity of this antibody being determined by the variable region derived from the murine DNA and its isotype determined by the constant region derived from the human DNA. For the methods of preparation of chimeric antibodies, it is possible, for example, to refer to the documents Verhoeyn et al. (BioEssays, 8:74, 1988), Morrison et al. (Proc. Natl. Acad. Sci. USA 82:6851-6855, 1984) or US 4,816,567.

More particularly, said antibody, or a functional fragment or derivative thereof, comprises a chimeric heavy chain variable domain of sequence comprising the amino acid sequence SEQ ID No. 46 or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 46.

SEQ ID No. 46: EVQLQQSGPELVKPGASVKISKTSQYIFTAYTMHWVRQSLGE  
SLDWIGGIKPNNGLANYNQKFKGKATLTVDKSSSTAYMDLRSLTSEDSAVYYC  
ARSEITTEFDYWGQGTALTVSS

More particularly, said antibody, or a functional fragment or derivative thereof, comprises a chimeric light chain variable domain of sequence comprising the amino acid sequence SEQ ID No. 47 or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 47.

SEQ ID No. 47: DIVLTQSPASLAVSLGQRATISCRASESVDSYANSFMHWYQQKP  
GQPPKLLIYRASNLESGIPARFSGSGSRTDFTLTINPVEADDVATYYCQQSKEDP  
LTFGSGTKLEMKR

More particularly, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [IgG2chim], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 22.

In the present application, the use of square brackets is not necessary and, as an example, the reference [224G11] [IgG2chim] must be considered as identical to 224G11IgG2chim. In a same way, to indicate that the antibody is a murine one, the

expression murine or the letter m can be added; to indicate that the antibody is a chimeric one, the expression chim or the letter c can be added and ; to indicate that the antibody is a humanized one, the expression hum, hz, Hz or the letter h can be added. As an example, the chimeric antibody 224G11IgG2 can be referred as c224G11IgG2,  
5 c224G11[IgG2], c[224G11]IgG2, c[224G11][IgG2], 224G11IgG2chim, 224G11[IgG2chim], [224G11]IgG2chim or [224G11][IgG2chim].

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [TH7chim], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID  
10 No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 28.

In the present application, the reference TH7 must be considered as identical to C7Δ6-9 or TH7C7Δ6-9. The symbol Δ means deletion.

In another aspect, a preferred antibody, or a divalent functional fragment or  
15 derivative thereof, according to the invention and named [224G11] [MHchim], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 23.

In another aspect, a preferred antibody, or a divalent functional fragment or  
20 derivative thereof, according to the invention and named [224G11] [MUP9Hchim], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 26.

In another aspect, a preferred antibody, or a divalent functional fragment or  
25 derivative thereof, according to the invention and named [224G11] [MMCHchim], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 24.

In another aspect, a preferred antibody, or a divalent functional fragment or  
30 derivative thereof, according to the invention and named [224G11] [C1], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a

light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 58.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [C2], comprises a  
5 heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 59.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [C3], comprises a  
10 heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 60.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [C5], comprises a  
15 heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 61.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [C6], comprises a  
20 heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 62.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [C7], comprises a  
25 heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 63.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [C9], comprises a  
30 heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 64.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [ $\Delta$ 1-3], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 65.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [C7 $\Delta$ 6], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 66.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [C6 $\Delta$ 9], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 67.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [C2 $\Delta$ 5-7], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 68.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [C5 $\Delta$ 2-6], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 69.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [C9 $\Delta$ 2-7], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 70.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [ $\Delta$ 5-6-7-8],

comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 71.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [IgG1/IgG2],  
5 comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 72.

The antibody of the invention, or a divalent functional fragment or derivative  
10 thereof, consists preferably of a human antibody.

The term "human antibody" includes all antibodies that have one or more variable and constant region derived from human immunoglobulin sequences. In a preferred embodiment, all of the variable and constant domains (or regions) are derived from human immunoglobulin sequence (fully human antibody). In other words, it  
15 includes any antibody which have variable and constant regions (if present) derived from human germline immunoglobulin sequences, i.e. which possesses an amino acid sequence which corresponds to that of an antibody produced by a human and/or has been made using any techniques for making human antibodies known by the man skill in the art.

In one embodiment, the human monoclonal antibodies are produced by a  
20 hybridoma which includes a B cell obtained from a transgenic non-human animal, e.g., a transgenic mouse, having a genome comprising a human heavy chain transgene and a light chain transgene fused to an immortalized cell.

As example for such transgenic mouse, it can be mentioned the  
25 XENOMOUSE<sup>TM</sup> which is an engineered mouse strain that comprises large fragments of the human immunoglobulin loci and is deficient in mouse antibody production (Green et al., 1994, Nature Genetics, 7:13-21). The XENOMOUSE<sup>TM</sup> produces an adult-like human repertoire of fully human antibodies, and generate antigen-specific human monoclonal antibodies. A second generation XENOMOUSE<sup>TM</sup> contains  
30 approximately 80% of the human antibody repertoire (Green & Jakobovits, 1998, J. Exp. Med., 188:483-495).



Any other technique known by the man skill in the art, such as phage display technique, can also be used for the generation of human antibody according to the invention.

The antibody of the invention, or a divalent functional fragment or derivative thereof, consists preferably of a humanized antibody.

By the expression "humanized antibody", it is intended to indicate an antibody which contains CDR regions derived from an antibody of nonhuman origin, the other parts of the antibody molecule being derived from one (or from several) human antibodies. Moreover, some of the residues of the segments of the skeleton (called FR) can be modified in order to conserve the affinity of the binding (Jones et al., Nature, 321:522-525, 1986; Verhoeyen et al., Science, 239:1534-1536, 1988; Riechmann et al., Nature, 332:323-327, 1988).

The humanized antibodies according to the invention or their fragments can be prepared by techniques known to the person skilled in the art (such as, for example, those described in the documents Singer et al., J. Immun. 150:2844-2857, 1992; Mountain et al., Biotechnol. Genet. Eng. Rev., 10: 1-142, 1992; or Bebbington et al., Bio/Technology, 10:169-175, 1992).

Other humanization methods are known by the man skill in the art as, for example, the "CDR Grafting" method described by Protein Design Lab (PDL) in the patent applications EP 0 451261, EP 0 682 040, EP 0 9127, EP 0 566 647 or US 5,530,101, US 6,180,370, US 5,585,089 and US 5,693,761. The following patent applications can also be mentioned: US 5,639,641; US 6,054,297; US 5,886,152 and US 5,877,293.

More particularly, said antibody, or a functional fragment or derivative thereof, comprises a humanized heavy chain variable domain of sequence comprising the amino acid sequence SEQ ID No. 4 or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 4.

SEQ ID No. 4: QVQLVQSGAEVKKPGASVKVSKASGYIFTAYTMHWVRQAPG  
QGLEWMGWIKPNNGLANYAQKFQGRVTMTRDTSISTAYMELSRLRSDDTAVY  
YCARSEITTEFDYWGQGTLVTVSS

More particularly, said antibody, or a functional fragment or derivative thereof, comprises a humanized light chain variable domain selected from the group of

sequences comprising the amino acid sequence SEQ ID No. 8, 9 or 10 or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 8, 9 or 10.

SEQ ID No. 8: DIVLTQSPDSLAVSLGERATINCKSSESVD SYANSFMHWYQQKP  
5 GQPPKLLIYRASTRESGVPDRFSGSGSRTDFTLTISLQAEDVAVYYCQQSKEDP  
LTFGGGTKVEIKR

SEQ ID No. 9: DIVMTQSPDSLAVSLGERATINCKSSESVD SYANSFMHWYQQKP  
GQPPKLLIYRASTRESGVPDRFSGSGSGTDFTLTISLQAEDVAVYYCQQSKEDP  
LTFGGGTKVEIKR

10 SEQ ID No. 10: DIVMTQSPDSLAVSLGERATINCKSSESVD SYANSFLHWYQQKP  
GQPPKLLIYRASTRESGVPDRFSGSGSGTDFTLTISLQAEDVAVYYCQQSKEDP  
LTFGGGTKVEIKR

More particularly, a preferred antibody, or a divalent functional fragment or  
15 derivative thereof, according to the invention and named [224G11] [IgG2Hz1],  
comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID  
No. 4, a light chain variable domain comprising the amino acid sequence SEQ ID No. 8,  
and a hinge region comprising the amino acid sequence SEQ ID No. 22.

In another aspect, a preferred antibody, or a divalent functional fragment or  
20 derivative thereof, according to the invention and named [224G11] [IgG2Hz2],  
comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID  
No. 4, a light chain variable domain comprising the amino acid sequence SEQ ID No. 9,  
and a hinge region comprising the amino acid sequence SEQ ID No. 22.

In another aspect, a preferred antibody, or a divalent functional fragment or  
25 derivative thereof, according to the invention and named [224G11] [IgG2Hz3],  
comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID  
No. 4, a light chain variable domain comprising the amino acid sequence SEQ ID  
No. 10, and a hinge region comprising the amino acid sequence SEQ ID No. 22.

In another aspect, a preferred antibody, or a divalent functional fragment or  
30 derivative thereof, according to the invention and named [224G11] [TH7Hz1],  
comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID

No. 4, a light chain variable domain comprising the amino acid sequence SEQ ID No. 8, and a hinge region comprising the amino acid sequence SEQ ID No. 28.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [TH7z2], comprises  
5 a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 4, a light chain variable domain comprising the amino acid sequence SEQ ID No. 9, and a hinge region comprising the amino acid sequence SEQ ID No. 28.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [TH7Hz3],  
10 comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 4, a light chain variable domain comprising the amino acid sequence SEQ ID No. 10, and a hinge region comprising the amino acid sequence SEQ ID No. 28.

In another aspect, antibodies of the invention can be described by their total heavy and light chains, respectively.

As example, the antibody [224G11] [IgG2chim] of the invention comprises a  
15 complete heavy chain comprising the amino acid sequence SEQ ID No. 50, or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 50, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80% identity after optimum alignment with the  
20 sequence SEQ ID No. 52.

As another example, the antibody [224G11] [TH7chim] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 51, or a sequence having at least 80% identity after optimum alignment with the sequence  
25 SEQ ID No. 51, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 52.

As another example, the antibody [224G11] [C1] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 88, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ  
30 ID No. 88, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 52.

As another example, the antibody [224G11] [C2] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 89, or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 89, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 52.

As another example, the antibody [224G11] [C3] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 90, or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 90, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 52.

As another example, the antibody [224G11] [C5] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 91, or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 91, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 52.

As another example, the antibody [224G11] [C6] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 92, or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 92, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 52.

As another example, the antibody [224G11] [C7] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 93, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 93, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 52.

As another example, the antibody [224G11] [C9] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 94, or a

sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 94, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 52.

5           As another example, the antibody [224G11] [ $\Delta$ 1-3] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 95, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 95, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80% identity after optimum alignment with the  
10           sequence SEQ ID No. 52.

          As another example, the antibody [224G11] [C7 $\Delta$ 6] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 96, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 96, and a complete light chain comprising the amino acid sequence SEQ ID No.  
15           52, or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 52.

          As another example, the antibody [224G11] [C6 $\Delta$ 9] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 97, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ  
20           ID No. 97, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 52.

          As another example, the antibody [224G11] [C2 $\Delta$ 5-7] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 98,  
25           or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 98, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 52.

          As another example, the antibody [224G11] [C5 $\Delta$ 2-6] of the invention  
30           comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 99, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 99, and a complete light chain comprising the amino acid sequence SEQ

ID No. 52, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 52.

As another example, the antibody [224G11] [C9Δ2-7] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 100, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 100, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 52.

As another example, the antibody [224G11] [Δ5-6-7-8] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 101, or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 101, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 52.

As another example, the antibody [224G11] [IgG1/IgG2] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 102, or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 102, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 52.

As another example, the antibody [224G11] [IgG2Hz1] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 36, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 36 and a complete light chain comprising the amino acid sequence SEQ ID No. 38, or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 38.

As another example, the antibody [224G11] [IgG2Hz2] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 36, or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 36 and a complete light chain comprising the amino acid sequence SEQ ID No. 39, or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 39.

As another example, the antibody [224G11] [IgG2Hz3] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 36, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 36 and a complete light chain comprising the amino acid sequence SEQ ID No. 40, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 40.

As another example, the antibody [224G11] [TH7Hz1] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 37, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 37 and a complete light chain comprising the amino acid sequence SEQ ID No. 38, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 38.

As another example, the antibody [224G11] [TH7Hz2] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 37, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 37 and a complete light chain comprising the amino acid sequence SEQ ID No. 39, or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 39.

As another example, the antibody [224G11] [TH7Hz3] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 37, or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 37 and a complete light chain comprising the amino acid sequence SEQ ID No. 40, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 40.

Other examples of antibodies, or derivatives thereof, according to the invention comprises complete heavy chains comprising an amino acid sequence selected in the group consisting of SEQ ID Nos. 88 to 102 (corresponding nucleotide sequences are SEQ ID Nos. 103 to 117).

By “functional fragment” of an antibody according to the invention, it is intended to indicate in particular an antibody fragment, such as Fv, scFv (sc for single chain), Fab, F(ab')<sub>2</sub>, Fab', scFv-Fc fragments or diabodies, or any fragment of which the half-life time would have been increased by chemical modification, such as the

addition of poly(alkylene) glycol such as poly(ethylene) glycol (“PEGylation”) (pegylated fragments called Fv-PEG, scFv-PEG, Fab-PEG, F(ab’)<sub>2</sub>-PEG or Fab’-PEG) (“PEG” for Poly(Ethylene) Glycol), or by incorporation in a liposome, said fragments having at least one of the characteristic CDRs of sequence SEQ ID Nos. 1 to 3 and 5 to 7 according to the invention, and, especially, in that it is capable of exerting in a general manner an even partial activity of the antibody from which it is descended, such as in particular the capacity to recognize and to bind to the c-Met, and, if necessary, to inhibit the activity of the c-Met.

Preferably, said functional fragments will be constituted or will comprise a partial sequence of the heavy or light variable chain of the antibody from which they are derived, said partial sequence being sufficient to retain the same specificity of binding as the antibody from which it is descended and a sufficient affinity, preferably at least equal to 1/100, in a more preferred manner to at least 1/10, of that of the antibody from which it is descended, with respect to the c-Met. Such a functional fragment will contain at the minimum 5 amino acids, preferably 6, 7, 8, 9, 10, 12, 15, 25, 50 and 100 consecutive amino acids of the sequence of the antibody from which it is descended.

Preferably, these functional fragments will be fragments of Fv, scFv, Fab, F(ab’)<sub>2</sub>, F(ab’), scFv-Fc type or diabodies, which generally have the same specificity of binding as the antibody from which they are descended. In a more preferred embodiment of the invention, these fragments are selected among divalent fragments such as F(ab’)<sub>2</sub> fragments. According to the present invention, antibody fragments of the invention can be obtained starting from antibodies such as described above by methods such as digestion by enzymes, such as pepsin or papain and/or by cleavage of the disulfide bridges by chemical reduction. In another manner, the antibody fragments comprised in the present invention can be obtained by techniques of genetic recombination likewise well known to the person skilled in the art or else by peptide synthesis by means of, for example, automatic peptide synthesizers such as those supplied by the company Applied Biosystems, etc.

By “divalent fragment”, it must be understood any antibody fragments comprising two arms and, more particularly, F(ab’)<sub>2</sub> fragments.

By “derivatives” of an antibody according to the invention, it is meant a binding protein comprising a protein scaffold and at least one of the CDRs selected from the



original antibody in order to maintain the binding capacity. Such compounds are well known by the man skilled in the art and will be described in more details in the following specification.

More particularly, the antibody, or one of its functional fragments or derivatives,  
5 according to the invention is characterized in that said derivative consists in a binding protein comprising a scaffold on which at least one CDR has been grafted for the conservation of the original antibody paratopic recognizing properties.

One or several sequences through the 6 CDR sequences described in the invention can be presented on a protein scaffold. In this case, the protein scaffold  
10 reproduces the protein backbone with appropriate folding of the grafted CDR(s), thus allowing it (or them) to maintain their antigen paratopic recognizing properties.

The man skilled in the art knows how to select the protein scaffold on which at least one CDR selected from the original antibody could be grafted. More particularly, it is known that, to be selected, such scaffold should display several features as follow  
15 (Skerra A., *J. Mol. Recogn.*, 13, 2000, 167-187):

- phylogenetically good conservation,
- robust architecture with a well known three-dimensional molecular organization (such as, for example, crystallography or NMR),
- small size,
- 20 - no or only low degree of post-translational modifications,
- easy to produce, express and purify.

Such protein scaffold can be, but without limitation, structure selected from the group consisting in fibronectin and preferentially the tenth fibronectin type III domain (FNfn10), lipocalin, anticalin (Skerra A., *J. Biotechnol.*, 2001, 74(4):257-75), the  
25 protein Z derivative from the domain B of staphylococcal protein A, thioredoxin A or any protein with repeated domain such as "ankyrin repeat" (Kohl et al., *PNAS*, 2003, vol. 100, No.4, 1700-1705), "armadillo repeat", "leucin-rich repeat" or "tetratricopeptide repeat".

It could also be mentioned scaffold derivative from toxins (such as, for example,  
30 scorpion, insect, plant or mollusc toxins) or protein inhibitors of neuronal nitric oxyde synthase (PIN).

As non limitative example of such hybrid constructions, it can be mentioned the

insertion of the CDR-H1 (heavy chain) of an anti-CD4 antibody, i.e. the 13B8.2 antibody, into one of the exposed loop of the PIN. The binding properties of the obtained binding protein remain similar to the original antibody (Bcs et al., BBRC 343, 2006, 334-344). It can also be mentioned the grafting of the CDR-H3 (heavy chain) of an anti-lysozyme VHH antibody on a loop of neocarzinostatine (Nicaise et al., 2004).

As above mentioned, such protein scaffold can comprise from 1 to 6 CDR(s) from the original antibody. In a preferred embodiment, but without any limitation, the man skilled in the art would select at least a CDR from the heavy chain, said heavy chain being known to be particularly implicated in the antibody specificity. The selection of the CDR(s) of interest will be evident for the man of the art with known method (BES et al., FEBS letters 508, 2001, 67-74).

As an evidence, these examples are not limitative and any other scaffold known or described must be included in the present specification.

According to a novel aspect, the present invention relates to an isolated nucleic acid, characterized in that it is chosen from the following nucleic acids:

- a) a nucleic acid, DNA or RNA, coding for an antibody, or one of its functional fragments or derivatives, according to the invention;
- b) a nucleic acid sequence comprising the sequences SEQ ID No. 11, SEQ ID No. 12, SEQ ID No. 13 and the sequences SEQ ID No. 15, SEQ ID No. 16 and SEQ ID No. 17;
- c) a nucleic acid sequence comprising the sequences SEQ ID No. 14 and SEQ ID No. 18, 19 or 20;
- d) the corresponding RNA nucleic acids of the nucleic acids as defined in b) or c);
- e) the complementary nucleic acids of the nucleic acids as defined in a), b) and c); and
- f) a nucleic acid of at least 18 nucleotides capable of hybridizing under conditions of high stringency with at least one of the CDRs of sequence SEQ ID Nos. 11 to 13 and 15 to 17.

According to still another aspect, the present invention relates to an isolated nucleic acid, characterized in that it is chosen from the following nucleic acids:

- a nucleic acid, DNA or RNA, coding for an antibody, or one of its functional

fragments or derivatives, according to the present invention and wherein the nucleic sequence coding for the hinge region of said antibody comprises or has a sequence selected from the group consisting of the sequences SEQ ID Nos. 29 to 35 and SEQ ID Nos. 73 to 87.

5 By nucleic acid, nucleic or nucleic acid sequence, polynucleotide, oligonucleotide, polynucleotide sequence, nucleotide sequence, terms which will be employed indifferently in the present invention, it is intended to indicate a precise linkage of nucleotides, which are modified or unmodified, allowing a fragment or a region of a nucleic acid to be defined, containing or not containing unnatural  
10 nucleotides, and being able to correspond just as well to a double-stranded DNA, a single-stranded DNA as to the transcription products of said DNAs.

It must also be understood here that the present invention does not concern the nucleotide sequences in their natural chromosomal environment, that is to say in the natural state. It concerns sequences which have been isolated and/or purified, that is to  
15 say that they have been selected directly or indirectly, for example by copy, their environment having been at least partially modified. It is thus likewise intended to indicate here the isolated nucleic acids obtained by genetic recombination by means, for example, of host cells or obtained by chemical synthesis.

A hybridization under conditions of high stringency signifies that the  
20 temperature conditions and ionic strength conditions are chosen in such a way that they allow the maintenance of the hybridization between two fragments of complementary DNA. By way of illustration, conditions of high stringency of the hybridization step for the purposes of defining the polynucleotide fragments described above are advantageously the following.

25 The DNA-DNA or DNA-RNA hybridization is carried out in two steps: (1) prehybridization at 42°C for 3 hours in phosphate buffer (20 mM, pH 7.5) containing 5 x SSC (1 x SSC corresponds to a 0.15 M NaCl + 0.015 M sodium citrate solution), 50 % of formamide, 7 % of sodium dodecyl sulfate (SDS), 10 x Denhardt's, 5 % of dextran sulfate and 1 % of salmon sperm DNA; (2) actual hybridization for 20 hours at  
30 a temperature dependent on the size of the probe (i.e.: 42°C, for a probe size > 100 nucleotides) followed by 2 washes of 20 minutes at 20°C in 2 x SSC + 2% of SDS, 1 wash of 20 minutes at 20°C in 0.1 x SSC + 0.1 % of SDS. The last wash is carried out

in 0.1 x SSC + 0.1 % of SDS for 30 minutes at 60°C for a probe size > 100 nucleotides. The hybridization conditions of high stringency described above for a polynucleotide of defined size can be adapted by the person skilled in the art for oligonucleotides of greater or smaller size, according to the teaching of Sambrook et al. (1989, Molecular cloning: a laboratory manual. 2nd Ed. Cold Spring Harbor).

The invention likewise relates to a vector comprising a nucleic acid according to the present invention.

The invention aims especially at cloning and/or expression vectors which contain a nucleotide sequence according to the invention.

The vectors according to the invention preferably contain elements which allow the expression and/or the secretion of the translated nucleotide sequences in a determined host cell. The vector must therefore contain a promoter, signals of initiation and termination of translation, as well as appropriate regions of regulation of transcription. It must be able to be maintained in a stable manner in the host cell and can optionally have particular signals which specify the secretion of the translated protein. These different elements are chosen and optimized by the person skilled in the art as a function of the host cell used. To this effect, the nucleotide sequences according to the invention can be inserted into autonomous replication vectors in the chosen host, or be integrative vectors of the chosen host.

Such vectors are prepared by methods currently used by the person skilled in the art, and the resulting clones can be introduced into an appropriate host by standard methods, such as lipofection, electroporation, thermal shock, or chemical methods.

The vectors according to the invention are, for example, vectors of plasmidic or viral origin. They are useful for transforming host cells in order to clone or to express the nucleotide sequences according to the invention.

The invention likewise comprises the host cells transformed by or comprising a vector according to the invention.

The host cell can be chosen from prokaryotic or eukaryotic systems, for example bacterial cells but likewise yeast cells or animal cells, in particular mammalian cells. It is likewise possible to use insect cells or plant cells.

The invention likewise concerns animals, except man, which comprise at least one cell transformed according to the invention.

According to another aspect, a subject of the invention is a process for production of an antibody, or one of its functional fragments according to the invention, characterized in that it comprises the following stages:

5 a) culture in a medium and appropriate culture conditions of a host cell according to the invention; and

b) the recovery of said antibodies, or one of their functional fragments, thus produced starting from the culture medium or said cultured cells.

The cells transformed according to the invention can be used in processes for preparation of recombinant polypeptides according to the invention. The processes for  
10 preparation of a polypeptide according to the invention in recombinant form, characterized in that they employ a vector and/or a cell transformed by a vector according to the invention, are themselves comprised in the present invention. Preferably, a cell transformed by a vector according to the invention is cultured under conditions which allow the expression of said polypeptide and said recombinant peptide  
15 is recovered.

As has been said, the host cell can be chosen from prokaryotic or eukaryotic systems. In particular, it is possible to identify nucleotide sequences according to the invention, facilitating secretion in such a prokaryotic or eukaryotic system. A vector according to the invention carrying such a sequence can therefore advantageously be  
20 used for the production of recombinant proteins, intended to be secreted. In effect, the purification of these recombinant proteins of interest will be facilitated by the fact that they are present in the supernatant of the cell culture rather than in the interior of the host cells.

It is likewise possible to prepare the polypeptides according to the invention by  
25 chemical synthesis. Such a preparation process is likewise a subject of the invention. The person skilled in the art knows the processes of chemical synthesis, for example the techniques employing solid phases [Steward et al., 1984, Solid phase peptide synthesis, Pierce Chem. Company, Rockford, 111, 2nd ed., (1984)] or techniques using partial solid phases, by condensation of fragments or by a classical synthesis in solution. The  
30 polypeptides obtained by chemical synthesis and being able to contain corresponding unnatural amino acids are likewise comprised in the invention.

The antibodies, or one of their functional fragments or derivatives, capable of being obtained by a process according to the invention are likewise comprised in the present invention.

The invention also concerns the antibody of the invention as a medicament.

5 The invention likewise concerns a pharmaceutical composition comprising by way of active principle a compound consisting of an antibody, or one of its functional fragments according to the invention, preferably mixed with an excipient and/or a pharmaceutically acceptable vehicle.

10 Another complementary embodiment of the invention consists in a composition such as described above which comprises, moreover, as a combination product for simultaneous, separate or sequential use, an anti-tumoral antibody.

15 Most preferably, said second anti-tumoral antibody could be chosen through anti-IGF-IR, anti-EGFR, anti-HER2/neu, anti-VEGFR, anti-VEGF, etc., antibodies or any other anti-tumoral antibodies known by the man skilled in the art. It is evident that the use, as second antibody, of functional fragments or derivatives of above mentioned antibodies is part of the invention.

As a most preferred antibody, anti-EGFR antibodies are selected such as for example the antibody C225 (Erbix).

20 "Simultaneous use" is understood as meaning the administration of the two compounds of the composition according to the invention in a single and identical pharmaceutical form.

"Separate use" is understood as meaning the administration, at the same time, of the two compounds of the composition according to the invention in distinct pharmaceutical forms.

25 "Sequential use" is understood as meaning the successive administration of the two compounds of the composition according to the invention, each in a distinct pharmaceutical form.

30 In a general fashion, the composition according to the invention considerably increases the efficacy of the treatment of cancer. In other words, the therapeutic effect of the anti-c-Met antibodies according to the invention is potentiated in an unexpected manner by the administration of a cytotoxic agent. Another major subsequent advantage produced by a composition according to the invention concerns the possibility of using

lower efficacious doses of active principle, which allows the risks of appearance of secondary effects to be avoided or to be reduced, in particular the effects of the cytotoxic agent.

In addition, this composition according to the invention would allow the  
5 expected therapeutic effect to be attained more rapidly.

The composition of the invention can also be characterized in that it comprises, moreover, as a combination product for simultaneous, separate or sequential use, a cytotoxic/cytostatic agent.

By "anti-cancer therapeutic agents" or "cytotoxic/cytostatic agents", it is  
10 intended a substance which, when administered to a subject, treats or prevents the development of cancer in the subject's body. As non limitative example of such agents, it can be mentioned alkylating agents, anti-metabolites, anti-tumor antibiotics, mitotic inhibitors, chromatin function inhibitors, anti-angiogenesis agents, anti-estrogens, anti-androgens or immunomodulators.

Such agents are, for example, cited in the 2001 edition of VIDAL, on the page  
15 devoted to the compounds attached to the cancerology and hematology column "Cytotoxics", these cytotoxic compounds cited with reference to this document are cited here as preferred cytotoxic agents.

More particularly, the following agents are preferred according to the invention.

20 "Alkylating agent" refers to any substance which can cross-link or alkylate any molecule, preferably nucleic acid (e.g., DNA), within a cell. Examples of alkylating agents include nitrogen mustard such as mechlorethamine, chlorambucol, melphalen, chlorydrate, pipobromen, prednimustin, disodic-phosphate or estramustine; oxazophorins such as cyclophosphamide, altretamine, trofosfamide, sulfosfosfamide or  
25 ifosfamide; aziridines or imine-ethylenes such as thiotepa, triethylenamine or altetramine; nitrosoarea such as carmustine, streptozocin, fotemustin or lomustine; alkyle-sulfonates such as busulfan, treosulfan or improsulfan; triazenes such as dacarbazine; or platinum complexes such as cis-platinum, oxaliplatin and carboplatin.

"Anti-metabolites" refer to substances that block cell growth and/or metabolism  
30 by interfering with certain activities, usually DNA synthesis. Examples of anti-metabolites include methotrexate, 5-fluoruracil, floxuridine, 5-fluorodeoxyuridine, capecitabine, cytarabine, fludarabine, cytosine arabinoside, 6-mercaptopurine (6-MP),

6-thioguanine (6-TG), chlorodesoxyadenosine, 5-azacytidine, gemcitabine, cladribine, deoxycoformycin and pentostatin.

“Anti-tumor antibiotics” refer to compounds which may prevent or inhibit DNA, RNA and/or protein synthesis. Examples of anti-tumor antibiotics include doxorubicin, daunorubicin, idarubicin, valrubicin, mitoxantrone, dactinomycin, mithramycin, plicamycin, mitomycin C, bleomycin, and procarbazine.

“Mitotic inhibitors” prevent normal progression of the cell cycle and mitosis. In general, microtubule inhibitors or taxoides such as paclitaxel and docetaxel are capable of inhibiting mitosis. Vinca alkaloid such as vinblastine, vincristine, vindesine and vinorelbine are also capable of inhibiting mitosis.

“Chromatin function inhibitors” or “topoisomerase inhibitors” refer to substances which inhibit the normal function of chromatin modeling proteins such as topoisomerase I or topoisomerase II. Examples of chromatin function inhibitors include, for topoisomerase I, camptothecine and its derivatives such as topotecan or irinotecan, and, for topoisomerase II, etoposide, etoposide phosphate and teniposide.

“Anti-angiogenesis agent” refers to any drug, compound, substance or agent which inhibits growth of blood vessels. Exemplary anti-angiogenesis agents include, but are by no means limited to, razoxin, marimastat, batimastat, prinomastat, tanomastat, ilomastat, CGS-27023A, halofuginon, COL-3, neovastat, BMS-275291, thalidomide, CDC 501, DMXAA, L-651582, squalamine, endostatin, SU5416, SU6668, interferon-alpha, EMD121974, interleukin-12, IM862, angiostatin and vitaxin.

“Anti-estrogen” or “anti-estrogenic agent” refer to any substance which reduces, antagonizes or inhibits the action of estrogen. Examples of anti-estrogen agents are tamoxifen, toremifene, raloxifene, droloxifene, iodoxyfene, anastrozole, letrozole, and exemestane.

“Anti-androgens” or “anti-androgen agents” refer to any substance which reduces, antagonizes or inhibits the action of an androgen. Examples of anti-androgens are flutamide, nilutamide, bicalutamide, sprironolactone, cyproterone acetate, finasteride and cimitidine.

“Immunomodulators” are substances which stimulate the immune system.

Examples of immunomodulators include interferon, interleukin such as aldesleukine, OCT-43, denileukin diflitox and interleukin-2, tumoral necrose factors such



as tasonermine or others immunomodulators such as lentinan, sizofiran, roquinimex, pidotimod, pegademase, thymopentine, poly I:C or levamisole in conjunction with 5-fluorouracil.

For more detail, the man skill in the art could refer to the manual edited by the  
5 “Association Française des Enseignants de Chimie Thérapeutique” and entitled “Traité de chimie thérapeutique”, vol. 6, Médicaments antitumoraux et perspectives dans le traitement des cancers, edition TEC & DOC, 2003.

Can also be mentioned as chemical agents or cytotoxic agents, all kinase inhibitors such as, for example, gefitinib or erlotinib.

10 In a particularly preferred embodiment, said composition as a combination product according to the invention is characterized in that said cytotoxic agent is coupled chemically to said antibody for simultaneous use.

In order to facilitate the coupling between said cytotoxic agent and said antibody according to the invention, it is especially possible to introduce spacer molecules  
15 between the two compounds to be coupled, such as poly(alkylene) glycols like polyethylene glycol, or else amino acids, or, in another embodiment, to use active derivatives of said cytotoxic agents into which would have been introduced functions capable of reacting with said antibody according to the invention. These coupling techniques are well known to the person skilled in the art and will not be expanded upon  
20 in the present description.

The invention relates, in another aspect, to a composition characterized in that one, at least, of said antibodies, or one of their functional fragments or derivatives, is conjugated with a cell toxin and/or a radioelement.

Preferably, said toxin or said radioelement is capable of inhibiting at least one  
25 cell activity of cells expressing the c-Met, in a more preferred manner capable of preventing the growth or the proliferation of said cell, especially of totally inactivating said cell.

Preferably also, said toxin is an enterobacterial toxin, especially Pseudomonas exotoxin A.

30 The radioelements (or radioisotopes) preferably conjugated to the antibodies employed for the therapy are radioisotopes which emit gamma rays and preferably

iodine<sup>131</sup>, yttrium<sup>90</sup>, gold<sup>199</sup>, palladium<sup>100</sup>, copper<sup>67</sup>, bismuth<sup>217</sup> and antimony<sup>211</sup>. The radioisotopes which emit beta and alpha rays can likewise be used for the therapy.

By toxin or radioelement conjugated to at least one antibody, or one of its functional fragments, according to the invention, it is intended to indicate any means  
5 allowing said toxin or said radioelement to bind to said at least one antibody, especially by covalent coupling between the two compounds, with or without introduction of a linking molecule.

Among the agents allowing binding in a chemical (covalent), electrostatic or noncovalent manner of all or part of the components of the conjugate, mention may  
10 particularly be made of benzoquinone, carbodiimide and more particularly EDC (1-ethyl-3-[3-dimethyl-aminopropyl]-carbodiimide hydrochloride), dimaleimide, dithiobis-nitrobenzoic acid (DTNB), N-succinimidyl S-acetyl thio-acetate (SATA), the bridging agents having one or more phenylazide groups reacting with the ultraviolets (U.V.) and preferably N-[-4-(azidosalicylamino)butyl]-3'-(2'-pyridyldithio)-propionamide (APDP),  
15 N-succinimid-yl 3-(2-pyridyldithio)propionate (SPDP), 6-hydrazino-nicotinamide (HYNIC).

Another form of coupling, especially for the radioelements, can consist in the use of a bifunctional ion chelator.

Among these chelates, it is possible to mention the chelates derived from EDTA  
20 (ethylenediaminetetraacetic acid) or from DTPA (diethylenetriaminepentaacetic acid) which have been developed for binding metals, especially radioactive metals, and immunoglobulins. Thus, DTPA and its derivatives can be substituted by different groups on the carbon chain in order to increase the stability and the rigidity of the ligand-metal complex (Krejcarek et al. (1977); Brechbiel et al. (1991); Gansow (1991);  
25 US patent 4,831,175).

For example diethylenetriaminepentaacetic acid (DTPA) and its derivatives, which have been widely used in medicine and in biology for a long time either in their free form, or in the form of a complex with a metallic ion, have the remarkable characteristic of forming stable chelates with metallic ions and of being coupled with  
30 proteins of therapeutic or diagnostic interest such as antibodies for the development of radioimmunoconjugates in cancer therapy (Meases et al., 1984; Gansow et al., 1990).

Likewise preferably, said at least one antibody forming said conjugate according to the invention is chosen from its functional fragments, especially the fragments amputated of their Fc component such as the scFv fragments.

As already mentioned, in a preferred embodiment of the invention, said  
5 cytotoxic/cytostatic agent or said toxin and/or a radioelement is coupled chemically to at least one of the elements of said composition for simultaneous use.

The present invention comprises the described composition as a medicament.

The present invention moreover comprises the use of the composition according to the invention for the preparation of a medicament.

10 In another aspect, the invention deals with the use of an antibody, or one of its functional fragments or derivatives, and/or of a composition as above described for the preparation of a medicament intended to inhibit the growth and/or the proliferation of tumor cells.

Another aspect of the invention consists in the use of an antibody, or one of its  
15 functional fragments or derivatives and/or of a composition, as described above or the use above mentioned, for the preparation of a medicament intended for the prevention or for the treatment of cancer.

Is also comprised in the present invention a method intended to inhibit the growth and/or the proliferation of tumor cells in a patient comprising the administration  
20 to a patient in need thereof of an antibody, or one of its functional fragments or derivatives according to the invention, an antibody produced by an hybridoma according to the invention or a composition according to the invention.

The present invention further comprises a method for the prevention or the treatment of cancer in a patient in need thereof, comprising the administration to the  
25 patient of an antibody, or one of its functional fragments or derivatives according to the invention, an antibody produced by an hybridoma according to the invention or a composition according to the invention.

In a particular preferred aspect, said cancer is a cancer chosen from prostate cancer, osteosarcomas, lung cancer, breast cancer, endometrial cancer, glioblastoma or  
30 colon cancer.

As explained before, an advantage of the invention is to allow the treatment of HGF dependent and independent Met-activation related cancers.

The invention, in yet another aspect, encompasses a method of *in vitro* diagnosis of illnesses induced by an overexpression or an underexpression of the c-Met receptor starting from a biological sample in which the abnormal presence of c-Met receptor is suspected, said method being characterized in that it comprises a step wherein said biological sample is contacted with an antibody of the invention, it being possible for said antibody to be, if necessary, labeled.

Preferably, said illnesses connected with an abnormal presence of c-Met receptor in said diagnosis method will be cancers.

Said antibody, or one of its functional fragments, can be present in the form of an immunoconjugate or of a labelled antibody so as to obtain a detectable and/or quantifiable signal.

The antibodies labelled according to the invention or their functional fragments include, for example, antibodies called immunoconjugates which can be conjugated, for example, with enzymes such as peroxidase, alkaline phosphatase, beta-D-galactosidase, glucose oxydase, glucose amylase, carbonic anhydrase, acetylcholinesterase, lysozyme, malate dehydrogenase or glucose 6-phosphate dehydrogenase or by a molecule such as biotin, digoxigenin or 5-bromodeoxyuridine. Fluorescent labels can be likewise conjugated to the antibodies or to their functional fragments according to the invention and especially include fluorescein and its derivatives, fluorochrome, rhodamine and its derivatives, GFP (GFP for "Green Fluorescent Protein"), dansyl, umbelliferone etc. In such conjugates, the antibodies of the invention or their functional fragments can be prepared by methods known to the person skilled in the art. They can be coupled to the enzymes or to the fluorescent labels directly or by the intermediary of a spacer group or of a linking group such as a polyaldehyde, like glutaraldehyde, ethylenediaminetetraacetic acid (EDTA), diethylene-triaminepentaacetic acid (DPTA), or in the presence of coupling agents such as those mentioned above for the therapeutic conjugates. The conjugates containing labels of fluorescein type can be prepared by reaction with an isothiocyanate.

Other conjugates can likewise include chemoluminescent labels such as luminol and the dioxetanes, bio-luminescent labels such as luciferase and luciferin, or else radioactive labels such as iodine<sup>123</sup>, iodine<sup>125</sup>, iodine<sup>126</sup>, iodine<sup>133</sup>, bromine<sup>77</sup>, technetium<sup>99m</sup>, indium<sup>111</sup>, indium<sup>113m</sup>, gallium<sup>67</sup>, gallium<sup>68</sup>, ruthenium<sup>95</sup>, ruthenium<sup>97</sup>,

ruthenium<sup>103</sup>, ruthenium<sup>105</sup>, mercury<sup>107</sup>, mercury<sup>203</sup>, rhenium<sup>99m</sup>, rhenium<sup>101</sup>, rhenium<sup>105</sup>, scandium<sup>47</sup>, tellurium<sup>121m</sup>, tellurium<sup>122m</sup>, tellurium<sup>125m</sup>, thulium<sup>165</sup>, thulium<sup>167</sup>, thulium<sup>168</sup>, fluorine<sup>18</sup>, yttrium<sup>199</sup>, iodine<sup>131</sup>. The methods known to the person skilled in the art existing for coupling the therapeutic radioisotopes to the antibodies either  
5 directly or via a chelating agent such as EDTA, DTPA mentioned above can be used for the radioelements which can be used in diagnosis. It is likewise possible to mention labelling with Na[I<sup>125</sup>] by the chloramine T method [Hunter W.M. and Greenwood F.C. (1962) Nature 194:495] or else with technetium<sup>99m</sup> by the technique of Crockford *et al.* (US patent 4,424,200) or attached via DTPA as described by Hnatowich (US patent  
10 4,479,930).

Thus, the antibody, or a functional fragment or derivative thereof, according to the invention can be employed in a process for the detection and/or the quantification of an overexpression or of an underexpression, preferably an overexpression, of the c-Met receptor in a biological sample, characterized in that it comprises the following steps:

- 15 a) the contacting of the biological sample with an antibody, or a functional fragment or derivative thereof, according to the invention; and  
b) the demonstration of the c-Met/antibody complex possibly formed.

In a particular embodiment, the antibody, or a functional fragment or derivative thereof, according to the invention, can be employed in a process for the detection  
20 and/or the quantification of the c-Met receptor in a biological sample, for the monitoring of the efficacy of a prophylactic and/or therapeutic treatment of c-Met-dependent cancer.

More generally, the antibody or a functional fragment or derivative thereof, according to the invention can be advantageously employed in any situation where the  
25 expression of the c-Met- receptor must be observed in a qualitative and/or quantitative manner.

Preferably, the biological sample is formed by a biological fluid, such as serum, whole blood, cells, a tissue sample or biopsies of human origin.

Any procedure or conventional test can be employed in order to carry out such a  
30 detection and/or dosage. Said test can be a competition or sandwich test, or any test known to the person skilled in the art dependent on the formation of an immune complex of antibody-antigen type. Following the applications according to the

invention, the antibody or a functional fragment or derivative thereof can be immobilized or labelled. This immobilization can be carried out on numerous supports known to the person skilled in the art. These supports can especially include glass, polystyrene, poly-propylene, polyethylene, dextran, nylon, or natural or modified cells.

5 These supports can be either soluble or insoluble.

By way of example, a preferred method brings into play immunoenzymatic processes according to the ELISA technique, by immunofluorescence, or radio-immunoassay (RIA) technique or equivalent.

Thus, the present invention likewise comprises the kits or sets necessary for  
10 carrying out a method of diagnosis of illnesses induced by an overexpression or an underexpression of the c-Met receptor or for carrying out a process for the detection and/or the quantification of an overexpression or of an underexpression of the c-Met receptor in a biological sample, preferably an overexpression of said receptor, characterized in that said kit or set comprises the following elements:

15 a) an antibody, or a functional fragment or derivative thereof, according to the invention;

b) optionally, the reagents for the formation of the medium favorable to the immunological reaction;

20 c) optionally, the reagents allowing the demonstration of c-Met/antibody complexes produced by the immunological reaction.

A subject of the invention is likewise the use of an antibody or a composition according to the invention for the preparation of a medicament intended for the specific targeting of a biologically active compound to cells expressing or overexpressing the c-Met receptor.

25 It is intended here by biologically active compound to indicate any compound capable of modulating, especially of inhibiting, cell activity, in particular their growth, their proliferation, transcription or gene translation.

A subject of the invention is also an *in vivo* diagnostic reagent comprising an antibody according to the invention, or a functional fragment or derivative thereof,  
30 preferably labelled, especially radiolabelled, and its use in medical imaging, in particular for the detection of cancer connected with the expression or the overexpression by a cell of the c-Met receptor.

The invention likewise relates to a composition as a combination product or to an anti-c-Met/toxin conjugate or radioelement, according to the invention, as a medicament.

Preferably, said composition as a combination product or said conjugate  
5 according to the invention will be mixed with an excipient and/or a pharmaceutically acceptable vehicle.

In the present description, pharmaceutically acceptable vehicle is intended to indicate a compound or a combination of compounds entering into a pharmaceutical composition not provoking secondary reactions and which allows, for example,  
10 facilitation of the administration of the active compound(s), an increase in its lifespan and/or in its efficacy in the body, an increase in its solubility in solution or else an improvement in its conservation. These pharmaceutically acceptable vehicles are well known and will be adapted by the person skilled in the art as a function of the nature and of the mode of administration of the active compound(s) chosen.

Preferably, these compounds will be administered by the systemic route, in  
15 particular by the intravenous route, by the intramuscular, intradermal, intraperitoneal or subcutaneous route, or by the oral route. In a more preferred manner, the composition comprising the antibodies according to the invention will be administered several times, in a sequential manner.

Their modes of administration, dosages and optimum pharmaceutical forms can  
20 be determined according to the criteria generally taken into account in the establishment of a treatment adapted to a patient such as, for example, the age or the body weight of the patient, the seriousness of his/her general condition, the tolerance to the treatment and the secondary effects noted.

Other characteristics and advantages of the invention appear in the continuation  
25 of the description with the examples and the figures wherein:

Figure 1: Effect of irrelevant IgG1 Mabs from mouse and human origin and PBS on c-Met receptor phosphorylation on A549 cells.

Figures 2A and 2B: Effect of murine and humanized 224G11 Mabs produced as  
30 a human IgG1/kappa isotype on c-Met receptor phosphorylation on A549 cells.

Figure 2A: agonist effect calculated as percentage versus maximal stimulation of c-Met phosphorylation by HGF [100 ng/ml].

Figure 2B: antagonist effect calculated as percentage of inhibition of the maximal stimulation of c-Met phosphorylation by HGF [100 ng/ml].

Figures 3A and 3B: Comparison between murine 224G11 Mab and chimeric 224G11 Mabs containing various engineered hinge regions, on c-Met receptor phosphorylation on A549 cells.

Figure 3A: agonist effect calculated as percentage versus maximal stimulation of c-Met phosphorylation by HGF [100 ng/ml].

Figure 3B: antagonist effect calculated as percentage of inhibition of the maximal stimulation of c-Met phosphorylation by HGF [100 ng/ml].

Figures 4A and 4B: Comparison between murine 224G11 Mab and chimeric and humanized 224G11 Mabs produced as a human IgG2/kappa isotype, on c-Met receptor phosphorylation on A549 cells.

Figure 4A: agonist effect calculated as percentage versus maximal stimulation of c-Met phosphorylation by HGF [100 ng/ml].

Figure 4B: antagonist effect calculated as percentage of inhibition of the maximal stimulation of c-Met phosphorylation by HGF [100 ng/ml].



Figures 5A and 5B: Comparison between murine 224G11 Mab and chimeric and humanized 224G11 Mabs produced as an engineered hinge mutant TH7IgG1/kappa, on c-Met receptor phosphorylation on A549 cells.

Figure 5A: agonist effect calculated as percentage versus maximal stimulation of c-Met phosphorylation by HGF [100 ng/ml].

Figure 5B: antagonist effect calculated as percentage of inhibition of the maximal stimulation of c-Met phosphorylation by HGF [100 ng/ml].

Figures 6A and 6B, Figures 7A and 7B, Figures 8A and 8B, Figures 9A and 9B, Figures 10A and 10B: BRET models with Figures A: c-Met dimerization model; and Figures B: c-Met activation model.

Figure 11: c-Met recognition by chimeric and humanized 224G11 forms.

Figure 12: Effect of murine and chimeric antibodies on HGF-induced proliferation of NCI-H441 cells *in vitro*. NCI-H441 cells were plated in serum-free medium. 24 hours after plating m224G11 and [224G11]chim were added either in absence or in presence of HGF. Black arrows indicate the wells plated with cells alone either in absence  or in presence  of HGF. A murine IgG1 (mIgG1) was introduced



as an isotype control.

Figure 13: *In vivo* comparison of murine and IgG1 chimeric 224G11 Mabs on the NCI-H441 xenograft model.





Figures 14A and 14B: Effect of the murine 224G11 Mab and of various  
5 chimeric and humanized versions of this antibody on HGF-induced proliferation of  
NCI-H441 cells in vitro. NCI-H441 cells were plated in serum-free medium. Twenty  
four hours after plating antibody to be tested were added either in absence or in  
presence of HGF. In panel (Figure 14A), the murine m224G11, chimeric IgG1  
[224G11]chim, humanized IgG1 [224G11] [Hz1], [224G11] [Hz2], [224G11] [Hz3]  
10 versions were shown. In panel (Figure 14B), the murine m224G11 and various chimeric  
IgG1 forms ([224G11] chim, [224G11] [MH chim], [224G11] [MUP9H chim],  
[224G11] [MMCH chim], [224G11] [TH7 chim]) were presented. Black arrows  
indicate the wells plated with cells alone either in absence  or in presence  of HGF.  
A murine IgG1 was introduced as a negative control for agonist activity. The m5D5 was  
15 used as a dose-dependent full agonist control.

Figure 15: Effect of the murine 224G11 Mab and of various chimeric and  
humanized versions of this antibody on HGF-induced proliferation of NCI-H441 cells  
in vitro. NCI-H441 cells were plated in serum-free medium. Twenty four hours after  
plating antibody to be tested were added either in absence or in presence of HGF. The  
20 murine m224G11, [224G11] chim, [224G11] [TH7 chim]) IgG1 chimeric forms and  
[224G11] [TH7 Hz1], [224G11] [TH7 Hz3],) were presented. Black arrows indicate the  
wells plated with cells alone either in absence  or in presence  of HGF. A murine  
IgG1 was introduced as a negative control for agonist activity. The m5D5 was used as a  
dose-dependent full agonist control.

25 Figure 16: *In vivo* comparison of murine, chimeric and humanized 224G11  
Mabs on the NCI-H441 xenograft model.

Figure 17A: agonist effect calculated as percentage versus maximal stimulation  
of c-Met phosphorylation by HGF [100 ng/ml].

Figure 17B: antagonist effect calculated as percentage of inhibition of the  
30 maximal stimulation of c-Met phosphorylation by HGF [100 ng/ml].

Figure 18: BRET models with c-Met activation model.

Figure 19: Effect of m224G11 and h224G11 on c-Met degradation on A549

cells. A) Mean of 4 independent experiments +/- s.e.m. B) Western blot image representative of the 4 independent experiments performed.

Figure 20: Effect of m224G11 and h224G11 on c-Met degradation on NCI-H441 cells. A) Mean of 4 independent experiments +/- s.e.m. B) Western blot image representative of the 4 independent experiments performed.

Figure 21: Set up of an ELISA to evaluate c-Met shedding.

Figure 22: *In vitro* evaluation of c-Met shedding on NCI-H441 cells treated for 5 days with m224G11. mIgG1 is an irrelevant antibody used as an isotype control.

Figure 23: *In vitro* evaluation of c-Met shedding on amplified Hs746T, MKN45 and EBC-1 cell lines treated for 5 days with m224G11. mIgG1 is an irrelevant antibody used as an isotype control. PMA is a shedding inducer used as a positive control.

Figure 24: *In vitro* evaluation of c-Met shedding on NCI-H441 and amplified Hs746T, MKN45 and EBC-1 cell lines treated for 5 days with m224G11. mIgG1 is an irrelevant antibody used as an isotype control. PMA is a shedding inducer used as a positive control.

Figure 25: Study of intrinsic phosphorylation of h224G11 on Hs746T cell line.

Figure 26: Study of intrinsic phosphorylation of h224G11 on NCI-H441 cell line. A) phospho-ELISA and B) Western analysis.

Figure 27: Study of intrinsic phosphorylation of h224G11 on Hs578T cell line. A) phospho-ELISA and B) Western analysis.

Figure 28: Study of intrinsic phosphorylation of h224G11 on NCI-H125 cell line. A) phospho-ELISA and B) Western analysis.

Figure 29: Study of intrinsic phosphorylation of h224G11 on T98G cell line. A) phospho-ELISA and B) Western analysis.

Figure 30: Study of intrinsic phosphorylation of h224G11 on MDA-MB-231 cell line. A) phospho-ELISA and B) Western analysis.

Figure 31: Study of intrinsic phosphorylation of h224G11 on PC3 cell line. A) phospho-ELISA and B) Western analysis.

Figure 32: Study of intrinsic phosphorylation of h224G11 on HUVEC cells.

Figure 33: *In vivo* comparison of the wild type murine 224G11 antibody with a chimeric hinge-engineered 224G11[C2D5-7] Mabs on the NCI-H441 xenograft model.

Figure 34: ADCC induction by h224G11 on both Hs746T and NCI-H441 cells.

<sup>51</sup>Cr-labeled Hs746T (A) or NCI-H441 (B) cells loaded (bold squares) or not (empty squares) with h224G11 were mixed with different ratio of human NK cells and incubated for 4 hr. Cells were harvested and cpm of <sup>51</sup>Cr released by lysis was counted. The results are plotted as percentage of lysis against the effector/target cell ratio. NL for non loaded cells.

Figure 35: h224G11 staining in tumor xenograft which expressed various level of c-Met (A: Hs746T amplified cell line for c-Met, B: NCI-H441 high level of c-Met expression and C: MCF-7 low level of c-Met).

### 10           **Example 1: Generation of antibodies against c-Met**

To generate anti-c-Met antibodies 8 weeks old BALB/c mice were immunized either 3 to 5 times subcutaneously with a CHO transfected cell line that express c-Met on its plasma membrane ( $20 \times 10^6$  cells/dose/mouse) or 2 to 3 times with a c-Met extracellular domain fusion protein (10-15  $\mu\text{g}/\text{dose}/\text{mouse}$ ) (R&D Systems, Catalog # 358MT) or fragments of this recombinant protein mixed with complete Freund adjuvant for the first immunization and incomplete Freund adjuvant for the following ones. Mixed protocols in which mice received both CHO-cMet cells and recombinant proteins were also performed. Three days before cell fusion, mice were boosted i.p. or i.v. with the recombinant protein or fragments. Then spleens of mice were collected and fused to SP2/0-Ag14 myeloma cells (ATCC) and subjected to HAT selection. Four fusions were performed. In general, for the preparation of monoclonal antibodies or their functional fragments, especially of murine origin, it is possible to refer to techniques which are described in particular in the manual "Antibodies" (Harlow and Lane, Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor NY, pp. 726, 1988) or to the technique of preparation of hybridomas described by Kohler and Milstein (Nature, 256:495-497, 1975).

Obtained hybridomas were initially screened by ELISA on the c-Met recombinant protein and then by FACS analysis on A549 NSCLC, BxPC3 pancreatic, and U87-MG glioblastoma cell lines to be sure that the produced antibodies will be able to also recognize the native receptor on tumor cells. Positive reactors on these 2 tests were amplified, cloned and a set of hybridomas was recovered, purified and screened for its ability to inhibit *in vitro* cell proliferation in the BxPC3 model.

For that purpose 50 000 BxPC3 cells were plated in 96 well plates in RPMI medium, 2 mM L. Glutamine, without SVF. 24 hours after plating, antibodies to be tested were added at a final concentration ranging from 0.0097 to 40 µg/ml 60 min before addition of 100 ng/ml of hHGF. After 3 days, cells were pulsed with 0.5 µCi of [3H]thymidine for 16 hours. The magnitude of [3H]thymidine incorporated into trichloroacetic acid-insoluble DNA was quantified by liquid scintillation counting. Results were expressed as raw data to really evaluate the intrinsic agonistic effect of each Mab.

Then antibodies inhibiting at least 50% cell proliferation were evaluated for their activity on c-Met dimerization and activation BRET analysis on transfected cells. c-Met receptor activity was quantified by measuring the Gab1 signalling molecule recruitment on activated c-Met. For that purpose, CHO stable cell lines expressing C-Met-Rluc or C-Met-Rluc and C-Met-K1100A-YFP for c-Met dimerization or C-Met-Rluc and a mutated form of Gab1 [Maroun et al., Mol. Cell. Biol., 1999, 19:1784-1799] fused to YFP for c-Met activation were generated. Cells were distributed in white 96 well microplates in DMEM-F12/FBS 5 % culture medium one or two days before BRET experiments. Cells were first cultured at 37°C with CO<sub>2</sub> 5 % in order to allow cell attachment to the plate. Cells were then starved with 200 µl DMEM/well overnight. Immediately prior to the experiment, DMEM was removed and cells quickly washed with PBS. Cells were incubated in PBS in the presence or absence of antibodies to be tested or reference compounds, 10 min at 37°C prior to the addition of coelenterazine with or without HGF in a final volume of 50 µl. After incubation for further 10 minutes at 37°C, light-emission acquisition at 485 nm and 530 nm was initiated using the Mithras luminometer (Berthold) (1s/wave length/well repeated 15 times).

BRET ratio has been defined previously [Angers et al., Proc. Natl. Acad. Sci. USA, 2000, 97:3684-3689] as:  $[(\text{emission at 530 nm}) - (\text{emission at 485 nm}) \times \text{Cf}] / (\text{emission at 485 nm})$ , where Cf corresponds to  $(\text{emission at 530 nm}) / (\text{emission at 485 nm})$  for cells expressing Rluc fusion protein alone in the same experimental conditions. Simplifying this equation shows that BRET ratio corresponds to the ratio 530/485 nm obtained when the two partners were present, corrected by the ratio 530/485 nm obtained under the same experimental conditions, when only the partner fused to *R. reniformis* luciferase was present in the assay. For the sake of readability,

results are expressed in milliBRET units (mBU); mBU corresponds to the BRET ratio multiplied by 1000.

After this second *in vitro* test, the antibody 224G11 i) without intrinsic activity as a whole molecule in the functional test of proliferation, ii) inhibiting significantly  
5 BxPC3 proliferation and iii) inhibiting c-Met dimerization was selected. In the experiments, the 5D5 Mab, generated by Genentech, and available at the ATCC, was added as a control for the intrinsic agonistic activity.

### **Example 2: Humanization process of mouse 224G11 Mab by CDR-grafting**

#### 10 1° Humanization of the light chain variable domain (VL)

As a preliminary step, the nucleotide sequence of 224G11 VL was compared to the murine germline gene sequences included in the IMGT database (<http://imgt.cines.fr>). Murine IGKV3-5\*01 and IGKJ4\*01 germline genes showing a sequence identity of 99.31 % for the V region and 94.28 % for the J region,  
15 respectively, have been identified. Regarding these high homologies, the 224G11VL nucleotide sequence has been used directly to search for human homologies, instead of corresponding mouse germlines.

In a second step, the human germline gene displaying the best identity with the 224G11VL has been searched to identify the best human candidate for the CDR  
20 grafting. For optimization of the selection, alignments between the amino acid sequences have been performed. The human IGKV4-1\*01 germline gene yielded a sequence identity of 67.30 %, but showed a different length for CDR1 (10 amino acids in 224G11 VL and 12 amino acids in IGKV4-1\*01). For the J region, the human IGKJ4\*02 germline gene (sequence identity of 77.14 %) was selected.

25 In a next step, mouse 224G11 VL CDR regions were engrafted into the above selected human framework sequences. Each amino acid position was analyzed for several criteria such as participation in VH/VL interface, in antigen binding or in CDR structure, localization of the residue in the 3D structure of the variable domain, CDR anchors, residues belonging to the Vernier zone. Three humanized versions,  
30 corresponding to SEQ ID No. 8, SEQ ID No. 9 and SEQ ID No. 10 were constructed, and containing respectively four (4, 39, 40, 84), two (39, 40) or one (40) murine residues in their FR regions and the CDRs corresponding to mouse 224G11 VL.

## 2°) Humanization of the heavy chain variable domain (VH)

As a preliminary step, the nucleotidic sequence of the 224G11 VH was compared to the murine germline genes sequences included in the IMGT database (<http://imgt.cines.fr>).

5 Murine IGHV1-18\*01, IGHD2-4\*01 and IGHJ2\*01 germline genes with a sequence identity of 92.70 % for the V region, 75.00 % for the D region and 89.36 % for the J region, respectively, have been identified. Regarding these high homologies, it has been decided to use directly the 224G11VH nucleotide sequences to search for human homologies, instead of corresponding mouse germlines.

10 In a second step, the human germline gene displaying the best identity with the 224G11 VH has been searched to identify the best human candidate for the CDR grafting. To this end, the nucleotidic sequence of 224G11 VH has been aligned with the human germline genes sequences belonging to the IMGT database. The human IGHV1-2\*02 V sequence exhibited a sequence identity of 75.00 % at the nucleotide level and  
15 64.30 % at the amino acid level. Looking for homologies for the J region led to the identification of the human IGHJ4\*04 germline gene with a sequence identity of 78.72 %.

In a next step, mouse 224G11 VH CDR regions were engrafted into the above selected human framework sequences. Each amino acid position was analyzed for  
20 several criteria such as participation in VH/VL interface, in antigen binding or in CDR structure, localization of the residue in the 3D structure of the variable domain, CDR anchors, residues belonging to the Vernier zone. One fully humanized form, corresponding to SEQ ID 4 was constructed; it contains exclusively human residues in its FR regions and the CDRs corresponding to mouse 224G11 VH.

25

### **Example 3: Engineering of improved hinge mutants**

It is well known by the skilled artisan that the hinge region strongly participates in the flexibility of the variable domain of immunoglobulins (see Brekke et al., 1995; Roux et al., 1997). During the chimerization process of 224G11 Mab, the mouse  
30 constant domain IGHG1 was replaced by the equivalent IGHG1 portion of human origin. Since the amino acid sequence of the hinge region were highly divergent, "murinization" of the hinge region was performed in order to keep its length and

rigidity. Since the human IGHG2 hinge region corresponds to the closest homologue of the mouse IGHG1 hinge, this sequence was as well considered. A series of 7 different hinge sequences were constructed (SEQ ID Nos. 22 to 28) by incorporating portions of the mouse IGHG1 and the human IGHG2 hinges into the human IGHG1 hinge portion.

5 Another series of hinge mutants was designed and constructed (SEQ ID Nos. 58 to 72) to evaluate the influence of either an additional cysteine and its position along the hinge domain, deletion of 1, 2, 3 or 4 amino acids along the hinge domain and a combination of these two parameters (cysteine addition and amino acid deletion).

#### 10 **Example 4: Production of humanized 224G11 Mab and engineered hinge Mab formats**

All above described Mab forms containing either chimeric, humanized and/or engineered hinge regions were produced upon transient transfection and by using the HEK293/EBNA system with a pCEP4 expression vector (InVitrogen, US).

15 The entire nucleotide sequences corresponding to the humanized versions of the variable domain of 224G11 Mab light (SEQ ID No. 18, SEQ ID No. 19 and SEQ ID No. 20) and heavy (SEQ ID No. 14) chains were synthesized by global gene synthesis (Genecust, Luxembourg). They were subcloned into a pCEP4 vector (InVitrogen, US) carrying the entire coding sequence of the constant domain [CH1-Hinge-CH2-CH3] of a  
20 human IgG1 or IgG2 immunoglobulin. Modification of the hinge region was performed by exchanging a {*NheI*-*BclI*} restriction fragment by the equivalent portion carrying the desired modifications, each respective {*NheI*-*BclI*} fragment being synthesized by global gene synthesis (Genecust, LU). All cloning steps were performed according to conventional molecular biology techniques as described in the Laboratory manual  
25 (Sambrook and Russel, 2001) or according to the supplier's instructions. Each genetic construct was fully validated by nucleotide sequencing using Big Dye terminator cycle sequencing kit (Applied Biosystems, US) and analyzed using a 3100 Genetic Analyzer (Applied Biosystems, US).

Suspension-adapted HEK293 EBNA cells (InVitrogen, US) were routinely  
30 grown in 250 ml flasks in 50 ml of serum-free medium Excell 293 (SAFC Biosciences) supplemented with 6 mM glutamine on an orbital shaker (110 rpm rotation speed). Transient transfection was performed with  $2 \cdot 10^6$  cells/ml using linear 25 kDa

polyethyleneimine (PEI) (Polysciences) prepared in water at a final concentration of 1 mg/ml mixed and plasmid DNA (final concentration of 1.25 µg/ml for heavy to light chain plasmid ratio of 1:1). At 4 hours post-transfection, the culture was diluted with one volume of fresh culture medium to achieve a final cell density of 10<sup>6</sup> cells/ml.

5 Cultivation process was monitored on the basis of cell viability and Mab production. Typically, cultures were maintained for 4 to 5 days. Mabs were purified using a conventional chromatography approach on a Protein A resin (GE Healthcare, US).

All different forms of Mabs were produced at levels suitable with functional evaluations. Productivity levels are typically ranging between 15 and 30 mg/l of  
10 purified Mabs.

#### **Example 5: Evaluation of c-Met phosphorylation status by a Phospho-c-Met-specific ELISA assay**

This functional assay allows to monitor modulation c-Met phosphorylation  
15 status either by Mabs alone or in the co-presence of HGF.

A549 cells were seeded in a 12MW plate in complete growth medium [F12K + 10 % FCS]. Cells were starved for 16 hours before stimulation with HGF [100 ng/ml], and each Mab to be tested was added at its final concentration of 30 µg/ml 15 minutes prior to ligand stimulation. Ice-cold lysis buffer was added 15 minutes after the addition  
20 of HGF to stop the phosphorylation reaction. Cells were scraped mechanically and cell lysates were collected by centrifugation at 13000 rpm for 10 min. at 4°C and correspond to the supernatant phase. Protein content was quantified using a BCA kit (Pierce) and stored at -20°C until use. The phosphorylation status of c-Met was quantified by ELISA. A goat anti-c-Met Mab (R&D, ref AF276) was used as a capture antibody  
25 (overnight coating at 4°C) and after a saturation step with a TBS-BSA 5% buffer (1 hour at room temperature (RT)), 25 µg of protein lysates were added to each well of the coated 96MW plate. After a 90 minutes incubation at RT, plates were washed four time and the detection antibody was added (anti-phospho-c-Met Mab, directed against the phosphorylated Tyr residues at position 1230, 1234 and 1235). After an additional  
30 hour incubation and 4 washes, an anti-rabbit antibody coupled to HRP (Biosource) was added for 1 hour at RT, and the luminescence detection was performed by adding Luminol. Luminescence readings were on a Mithras LB920 multimode plate reader



(Berthold).

Both basal and HGF [100 ng/ml]-induced c-Met receptor phosphorylation level were unaffected neither by PBS treatment, nor by the addition of mouse or human Mabs which do not target human c-Met receptor (Figure 1). On the other hand, mouse (m) 224G11 Mab strongly inhibited HGF [100 ng/ml]-induced c-Met phosphorylation (Figure 2B) without altering by itself receptor phosphorylation (Figure 2A). Surprisingly, the chimeric form of 224G11 Mab (224G11chim/IgG1), meaning variable domain (VH+VL) from m224G11 combined with human constant domain IgG1/kappa yielded strong (17 % of maximal HGF effect, Figure 2A) agonist activity associated with a reduced antagonist efficacy (54 % inhibition of HGF maximal effect compared to the m224G11 that yields 75% inhibition of HGF maximum effect, Figure 2B). Three humanized forms of 224G11 Mab, [224G11]Hz1/IgG1, [224G11]Hz2/IgG1 and [224G11]Hz3/IgG1, also constructed on a human IgG1/kappa backbone, yielded also decreased antagonist efficacy and significant agonist activity (11 to 24 % of maximal HGF level) as compared to mouse 224G11 (Figures 2A and 2B). A series of engineered versions of the heavy chain hinge domain were constructed and assayed in the c-Met receptor phosphorylation assay. As shown in Figure 3A, an important reduction of the agonist effect associated with the hIgG1/kappa isotype was observed for both the IgG2-based construct and for engineered IgG1/kappa constructs [MH, MUP9H and TH7]. A concomitant increase in antagonist efficacy was as well obtained. The hIgG1/kappa-based TH7 hinge mutant, with the most human sequence, was selected to complete the humanization process. In a next step, three humanized versions of 224G11 Mab variable domain were generated by combination to either a human IgG2/kappa or an IgG1/kappa-based TH7 engineered hinge constant domain. For the hIgG2/kappa humanized constructs, the humanized version Hz3 yielded strong agonism (Figure 4A), and for all three humanized versions, the antagonist efficacy was below that observed with murine 224G11 Mab and comparable to the chimeric hIgG1-based Mab (56-57 % inhibition of HGF effect, Figure 4B). On the other hand, combination of the three humanized versions Hz1, Hz2 or Hz3 to the engineered IgG1/TH7 mutant almost fully restored the properties of mouse 224G11 Mab in terms of weak agonist activity (5-6 % of HGF effect) and strong antagonist efficacy (68 to 72 % inhibition of HGF effect) of c-Met receptor phosphorylation (Figures 5A and 5B). These variants were highly

improved as compared to chimeric IgG1-based 224G11 Mab but also to IgG2-based humanized forms.

A second series of engineered versions of the heavy chain hinge domain was constructed and assayed in the c-Met receptor phosphorylation assay. As shown in figure 17A, all those new versions (c224G11[C2], c224G11[C3], c224G11[C5], c224G11[C6], c224G11[C7], c224G11[Δ1-3], c224G11[C7Δ6], c224G11[C6Δ9], c224G11[C2Δ5-7], c224G11[C5Δ2-6], c224G11[C9Δ2-7] and c224G11[Δ5-6-7-8]) exhibited weaker agonist effect than c224G11 since their agonism activities are comprised between 6 and 14% of the HGF effect compared to 23% for c224G11. As c224G11[TH7], all those new versions exhibited a concomitant increase in antagonist efficacy [figure 17B]. Those results showed that engineering of the heavy chain domain by point mutation and/or deletion could modify agonistic/antagonistic properties of an antibody.

#### Example 6: BRET analysis

In a first set of experiments, it had been control that irrelevant mouse IgG1, human IgG1 and human IgG2 had no effect of HGF induced BRET signal in both BRET models (representative experiment out of 12 independent experiments; Figure 6). These Mabs are forthwith cited as controls.

The effect of a IgG1 chimeric form of mouse 224G11 Mab ([224G11]chim) on both c-Met dimerization and c-met activation BRET model was evaluated. While mouse 224G11 Mab inhibited 59.4% of the HGF induced BRET signal on c-Met dimerization model, [224G11]chim Mab inhibited only 28.9% (Figure 7A). [224G11]chim antibody was also less effective in inhibiting HGF induced c-Met activation since [224G11]chim and m224G11 antibodies inhibited respectively 34.5% and 56.4% of HGF induced BRET signal (Figure 7B). Moreover, m224G11 alone had no effect on c-Met activation while [224G11]chim had a partial agonist effect on c-Met activation corresponding to 32.9% of the HGF induced signal. This partial agonist effect of the [224G11]chim was also seen on c-Met dimerization BRET model since [224G11]chim alone induced a BRET increase corresponding to 46.6% of HGF-induced signal *versus* 21.3% for m224G11 (Figure 7A).

In Figures 8A and 8B, hinge mutated chimeric forms of 224G11 antibody

showed a greater inhibitory effect on HGF induced BRET signal than [224G11]chim since they showed a 59.7%, 64.4%, 53.2% and 73.8% inhibition of the HGF induced activation BRET signal (Figure 8B) and 61.8%, 64.4% 52.5% and 64.4% inhibition of the HGF induced c-Met dimerization BRET signal (Figure 8A) for [224G11][MH chim], [224G11][MUP9H chim], [224G11][MMCH chim] and [224G11][TH7 chim] respectively. Contrary to [224G11]chim, which had a partial agonist effect on c-Met activation, hinge mutated chimerical forms of 224G11 antibody showed no significant effect on c-Met activation alone (5.1%, 7.6%, -2.0% and -6.9% respectively) as observed for m224G11.

10 In Figure 9B, like the [224G11] [TH7 chim], the 3 humanized versions of 224G11 IgG1 antibody with the TH7 hinge induced no significant increased of BRET signal in activation model when tested alone and showed a strong inhibition of HGF induced BRET signal: 59.9%, 41.8% and 57.9% for the Hz1, Hz2 and Hz3 forms respectively. Moreover, [224G11] [TH7 Hz1], [224G11] [TH7 Hz2] and [224G11] [TH7 Hz3] inhibited HGF induced BRET signal on dimerization model of 52.2%, 15 35.8% and 49.4% respectively (figure 9A).

Contrary to [224G11]chim, the chimeric form of 224G11 IgG2 antibody ([224G11] [IgG2 chim]) showed no partial agonist effect alone and inhibited 66.3% of the HGF effect on c-Met activation model (Figure 10B). On c-Met dimerization model, 20 [224G11] [IgG2 chim] inhibited 62.4% of the HGF induced BRET signal (Figure 10A).

The agonist efficacy of the second series of engineered versions of the heavy chain hinge domain was evaluated in c-Met activation BRET model (Figure 18). In contrast to c224G11, which had a partial agonist effect on c-Met activation, c224G11[C2], c224G11[C3], c224G11[C5], c224G11[C6], c224G11[C7], 25 c224G11[Δ1-3], c224G11[C7Δ6], c224G11[C6Δ9], c224G11[C2Δ5-7], c224G11[C5Δ2-6], c224G11[C9Δ2-7] and c224G11[Δ5-6-7-8] hinge mutated chimeric forms of 224G11 antibody showed no significant effect on c-Met activation alone.

#### **Example 7: c-Met recognition by chimeric and humanized 224G11 forms**

30 A direct ELISA has been set up to determine the binding ability of the various chimeric and humanized forms on the recombinant c-Met. Briefly recombinant dimeric c-Met from R&D Systems was coated at 1.25 μg/ml on 96-well Immunlon II plates.

After an overnight incubation at 4°C, wells were saturated with a 0.5% gelatine/PBS solution. Plates were then incubated for 1 hour at 37°C before addition of 2 fold dilutions of antibodies to be tested. Plates were incubated an additional hour before addition of a goat anti-mouse IgG HRP for detecting the murine antibody and a goat anti-human Kappa light chain HRP for chimeric and humanized antibody recognition. Plates were incubated for one hour and the peroxidase substrate TMB Uptima was added for 5 mn before neutralization with H<sub>2</sub>SO<sub>4</sub> 1M. Results presented in Figure 11 showed that all tested forms were comparable for c-Met recognition.

10           **Example 8: Effect of murine and chimeric 224G11 on HGF-induced proliferation of NCI-H441 cells in vitro**

NCI-H441 cells from ATCC were routinely cultured in RPMI 1640 medium (Invitrogen Corporation, Scotland, UK), 10% FCS (Invitrogen Corporation), 1% L-Glutamine (Invitrogen corporation). For proliferation assays, cells were split 3 days before use so that they were in the confluent phase of growth before plating. NCI-H441 cells were plated in 96-well tissue culture plates at a density of 3.75x10<sup>4</sup> cells/well in 200 µl of serum free medium (RPMI 1640 medium plus 1% L-Glutamine). Twenty four hours after plating, antibodies to be tested were added to NCI-H441 and incubated at 37°C for thirty minutes before adding HGF at a final concentration of 400 ng/ml (5 nM) for 142 additional hours. The dose range tested for each antibody is from 10 to 0.0097 µg/ml (final concentration in each well). In this experiment, a murine IgG1 Mab was added as a murine isotype control and the tested antibodies were the following one: m224G11 and its human IgG1 chimeric form identified as [224G11]chim. Wells plated with cells alone +/- HGF were also included. Then cells were pulsed with 0.25 µCi of [<sup>3</sup>H]Thymidine (Amersham Biosciences AB, Uppsala, Sweden) for 7 hours and 30 minutes. The magnitude of [<sup>3</sup>H]Thymidine incorporated in trichloroacetic acid-insoluble DNA was quantified by liquid scintillation counting. Results are expressed as non transformed cpm data to better evaluate the potential intrinsic agonist activity that could occur with anti-c-Met Mabs when added alone to tumour cell.

30           Results described in Figure 12 demonstrated that, as expected, the murine antibody m224G11 displayed no agonist effect when added alone to cancer cells whatever the tested dose. No significant inhibition of the HGF-induced proliferation

was observed with the isotype control regarding to the cpm variations observed for this compound in this experiment. When added alone, the m224G11 antibody did not show any agonist effect compared to the mIgG1 isotype control Mab or cells alone. A dose dependent anti-proliferative activities reaching 78% was observed for m224G11 (% inhibition calculation:  $100 - [(cpm\ cells + Mab\ to\ be\ tested - mean\ cpm\ background\ mIgG1) \times 100 / (mean\ cpm\ cells + HGF - mean\ cpm\ cells\ alone)]$ ). Surprisingly, the chimeric form of the 224G11 Mabs induced a significant, dose dependent agonist effect when added alone. This agonist effect had an impact on the *in vitro* inhibition of HGF-induced proliferation that shifted from 78% for the murine 224G11 to 50% for its chimeric form. To determine whether such “lower” *in vitro* intrinsic agonist activity was compatible with an unchanged *in vivo* effect, both m224G11 and [224G11]chim were produced for *in vivo* testing. As, in previous studies, the 30 µg/mice dose had demonstrated a significant *in vivo* activity, that dose was selected for *in vivo* evaluation.

15           **Example 9: *In vivo* comparison of murin and chimeric 224G11 Mabs on the NCI-H441 xenograft model**

NCI-H441 is derived from papillary lung adenocarcinoma, expresses high levels of c-Met, and demonstrates constitutive phosphorylation of c-Met RTK.

To evaluate the *in vivo* effect of antibodies on the NCI-H441 xenograft model, six to eight weeks old athymic mice were housed in sterilized filter-topped cages, maintained in sterile conditions and manipulated according to French and European guidelines. Mice were injected subcutaneously with  $9 \times 10^6$  cells. Then, six days after cell implantation, tumors were measurable (approximately 100 mm<sup>3</sup>), animals were divided into groups of 6 mice with comparable tumor size and treated first with a loading dose of 60 µg of antibody/mice and then twice a week with 30 µg/dose of each antibody to be tested. The mice were followed for the observation of xenograft growth rate. Tumor volume was calculated by the formula:  $\pi (Pi)/6 \times length \times width \times height$ . Results described in Figure 13 demonstrate that the murine Mab devoided of agonist activity *in vivo* behave, as expected, as potent antagonist even at the low tested dose. In contrast to what observed with the murine Mab, the chimeric one displayed a very transient *in vivo* activity and tumor completely escaped to the treatment at D20 post cell injection. This experiment demonstrates clearly that the increase of *in vitro* agonist

effect that resulted in a decrease of antagonist activity was also responsible for a significant *in vivo* loss of antagonist activity.

**Example 10: Effect of the murine 224G11 Mab and of various chimeric and humanized versions of this antibody on HGF-induced proliferation of NCI-H441 cells *in vitro***

NCI-H441 cells from ATCC were routinely cultured in RPMI 1640 medium (Invitrogen Corporation, Scotland, UK), 10% FCS (Invitrogen Corporation), 1% L-Glutamine (Invitrogen Corporation). For proliferation assays, cells were split 3 days before use so that they were in the confluent phase of growth before plating. NCI-H441 cells were plated in 96-well tissue culture plates at a density of  $3.75 \times 10^4$  cells/well in 200  $\mu$ l of serum free medium (RPMI 1640 medium plus 1% L-Glutamine). Twenty four hours after plating, antibodies to be tested were added to NCI-H441 and incubated at 37°C for thirty minutes before adding HGF at a final concentration of 400 ng/ml (5 nM) for 142 additional hours. The dose range tested for each antibody is from 10 to 0.0097  $\mu$ g/ml (final concentration in each well). In this experiment, murine IgG1 Mab was added as a murine isotype control and as an agonist negative control. The tested antibodies were the following one: i) m224G11, ii) its human IgG1 chimeric forms respectively identified as [224G11] chim, [224G11] [MH chim], [224G11] [MUP9H chim], [224G11] [MMCH chim], [224G11] [TH7 chim] iii) its humanized IgG1 forms respectively described as [224G11] [Hz1], [224G11] [Hz2], [224G11] [Hz3]. Wells plated with cells alone +/- HGF were also included. The 5D5 whole antibody from Genentech commercially available at the ATCC as an hybridoma cell line was introduced as a full agonist positive control and thereafter called m5D5. Then cells were pulsed with 0.25  $\mu$ Ci of [<sup>3</sup>H]Thymidine (Amersham Biosciences AB, Uppsala, Sweden) for 7 hours and 30 minutes. The magnitude of [<sup>3</sup>H]Thymidine incorporated in trichloroacetic acid-insoluble DNA was quantified by liquid scintillation counting. Results are expressed as non transformed cpm data to better evaluate the potential intrinsic agonist activity that could occur with anti-c-Met Mabs when added alone to tumour cell.

Results described in Figure 14A demonstrated that as expected neither the isotype control nor the m224G11 displayed any agonist activity on NCI-H441

proliferation. The isotype control was without effect on HGF-induced cell proliferation whereas m224G11 showed a 66% inhibition when added at the final concentration of 10 µg/ml. The m5D5 used as an agonist control showed, as expected, a full dose dependent agonist effect when added alone to the cells. As already observed, the [224G11] chim Mab displayed a significant dose- dependent agonist effect and, a decreased inhibitory activity of this chimeric form was observed: 19% instead of 66% for the murine form. When added alone, the 3 IgG1 humanized Mabs demonstrated dose dependent agonist effects compared to the m224G11 form. [224G11] [Hz1], [224G11] [Hz2] and [224G11] [Hz3] had comparable antagonist activities about 46, 30 and 35%. These activities are significantly lower than the one observed for m224G11. In Figure 14B, various IgG1 chimeric forms were tested. Compared to [224G11] chim form which displayed a dose-dependent agonist effect when added alone to NCI-H441 cells, the [224G11] [MH chim], [224G11] [MUP9H chim], [224G11] [MMCH chim], [224G11] [TH7 chim] forms were without significant intrinsic agonist effect. Their antagonist activity was higher than the one observed for the m224G11 Mab (57%) with inhibitions reaching 79, 78, 84 and 93% respectively for [224G11] [MH chim], [224G11] [MUP9H chim], [224G11] [MMCH chim] and [224G11] [TH7 chim].

**Example 11: *In vitro* effect of various IgG1 humanized form of the 224G11 Mab**

NCI-H441 cells from ATCC were routinely cultured in RPMI 1640 medium (Invitrogen Corporation, Scotland, UK), 10% FCS (Invitrogen Corporation), 1% L-Glutamine (Invitrogen Corporation). For proliferation assays, cells were split 3 days before use so that they were in the confluent phase of growth before plating. NCI-H441 cells were plated in 96-well tissue culture plates at a density of  $3.75 \times 10^4$  cells/well in 200 µl of serum free medium (RPMI 1640 medium plus 1% L-Glutamine). Twenty four hours after plating, antibodies to be tested were added to NCI-H441 and incubated at 37°C for thirty minutes before adding HGF at a final concentration of 400 ng/ml (5 nM) for 142 additional hours. The dose range tested for each antibody is from 10 to 0.0097 µg/ml (final concentration in each well). In this experiment, murine IgG1 Mab was added as a background negative control for agonist activity and the tested antibodies were the following one: i) m224G11, ii) its human IgG1 chimeric forms

respectively identified as [224G11] chim, [224G11] [TH7 chim] iii) its humanized IgG1 forms respectively described as [224G11] [TH7 Hz1], [224G11] [TH7 Hz3]. Wells plated with cells alone +/- HGF were also included. The 5D5 whole antibody from Genentech commercially available at the ATCC as an hybridoma cell line was introduced as a full agonist positive control and thereafter called m5D5. Then cells were pulsed with 0.25  $\mu$ Ci of [<sup>3</sup>H]Thymidine (Amersham Biosciences AB, Uppsala, Sweden) for 7 hours and 30 minutes. The magnitude of [<sup>3</sup>H]Thymidine incorporated in trichloroacetic acid-insoluble DNA was quantified by liquid scintillation counting. Results are expressed as non transformed cpm data to better evaluate the potential intrinsic agonist activity that could occur with anti-c-Met Mabs when added alone to tumour cell.

Figure 15 showed that the m224G11 Mab displayed the usual inhibitory effect (74% inhibition). The chimeric IgG1 form [224G11] chim had as expected a dose dependent intrinsic agonist effect and a lower antagonist effect compared to the murin form: 33% versus 74% inhibition. The [224G11] [TH7 chim] had a very weak agonist activity in this experiment. However it displayed a high inhibitory effect (81 %) close to the one noticed for the murine Mab. The 2 humanized forms had no intrinsic agonist effect and had an antagonist activity close to the ones observed for the murine Mab or the [224G11] [TH7 chim] with respectively 67 and 76% inhibition for [224G11] [TH7 Hz1] and [224G11] [TH7 Hz3].

**Example 12: *In vivo* comparison of murin, chimeric and humanized 224G11 Mabs bearing either the wild type or the TH7- engineered hinge (NCI-H441 xenograft model).**

NCI-H441 is derived from papillary lung adenocarcinoma, expresses high levels of c-Met, and demonstrates constitutive phosphorylation of c-Met RTK.

To evaluate the necessity of hinge engineering to save *in vivo* activity of the 224G11 murine antibody, six to eight weeks old athymic mice were housed in sterilized filter-topped cages, maintained in sterile conditions and manipulated according to French and European guidelines. Mice were injected subcutaneously with  $9 \times 10^6$  NCI-H441 cells. Then, six days after cell implantation, tumors were measurable (approximately 100 mm<sup>3</sup>), animals were divided into groups of 6 mice with comparable



tumor size and treated first with a loading dose of 2 mg of antibody/mice and then twice a week with a 1 mg/dose of each antibody to be tested. Ten antibodies were evaluated in this experiment including the m224G11, the chimeric form displaying the wild type hinge (c224G11), the TH7-engineered chimeric form (224G11[TH7 chim]), three  
5 humanized form bearing the wild type hinge (224G11[IgG1 Hz1], 224G11[IgG1 Hz2] and 224G11[IgG1 Hz3]) and the three corresponding TH7-engineered forms (224G11[TH7 Hz1], 224G11[TH7 Hz2] and 224G11[TH7 Hz3]). Mice were followed for the observation of xenograft growth rate.

Tumor volume was calculated by the formula:  $\pi (Pi)/6 \times \text{length} \times \text{width} \times$   
10 height.

Results described in figure 16 demonstrate that the murine Mab devoid of any agonist activity *in vitro* behave, as expected, as potent *in vivo* antagonist. In contrast to what observed with the murine Mab, both chimeric and humanized forms bearing the wild type hinge displayed only a very transient *in vivo* activity. In any cases the  
15 substitution of the wild type hinge by the TH7-engineered one resulted in a complete restoration of the *in vivo* activity observed with murine antibodies. This experiment demonstrates clearly that the increase of *in vitro* agonist effect that resulted in a decrease of antagonist activity was also responsible of a significant *in vivo* loss of antagonist activity. It also demonstrates that the use of a TH7-engineered region instead  
20 of the wild type one is needed for keeping the *in vivo* properties of the murine Mab.

### **Example 13: Effect of m224G11 and its humanized form h224G11 on c-Met downregulation *in vitro***

In the following examples, for the avoidance of doubt, the expression h224G11  
25 refers to the humanized form 224G11 [TH7 Hz3] of the antibody of the invention.

Two cell lines have been selected to address the activity of anti-c-Met antibodies on c-Met receptor degradation. A549 (#HTB-174) and NCI-H441 (#CCL-185) are two NSCLC cell lines from the ATCC collection. NCI-H441 cells were seeded in RPMI 1640 + 1% L-glutamine + 10% heat-inactivated FBS, at  $3 \times 10^4$  cells/cm<sup>2</sup> in six-well  
30 plates for 24 h at 37°C in a 5% CO<sub>2</sub> atmosphere. A549 cells were seeded in F12K + 10% heat-inactivated FBS, at  $2 \times 10^4$  cells/cm<sup>2</sup> in six-well plates for 24 h at 37°C in a 5% CO<sub>2</sub> atmosphere.

Then, cells were washed twice with phosphate buffer saline (PBS) before being serum-starved for 24 additional hours. Anti-c-Met antibodies (10 µg/ml), irrelevant mIgG1(10 µg/ml), or HGF (400 ng/mL) were added in serum-free DMEM medium at 37°C. After either 4 hours or 24 hours of incubation, the medium was gently removed  
5 and cells washed twice with cold PBS. Cells were lysed with 500 µL of ice-cold lysis buffer [50 mM Tris-HCl (pH 7.5); 150 mM NaCl; 1% Nonidet P40; 0.5% deoxycholate; and 1 complete protease inhibitor cocktail tablet plus 1% antiphosphatases]. Cell lysates were shaken for 90 min at 4°C and cleared at 15 000 rpm for 10 minutes. At this stage, cell lysates could be stored at -20°C until needed for western blot analysis. Protein  
10 concentration was quantified using BCA. Whole cell lysates (5µg in 20 µl) were separated by SDS-PAGE and transferred to nitrocellulose membrane. Membranes were saturated for 1 h at RT with TBS-Tween 20 0.1%(TBST); 5% non-fat dry milk and probed with anti-c-Met antibody (dilution 1/1000) overnight at 4°C in TBST-5% non-fat dry milk. Antibodies were diluted in tris-buffered saline-0.1% tween 20 (v/v)  
15 (TBST) with 1 % non-fat dry milk. Then, membranes were washed with TBST and incubated with peroxylase-conjugated secondary antibody (dilution 1:1000) for 1 h at RT. Immunoreactive proteins were visualized with ECL (Pierce # 32209). After c-Met visualization, membranes were washed once again with TBST and incubated for 1 h at RT with mouse anti-GAPDH antibody (dilution 1/200 000) in TBST-5% non-fat dry  
20 milk. Then, membranes were washed in TBST and incubated with peroxylase-conjugated secondary antibodies, for 1 h at RT. Membranes were washed and GAPDH was revealed using ECL. Band intensity was quantified by densitometry.

Results presented in figures 19A and 20A demonstrated that both m224G11 and h224G11 are able to significantly downregulate c-Met, in a dose-dependant way, in  
25 both A549 and NCI-H441 cell lines. The downregulation is already significant after a 4 hour incubation time and still increased at 24 hour. Histograms presented in figures 19A and 20A corresponds to mean values or respectively 4 and 3 independent experiments. Western blot images corresponding to one significant experiment were included in figures 19B and 20B.

30

**Example 14: Effect of m224G11 and its humanized form h224G11 on c-Met shedding *in vitro***

Soluble shedded forms of the c-Met receptor occur naturally in the serum of mice xenografted with human tumor or in serum of human patient carrying tumors expressing c-Met. Moreover, antibodies directed against c-Met such as the DN30 Mab, are described as shedding inducers of c-Met in *in vitro* experiments. To determine whether the m224G11 has such a property, cells were seeded in six-well plates in 10% FCS medium. When they reached approximately 80% confluence, medium was removed and fresh complete culture medium +/- compounds to be tested was added. Cells were incubated 72 additional hours with either m224G11, an isotype control mIgG1 or PBS. PMA (phorbol myristate acetate) was introduced as a shedding inducer. HGF was also tested on cells to determine the impact of c-Met ligand on natural occurring shedding. Then supernatants were collected and filtered on 0.2 µm before use in an ELISA test which soluble forms of c-Met were captured with an anti-c-Met antibody that does not recognize the same epitope as either m224G11 or the c11E1 (Figure 21). Moreover, cells from each well were washed once with PBS and lysed to determine protein concentration. For the ELISA, 224D10 was used as a capture antibody and after plate saturation, filtered supernatants from six well plates were added in the ELISA test. A monomeric c-Met form was used as a positive control. After supernatant incubation, plates were washed to remove the unbound c-Met and c11E1 was used to detect c-Met captured by the 224G11 Mab. The revelation of the test was finally performed by addition of an HRP-conjugated anti-hFc polyclonal antibody.

Results shown in figure 22 indicate that a natural shedding of c-Met occurred when cells were cultured for 72 hours *in vitro*. No effect of the mIgG1 was observed. However, the addition of m224G11 seemed to inhibit c-Met shedding. These results were confirmed for 3 other cells lines (Hs746T, EBC1 and MKN45) in figure 23. In that second experiment, the PMA, added as a positive shedding inducer, increased significantly, as expected, c-Met shedding at least in 2 cell lines (Hs746T and MKN45). Finally, in a third experiment (Figure 24), HGF was introduced as a control. No additional shedding was induced by HGF compared as cells alone or cells + mIgG1. Once again, a significant inhibition of c-Met shedding was observed with m224G11.

30

#### **Example 15: Intrinsic effect of h224G11 Ab on various cell lines**

In previous experiments described in this patent, it has been demonstrated that in

contrast to what was observed with other antibodies such as 5D5, the m224G11 and its humanized form h224G11 do not display significant intrinsic activity tumor cell lines. To extend this property to other cell lines, western blot and phospho-ELISA experiments were performed with the antibody alone, added for various times, on a set of cancer cell lines, with variable levels of c-Met expression, including Hs746T, NCI-H441, Hs578T, NCI-H125, T98G, MDA-MB-231, PC3. The same test was also performed in a normal cell: HUVEC.

Method for the phospho cMet ELISA assay was already described in example 5 of the present patent application. For the western analysis, protein lysates were made from pelleted cells by incubation in lysis buffer with proteases and phosphatase inhibitors [10nM Tris (pH 7.4), 150 mM NaCl, 1mM EDTA, 1 mM EGTA, 0.5% Nonidet P40, 100 mM sodium fluoride, 10 mM sodium pyrophosphate, 2 mM sodium orthovanadate, 2 mM PMSF, 10 mg/ml leupeptin, 10 mg/ml aprotinin] at 4°C. Protein lysates were cleared of cellular debris by centrifugation, resolved by electrophoresis on 8% SDS-PAGE gels, and electrotransferred to a nitrocellulose membrane. For c-Met experiments, lysates were immunoprecipitated for specific protein of interest before electrophoresis and transfer.

Results presented in figures 25 to 32 demonstrate once again that no intrinsic activity of the h224G11 antibody was observed in the tested cells.

20

**Example 16: In vivo comparison of the murin wild type 224G11 with a chimeric hinge-engineered 224G11 form described as 224G11[C2D5-7] (NCI-H441 xenograft model)**

NCI-H441 is derived from papillary lung adenocarcinoma, expresses high levels of c-Met, and demonstrates constitutive phosphorylation of c-Met RTK.

To evaluate the necessity of hinge engineering to save in vivo activity of the 224G11 murine antibody, six to eight weeks old athymic mice were housed in sterilized filter-topped cages, maintained in sterile conditions and manipulated according to French and European guidelines. Mice were injected subcutaneously with  $9 \times 10^6$  NCI-H441 cells. Then, six days after cell implantation, tumors were measurable (approximately 100 mm<sup>3</sup>), animals were divided into groups of 6 mice with comparable tumor size and treated first with a loading dose of 2 mg of antibody/mice and then twice

30

a week with a 1 mg/dose of each antibody to be tested. Mice were followed for the observation of xenograft growth rate. Tumor volume was calculated by the formula:  $\pi$  (Pi)/6 X length X width X height. Results described in figure 33 demonstrate that the murine Mab devoid of any agonist activity in vitro behave, as expected, as a potent in vivo antagonist. As suggested by the results obtained in vitro, in phosphorylation assays, the c224G11[C2D5-7] hinge-engineered antibody, that did not display a significant agonist effect, demonstrate a strong in vivo activity, comparable to the one of the m224G11 on the NCI-H441 xenograft model.

#### 10           **Example 17: Evaluation of h224G11 in an ADCC test**

As h224G11 is of IgG1 isotype, ADCC could be part of its *in vivo* efficacy in human. An *in vitro* [<sup>51</sup>Cr] release cytotoxicity assay was performed using either Hs746T or NCI-H441 cells as target cells and NK cells purified from human peripheral blood mononuclear lymphocytes.

15           Briefly, one million Hs746T or NCI-H441 target cells were incubated with or without 20 µg of h224G11 Ab in presence of 100 µCi of <sup>51</sup>Chromium (Perkin Elmer) for 1 hr. Then, 4 X 10<sup>3</sup> cells were plated with an increasing number of human natural killer (NK) cells isolated from peripheral blood mononuclear cells (PBMC) using a negative selection (Stemcell Technologies). Cells were incubated together for 4 additional hours at 37°C. Percent of cell lysis was calculated following the formula: [(experimental <sup>51</sup>Cr release - spontaneous <sup>51</sup>Cr release)/(full <sup>51</sup>Cr release - spontaneous <sup>51</sup>Cr release)] X 100. Spontaneous release represents the counts obtained when the target cells were cultured in absence of natural killer cells. Full release represents the counts obtained when the target cells were lysed with 1% Triton X-100. h224G11 significantly enhanced lysis of both Hs746T (Figure 34a) and NCI-H441 (Figure 34b) cells by 62.9% and 63.2%, respectively, at a ratio NK/Target cells of 100.

#### **Example 18: Immunohistochemical Studies (IHC)**

30           Procedures of Paraffin Embedded Tumors IHC Staining: 8 to 12 µM sections of frozen tumor were and immediately fixed in pre cooled acetone -20°C for 3 minutes. Slides were then cooled at room temperature for 30 minutes to 1 hour. After 2 washes in PBS the Endogenous peroxidase activity was blocked using Peroxidase Blocking

Reagent (Dako K4007) for five minutes. Sections were washed with PBS and incubated in avidin/biotin blocking reagent (Dako X0590) just before saturation of the non specific sites in PBS-BSA 4% for 30 minutes at room temperature. Then, slides were incubated with the biotinylated h224G11 (50 to 10  $\mu\text{g/ml}$ ) or human biotinylated IgG1/kappa (50 to 10  $\mu\text{g/ml}$ , the Binding Site) as negative control 2 hours at room temperature.

Sections were washed with PBS and incubated with Streptavidin-peroxydase complex universal (Dako K0679) for 30 to 45 minutes. 3-Amino-9-Ethylcarbazole was used for development of a red reaction product (Sigma). The slides were immersed in hematoxylin for 4 minutes to counterstain (Dako S3309).

Results are represented in Figure 35.

h224G11 differentially stains the cell membrane of various tumor types. In this immunohistochemistry procedure, the red reaction product correlates to positive staining of the cell membrane and lack of red reaction product correlates to negative staining and no visualization of the cell membrane. The IgG control, human IgG1/kappa is an isotype matched control.

A reference herein to a patent document or other matter which is given as prior art is not to be taken as an admission that that document or matter was known or that the information it contains was part of the common general knowledge as at the priority date of any of the claims.

Throughout the description and claims of the specification, the word "comprise" and variations of the word, such as "comprising" and "comprises", is not intended to exclude other additives, components, integers or steps.

## PCT

Print Out (Original in Electronic Form)  
 (This sheet is not part of and does not count as a sheet of the international application)

0-1	<b>Form PCT/RO/134 (SAFE) Indications Relating to Deposited Microorganism(s) or Other Biological Material (PCT Rule 13bis)</b>	
0-1-1	Prepared Using	<b>PCT Online Filing Version 3.5.000.204 MT/FOP 20020701/0.20.5.9</b>
0-2	<b>International Application No.</b>	
0-3	<b>Applicant's or agent's file reference</b>	<b>355644D27117</b>

<b>1</b>	<b>The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:</b>	
1-1	page	<b>15</b>
1-2	line	<b>18</b>
<b>1-3</b>	<b>Identification of deposit</b>	
1-3-1	Name of depositary institution	<b>CNCM Collection nationale de cultures de micro-organismes</b>
1-3-2	Address of depositary institution	<b>Institut Pasteur, 28, rue du Dr Roux, 75724 Paris Cedex 15, France</b>
1-3-3	Date of deposit	<b>14 March 2007 (14.03.2007)</b>
1-3-4	Accession Number	<b>CNCM I-3731</b>
<b>1-5</b>	<b>Designated States for Which Indications are Made</b>	<b>All designations</b>

## FOR RECEIVING OFFICE USE ONLY

0-4	<b>This form was received with the international application: (yes or no)</b>	Yes-RO/EP
0-4-1	Authorized officer	Bruno Gatinet - epo H04 -

## FOR INTERNATIONAL BUREAU USE ONLY

0-5	<b>This form was received by the International Bureau on:</b>	
0-5-1	Authorized officer	

**THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS**

1. A monoclonal antibody, or a divalent functional fragment or derivative thereof, capable to inhibit the c-Met dimerization, said antibody comprising a heavy chain comprising CDR-H1, CDR-H2 and CDR-H3 with respectively the amino acid sequences SEQ ID Nos. 1, 2 and 3; and a light chain comprising CDR-L1, CDR-L2 and CDR-L3 with respectively the amino acid sequences SEQ ID Nos. 5, 6 and 7, said antibody being further characterized in that it also comprises a hinge region comprising the amino acid sequence selected from the group consisting of SEQ ID Nos. 22 to 28 and SEQ ID Nos. 58 to 72.

2. The antibody of claim 1, or a divalent functional fragment or derivative thereof, wherein said hinge region comprises an amino acid sequence selected from the group consisting of SEQ ID Nos. 24, 26, 28, 59, 60, 61, 62, 63, 65, 66, 67, 68, 69, 70 and 71.

3. The antibody of claim 1, or a divalent functional fragment or derivative thereof, wherein said antibody consists of a chimeric antibody.

4. The antibody of claim 1, or a divalent functional fragment or derivative thereof, wherein said antibody consists of a humanized antibody.

5. The antibody of claim 4, or a divalent functional fragment or derivative thereof, comprising a heavy chain variable domain of sequence comprising the amino acid sequence SEQ ID No. 4; and a light chain variable domain of sequence comprising the amino acid sequence SEQ ID No. 8, 9 or 10.

6. The antibody of claim 5, or a divalent functional fragment or derivative thereof, comprising a heavy chain variable domain of sequence comprising the amino acid sequence SEQ ID No. 4; a light chain variable domain of sequence comprising the amino acid sequence SEQ ID No. 8; and a hinge region comprising the amino acid sequence SEQ ID No. 28.

7. The antibody of claim 5, or a divalent functional fragment or derivative thereof, comprising a heavy chain variable domain of sequence comprising the amino acid sequence SEQ ID No. 4; a light chain variable domain of sequence comprising the amino acid sequence SEQ ID No. 9; and a hinge region comprising the amino acid sequence SEQ ID No. 28.



8. The antibody of claim 5, or a divalent functional fragment or derivative thereof, comprising a heavy chain variable domain of sequence comprising the amino acid sequence SEQ ID No. 4; a light chain variable domain of sequence comprising the amino acid sequence SEQ ID No. 10; and a hinge region comprising the amino acid sequence SEQ ID No. 28.

9. An isolated nucleic acid, wherein it is chosen from the following nucleic acids:

a) a nucleic acid, DNA or RNA, coding for an antibody, or a divalent functional fragment or derivative thereof, as claimed in one of claims 1 to 8;

b) a nucleic acid as defined in a) comprising a DNA sequence comprising the sequences SEQ ID No. 11, SEQ ID No. 12, SEQ ID No. 13 and the sequences SEQ ID No. 15, SEQ ID No. 16 and SEQ ID No. 17;

c) a nucleic acid as defined in a) or b) comprising a DNA sequence comprising the sequences SEQ ID No. 14 and SEQ ID No. 18, 19 or 20;

d) the corresponding RNA nucleic acids of the nucleic acids as defined in b) or c);

e) the complementary nucleic acids of the nucleic acids as defined in a), b) and c).

10. An isolated nucleic acid according to claim 9, wherein the nucleic sequence coding for the hinge region of said antibody comprises the nucleic acid sequence SEQ ID Nos. 31 to 35 and SEQ ID Nos. 73 to 87 when, said hinge region has respectively one of the sequence SEQ ID Nos. 24 to 28 and SEQ ID Nos. 58 to 72.

11. A vector comprising a nucleic acid as claimed in claim 9 or 10.

12. A host cell comprising a vector as claimed in claim 11.

13. A transgenic animal with the exception of human being comprising at least one cell transformed by a vector as claimed in claim 11.

14. A process for production of an antibody, or a divalent functional fragment or derivative thereof, as claimed in one of claims 1 to 8, wherein it comprises the following stages:

a) culture in a medium and appropriate culture conditions of a cell as claimed in claim 11; and

b) the recovery of said antibody, or a divalent functional fragment or derivative thereof, thus produced starting from the culture medium or said cultured cells.

15. An antibody, or a divalent functional fragment or derivative thereof, obtained by a process as claimed in claim 14.

16. The antibody of claims 1 to 8 and 15 as a medicament.

17. A composition comprising by way of active principle a compound consisting of an antibody, or a divalent functional fragment or derivative thereof, as claimed in one of claims 1 to 8 or 15.

18. The composition of claim 17, wherein it comprises, moreover, as a combination product for simultaneous, separate or sequential use, an anti-tumoral antibody.

19. The composition of claim 17, wherein it comprises, moreover, as a combination product for simultaneous, separate or sequential use, a cytotoxic/cytostatic agent.

20. The composition of claim 19, wherein said cytotoxic/cytostatic agent is coupled chemically to said antibody, or said divalent functional fragment or derivative thereof for simultaneous use.

21. The composition of claim 20, wherein said cytotoxic/cytostatic agent is selected from the group consisting of alkylating agents, anti-metabolites, anti-tumor antibiotics, mitotic inhibitors, chromatin function inhibitors, anti-angiogenesis agents, anti-oestrogenic agents, anti-androgen agents and immunomodulators.

22. The composition of claim 21, wherein said cytotoxic/cytostatic agent is a mitotic inhibitor.

23. The composition of claim 18, wherein one, at least, of said antibodies, or a divalent functional fragment or derivative thereof, is conjugated with a cell toxin and/or a radioelement.

24. The composition of claim 23, wherein said toxin and/or a radioelement is coupled chemically to at least one of the elements of said composition for simultaneous use.

25. The composition as claimed in one of claims 17 to 24 as a medicament.

26. The use of an antibody, or a divalent functional fragment or derivative thereof, of claims 1 to 8 or 15 or of a composition of claims 17 to 24 for the preparation of a medicament intended to inhibit the growth and/or the proliferation of tumor cells.

27. The use of an antibody, or a divalent functional fragment or derivative thereof, of claims 1 to 8 or 15 or of a composition of claims 17 to 24 for the preparation of a medicament intended for the prevention or for the treatment of cancer.

28. The use of claim 27, wherein said cancer is prostate cancer, osteosarcomas, lung cancer, breast cancer, endometrial cancer, glioblastoma or colon cancer.

29. The use of claim 27 or 28, wherein said cancer is a HGF dependant and independent Met-activation related cancer.

30. A method of in vitro diagnosis of illnesses induced by an overexpression or an underexpression of the c-Met receptor starting from a biological sample in which the abnormal presence of c-Met receptor is suspected, wherein said method comprises a step wherein said biological sample is contacted with an antibody of claims 1 to 8 or 15.

31. The method of claim 30 wherein the said antibody is labeled.

32. The monoclonal antibody, or a divalent functional fragment or derivative thereof of claim 1, as herein described with reference to the Examples and/or Figures.




FIGURE 1




FIGURE 2A




FIGURE 2B




FIGURE 3A




FIGURE 3B




FIGURE 4A




FIGURE 4B




FIGURE 5A




FIGURE 5B



6/35




FIGURE 6A




FIGURE 6B




FIGURE 7A




FIGURE 7B




FIGURE 8A




FIGURE 8B




FIGURE 9A




FIGURE 9B




FIGURE 10A




FIGURE 10B





FIGURE 11




FIGURE 12




FIGURE 13




FIGURE 14A






FIGURE 14B




FIGURE 15




FIGURE 16




FIGURE 17A




FIGURE 17B




FIGURE 18




FIGURE 19




FIGURE 20



ELISA




FIGURE 21




FIGURE 22




FIGURE 23




FIGURE 24




FIGURE 25




FIGURE 26




FIGURE 27




FIGURE 28




FIGURE 29






FIGURE 30




FIGURE 31




FIGURE 32




FIGURE 33




FIGURE 34




FIGURE 35

SEQUENCE LISTING

<110> PIERRE FABRE MEDICAMENT

Goetsch, Liliane

Wurch, Thierry

Bes, Cédric

<120> Novel anti-cMet antibody

<130> D27117

<150> IB2008/055663

<151> 2008-12-02

<150> US 61/184,502

<151> 2009-06-05

<160> 117

<170> PatentIn version 3.3

<210> 1

<211> 8

<212> PRT

<213> mus musculus

<400> 1

Gly Tyr Ile Phe Thr Ala Tyr Thr

1 5

<210> 2

<211> 8

<212> PRT

<213> mus musculus

<400> 2

Ile Lys Pro Asn Asn Gly Leu Ala

1 5

<210> 3

<211> 11

<212> PRT

<213> mus musculus

<400> 3

Ala Arg Ser Glu Ile Thr Thr Glu Phe Asp Tyr

1 5 10

<210> 4  
<211> 118  
<212> PRT  
<213> mus musculus

<400> 4

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala  
1            5            10            15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ile Phe Thr Ala Tyr  
          20            25            30

Thr Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met  
          35            40            45

Gly Trp Ile Lys Pro Asn Asn Gly Leu Ala Asn Tyr Ala Gln Lys Phe  
          50            55            60

Gln Gly Arg Val Thr Met Thr Arg Asp Thr Ser Ile Ser Thr Ala Tyr  
          65            70            75            80

Met Glu Leu Ser Arg Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys  
          85            90            95

Ala Arg Ser Glu Ile Thr Thr Glu Phe Asp Tyr Trp Gly Gln Gly Thr  
          100            105            110

Leu Val Thr Val Ser Ser  
          115

<210> 5  
<211> 10  
<212> PRT  
<213> mus musculus

<400> 5

Glu Ser Val Asp Ser Tyr Ala Asn Ser Phe  
1            5            10

<210> 6  
<211> 3  
<212> PRT



<213> mus musculus

<400> 6

Arg Ala Ser

1

<210> 7

<211> 9

<212> PRT

<213> mus musculus

<400> 7

Gln Gln Ser Lys Glu Asp Pro Leu Thr

1

5

<210> 8

<211> 112

<212> PRT

<213> mus musculus

<400> 8

Asp Ile Val Leu Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly

1

5

10

15

Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Glu Ser Val Asp Ser Tyr

20

25

30

Ala Asn Ser Phe Met His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro

35

40

45

Lys Leu Leu Ile Tyr Arg Ala Ser Thr Arg Glu Ser Gly Val Pro Asp

50

55

60

Arg Phe Ser Gly Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Ser

65

70

75

80

Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln Ser Lys

85

90

95

Glu Asp Pro Leu Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg

100

105

110

<210> 9  
<211> 112  
<212> PRT  
<213> mus musculus

<400> 9

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly  
1           5           10           15

Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Glu Ser Val Asp Ser Tyr  
20           25           30

Ala Asn Ser Phe Met His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro  
35           40           45

Lys Leu Leu Ile Tyr Arg Ala Ser Thr Arg Glu Ser Gly Val Pro Asp  
50           55           60

Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser  
65           70           75           80

Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln Ser Lys  
85           90           95

Glu Asp Pro Leu Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg  
100           105           110

<210> 10  
<211> 112  
<212> PRT  
<213> mus musculus

<400> 10

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly  
1           5           10           15

Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Glu Ser Val Asp Ser Tyr  
20           25           30

Ala Asn Ser Phe Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro  
35           40           45

Lys Leu Leu Ile Tyr Arg Ala Ser Thr Arg Glu Ser Gly Val Pro Asp  
50 55 60

Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser  
65 70 75 80

Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln Ser Lys  
85 90 95

Glu Asp Pro Leu Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg  
100 105 110

<210> 11  
<211> 24  
<212> DNA  
<213> mus musculus

<400> 11  
ggctacatct tcacagcata cacc 24

<210> 12  
<211> 24  
<212> DNA  
<213> mus musculus

<400> 12  
attaaaccca acaatgggct ggcc 24

<210> 13  
<211> 33  
<212> DNA  
<213> mus musculus

<400> 13  
gccaggagcg aaattacaac agaattcgat tac 33

<210> 14  
<211> 354  
<212> DNA  
<213> mus musculus

<400> 14  
caggctccagc tgggtgcaatc cggcgcagag gtgaagaagc caggcgcttc cgtgaaggtg 60

agctgtaagg cctctggcta catcttcaca gcatacacca tgcactgggt gaggcaagct 120

cctgggcagg gactggagtg gatgggatgg attaaaccca acaatgggct ggccaactac 180

gcccagaaat tccagggtag ggtcactatg acaaggata ccagcatcag caccgcatat 240

atggagctga gcaggctgag gtctgacgac actgctgtct attattgcgc caggagcga 300

attacaacag aattcgatta ctgggggcag ggcaccctgg tgaccgtgc ctct 354

<210> 15

<211> 30

<212> DNA

<213> mus musculus

<400> 15

gaatctgtgg actcttacgc aaacagcttt 30

<210> 16

<211> 9

<212> DNA

<213> mus musculus

<400> 16

agggettct 9

<210> 17

<211> 27

<212> DNA

<213> mus musculus

<400> 17

cagcagtcca aggaggacc cctgact 27

<210> 18

<211> 333

<212> DNA

<213> lus musculus

<400> 18

gacattgtgc tgaccagtc tccgatagc ctggccgtgt ccctgggcga gagggtacc 60

atcaactgta aaagctccga atctgtggac tcttacgcaa acagctttat gactggtat 120

cagcaaaagc caggccaacc tccaaagctg ctgatttaca gggcttctac caggagagc 180

ggcgtgcccc ataggttcag cggatctggc agcaggaccg actttacact gaccatctcc 240

agcctgcagg ccgaagatgt ggcagtctat tactgccage agtccaagga ggacccctg 300

acttccgggg gtggtactaa agtggagatc aag 333

<210> 19  
<211> 333  
<212> DNA  
<213> mus musculus

<400> 19  
gacattgtga tgaccagtc tccgatage ctggccgtgt ccctgggcga gagggctacc 60  
atcaactgta aaagctccga atctgtggac tcttacgcaa acagctttat gactggtat 120  
cagcaaaagc caggccaacc tccaaagctg ctgatttaca gggcttctac caggagagc 180  
ggcgtgcccc ataggttcag cggatctggc agcggcaccg actttacact gaccatctcc 240  
agcctgcagg ccgaagatgt ggcagtctat tactgccagc agtccaagga ggacccctg 300  
acttccgggg gtggtactaa agtggagatc aag 333

<210> 20  
<211> 333  
<212> DNA  
<213> mus musculus

<400> 20  
gacattgtga tgaccagtc tccgatage ctggccgtgt ccctgggcga gagggctacc 60  
atcaactgta aaagctccga atctgtggac tcttacgcaa acagctttct gactggtat 120  
cagcaaaagc caggccaacc tccaaagctg ctgatttaca gggcttctac caggagagc 180  
ggcgtgcccc ataggttcag cggatctggc agcggcaccg actttacact gaccatctcc 240  
agcctgcagg ccgaagatgt ggcagtctat tactgccagc agtccaagga ggacccctg 300  
acttccgggg gtggtactaa agtggagatc aag 333

<210> 21  
<211> 13  
<212> PRT  
<213> artificial sequence

<220>  
<223> consensus artificial hinge proteic sequence

<220>  
<221> MUTAGEN  
<222> (1)..(1)  
<223> Xaa is P or R

<220>  
<221> MUTAGEN

<222> (2)..(2)  
<223> Xaa is K or R

<220>  
<221> MUTAGEN  
<222> (3)..(3)  
<223> Xaa is S, D or nothing

<220>  
<221> MUTAGEN  
<222> (5)..(5)  
<223> Xaa is G, D or nothing

<220>  
<221> MUTAGEN  
<222> (7)..(7)  
<223> Xaa is K, H or V

<220>  
<221> MUTAGEN  
<222> (8)..(8)  
<223> Xaa is E, P, T or nothing

<220>  
<221> MUTAGEN  
<222> (10)..(10)  
<223> Xaa is P or I

<220>  
<221> MUTAGEN  
<222> (11)..(11)  
<223> Xaa is P or nothing

<220>  
<221> MUTAGEN  
<222> (13)..(13)  
<223> Xaa is P or T

<400> 21

Xaa Xaa Xaa Cys Xaa Cys Xaa Xaa Cys Xaa Xaa Cys Xaa  
1 5 10

<210> 22  
<211> 11  
<212> PRT  
<213> artificial sequence

<220>  
<223> artificial hinge derived from IgG2

<400> 22

Arg Lys Cys Cys Val Glu Cys Pro Pro Cys Pro  
1 5 10

<210> 23  
<211> 12  
<212> PRT  
<213> artificial sequence

<220>  
<223> artificial hinge derived from IgG1

<400> 23

Pro Arg Asp Cys Gly Cys Lys Pro Cys Ile Cys Thr  
1 5 10

<210> 24  
<211> 12  
<212> PRT  
<213> artificial sequence

<220>  
<223> artificial hinge derived from IgG1

<400> 24

Pro Lys Ser Cys Gly Cys Lys Pro Cys Ile Cys Thr  
1 5 10

<210> 25  
<211> 12  
<212> PRT  
<213> artificial sequence

<220>  
<223> artificial hinge derived from IgG1

<400> 25

Pro Lys Ser Cys Gly Cys Lys Pro Cys Ile Cys Pro  
1 5 10

<210> 26  
<211> 13  
<212> PRT  
<213> artificial sequence

<220>  
<223> artificial hinge derived from IgG1

<400> 26

Pro Arg Asp Cys Gly Cys Lys Pro Cys Pro Pro Cys Pro  
1           5           10

<210> 27

<211> 13

<212> PRT

<213> artificial sequence

<220>

<223> artificial hinge derived from IgG1

<400> 27

Pro Arg Asp Cys Gly Cys His Thr Cys Pro Pro Cys Pro  
1           5           10

<210> 28

<211> 12

<212> PRT

<213> artificial sequence

<220>

<223> artificial hinge derived from IgG1

<400> 28

Pro Lys Ser Cys Asp Cys His Cys Pro Pro Cys Pro  
1           5           10

<210> 29

<211> 33

<212> DNA

<213> artificial sequence

<220>

<223> artificial hinge derived from IgG2

<400> 29

aggaagtgct gtgtggagtg cccccctgc cca

33

<210> 30

<211> 36

<212> DNA

<213> artificial sequence

<220>

<223> artificial sequence derived from IgG1



<400> 30  
ccccgggact gtgggtgcaa gccttgcaatt tgtacc 36

<210> 31  
<211> 36  
<212> DNA  
<213> artificial sequence

<220>  
<223> artificial hinge derived from IgG1

<400> 31  
cccaagagct gtgggtgcaa gccttgcaatt tgtacc 36

<210> 32  
<211> 36  
<212> DNA  
<213> artificial sequence

<220>  
<223> artificial hinge derived from IgG1

<400> 32  
ccaaagagct gcgggtgcaa gccttgatc tgtccc 36

<210> 33  
<211> 39  
<212> DNA  
<213> artificial sequence

<220>  
<223> artificial hinge derived from IgG1

<400> 33  
ccacgggact gtgggtgcaa gcctgcctt ccgtgtcca 39

<210> 34  
<211> 39  
<212> DNA  
<213> artificial sequence

<220>  
<223> artificial hinge derived from IgG1

<400> 34  
cccagagact gtgggtgtca cacctgcctt ccttgcct 39

<210> 35  
<211> 36

<212> DNA  
<213> artificial sequence

<220>  
<223> artificial hinge derived from IgG1

<400> 35  
ccccaaaagct gcgattgccca ctgtcctcca tgtcca 36

<210> 36  
<211> 443  
<212> PRT  
<213> mus musculus

<400> 36

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala  
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ile Phe Thr Ala Tyr  
20 25 30

Thr Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met  
35 40 45

Gly Trp Ile Lys Pro Asn Asn Gly Leu Ala Asn Tyr Ala Gln Lys Phe  
50 55 60

Gln Gly Arg Val Thr Met Thr Arg Asp Thr Ser Ile Ser Thr Ala Tyr  
65 70 75 80

Met Glu Leu Ser Arg Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys  
85 90 95

Ala Arg Ser Glu Ile Thr Thr Glu Phe Asp Tyr Trp Gly Gln Gly Thr  
100 105 110

Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro  
115 120 125

Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly  
130 135 140

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn

145            150            155            160  
 Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln  
           165            170            175  
 Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser  
           180            185            190  
 Asn Phe Gly Thr Gln Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser  
           195            200            205  
 Asn Thr Lys Val Asp Lys Thr Val Glu Arg Lys Cys Cys Val Glu Cys  
           210            215            220  
 Pro Pro Cys Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe  
           225            230            235            240  
 Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val  
           245            250            255  
 Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Gln Phe  
           260            265            270  
 Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro  
           275            280            285  
 Arg Glu Glu Gln Phe Asn Ser Thr Phe Arg Val Val Ser Val Leu Thr  
           290            295            300  
 Val Val His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val  
           305            310            315            320  
 Ser Asn Lys Gly Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr  
           325            330            335  
 Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg  
           340            345            350  
 Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly  
           355            360            365

Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro  
370 375 380

Glu Asn Asn Tyr Lys Thr Thr Pro Pro Met Leu Asp Ser Asp Gly Ser  
385 390 395 400

Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln  
405 410 415

Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His  
420 425 430

Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly  
435 440

<210> 37

<211> 445

<212> PRT

<213> mus musculus

<400> 37

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala  
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ile Phe Thr Ala Tyr  
20 25 30

Thr Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met  
35 40 45

Gly Trp Ile Lys Pro Asn Asn Gly Leu Ala Asn Tyr Ala Gln Lys Phe  
50 55 60

Gln Gly Arg Val Thr Met Thr Arg Asp Thr Ser Ile Ser Thr Ala Tyr  
65 70 75 80

Met Glu Leu Ser Arg Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys  
85 90 95

Ala Arg Ser Glu Ile Thr Thr Glu Phe Asp Tyr Trp Gly Gln Gly Thr  
100 105 110

Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro  
115 120 125

Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly  
130 135 140

Cys Leu Val Lys Asp Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser  
145 150 155 160

Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser  
165 170 175

Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser  
180 185 190

Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn  
195 200 205

Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Cys His Cys  
210 215 220

Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu  
225 230 235 240

Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu  
245 250 255

Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys  
260 265 270

Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys  
275 280 285

Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu  
290 295 300

Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys  
305 310 315 320

Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys  
325 330 335

Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser  
340 345 350

Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys  
355 360 365

Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln  
370 375 380

Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly  
385 390 395 400

Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln  
405 410 415

Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn  
420 425 430

His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
435 440 445

<210> 38

<211> 218

<212> PRT

<213> mus musculus

<400> 38

Asp Ile Val Leu Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly  
1 5 10 15

Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Glu Ser Val Asp Ser Tyr  
20 25 30

Ala Asn Ser Phe Met His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro  
35 40 45

Lys Leu Leu Ile Tyr Arg Ala Ser Thr Arg Glu Ser Gly Val Pro Asp  
50 55 60

Arg Phe Ser Gly Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Ser  
65 70 75 80

Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln Ser Lys  
85 90 95

Glu Asp Pro Leu Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg  
100 105 110

Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln  
115 120 125

Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr  
130 135 140

Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser  
145 150 155 160

Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr  
165 170 175

Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys  
180 185 190

His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro  
195 200 205

Val Thr Lys Ser Phe Asn Arg Gly Glu Cys  
210 215

<210> 39  
<211> 218  
<212> PRT  
<213> mus musculus

<400> 39

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly  
1 5 10 15

Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Glu Ser Val Asp Ser Tyr  
20 25 30

Ala Asn Ser Phe Met His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro  
35 40 45

Lys Leu Leu Ile Tyr Arg Ala Ser Thr Arg Glu Ser Gly Val Pro Asp  
50 55 60

Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser  
65 70 75 80

Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln Ser Lys  
85 90 95

Glu Asp Pro Leu Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg  
100 105 110

Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln  
115 120 125

Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr  
130 135 140

Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser  
145 150 155 160

Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr  
165 170 175

Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys  
180 185 190

His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro  
195 200 205

Val Thr Lys Ser Phe Asn Arg Gly Glu Cys  
210 215

<210> 40

<211> 218

<212> PRT

<213> mus musculus



<400> 40

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly  
1 5 10 15

Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Glu Ser Val Asp Ser Tyr  
20 25 30

Ala Asn Ser Phe Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro  
35 40 45

Lys Leu Leu Ile Tyr Arg Ala Ser Thr Arg Glu Ser Gly Val Pro Asp  
50 55 60

Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser  
65 70 75 80

Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln Ser Lys  
85 90 95

Glu Asp Pro Leu Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg  
100 105 110

Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln  
115 120 125

Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr  
130 135 140

Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser  
145 150 155 160

Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr  
165 170 175

Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys  
180 185 190

His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro  
195 200 205

Val Thr Lys Ser Phe Asn Arg Gly Glu Cys  
210 215

<210> 41  
<211> 1332  
<212> DNA  
<213> mus musculus

<400> 41  
caggtccage tggtgcaatc cggcgcagag gtgaagaagc caggcgcttc cgtgaagggtg 60  
agctgtaagg cctctggcta catcttcaca gcatacacca tgcactgggt gaggcaagct 120  
cctgggcagg gactggagtg gatgggatgg attaaacca acaatgggct ggccaactac 180  
gcccagaaat tccagggtag ggtcactatg acaagggata ccagcatcag caccgcatat 240  
atggagctga gcaggctgag gtctgacgac actgtgtct attattgcgc caggagcgaa 300  
attacaacag aattcgatta ctgggggcag ggcaccctgg tgaccgtgtc ctctgccage 360  
accaagggcc caagcgtgtt cccgetagec cctgtctcca gaagcaccag cgagagcaca 420  
gccgcctgg gctgectggt gaaggactac ttccccgagc cctgaccgt gtcctggaac 480  
agcggagccc tgaccagcgg cgtgcacacc ttccccgccg tgctgcagag cagcggcctg 540  
tacagcctga gcagcgtggt gaccgtgcca agcagcaact tggcaccca gacctacacc 600  
tgtaacgtgg accacaagcc cagcaacacc aaggtggaca agaccgtgga gaggaagtgc 660  
tgtgtggagt gccccctg cccagcccc ccagtggccg gaccagcgt gttctgttc 720  
cccccaagc ccaaggacac cctgatgatc agcagaacc cagaggtgac ctgtgtggtg 780  
gtggacgtgt cccacgagga ccccagggtg cagttcaact ggtacgtgga cggcgtggag 840  
gtgcacaacg ccaagaccaa gccagagag gagcagttta acagcacctt ccgggtggtg 900  
tccgtgctga ccgtggtgca ccaggactgg ctgaacggca aggagtacaa gtgtaaggtc 960  
tccaacaagg gcctgccage ccccatcgaa aagaccatca gcaagaccaa gggacagcca 1020  
agagagccac aggtctacac cctgcccccc agcagggagg agatgaccaa gaaccaggtg 1080  
tccctgacct gtctggtgaa gggcttctac ccaagcgaca tcgccgtgga gtgggagagc 1140  
aacggccagc ccgagaacaa ctacaagacc acccccccaa tgctggacag cgacggcagc 1200  
ttcttctgt acagcaagct gacagtggac aagagcagat ggcagcaggg caactgttc 1260  
agctgctccg tgatgcacga ggcctgcac aaccactaca cccagaagag cctgagcctg 1320

tccccaggct ga

1332

<210> 42

<211> 1341

<212> DNA

<213> mus musculus

<400> 42

caggctccagc tgggtgcaatc cggcgcagag gtgaagaagc caggcgcttc cgtgaagggtg 60  
agctgtaagg cctctggcta catcttcaca gcatacacca tgcactgggt gaggcaagct 120  
cctgggcagg gactggagtg gatgggatgg attaaacca acaatgggct ggccaactac 180  
gcccagaaat tccagggtag ggtcactatg acaagggata ccagcatcag caccgcatat 240  
atggagctga gcaggctgag gtctgacgac actgctgtct attattgcgc caggagcgaa 300  
attacaacag aattcgatta ctgggggcag ggcaccctgg tgaccgtgtc ctctgccagc 360  
accaagggcc caagcgtgtt cccgctagcc ccgtctcaa agagtacctc aggcggaact 420  
gcccgtcttg gttgccttgt gaaggactat ttctctgagc ccgtgacggt cagetggaac 480  
tctggcgcac tgactagcgg cgtacacaca ttccctgcgg tgcctcaaag ttccgggctg 540  
tactcactgt cctccgtgtg gaccgttcca tctagttccc tcgggacaca gacatacatc 600  
tgtaatgtga atcataagcc ttcaaacacc aaggctgaca aacgggtcga gcccanaaage 660  
tgcgaltgcc actgtcctec atgtccagcc cccgaactgt tgggcggacc gagegtgttc 720  
ttgtttccac ccaagcccaa agataccctc atgatcagca gaacccccga ggtgacctgt 780  
gtggtggtgg acgtgtccca cgaggacca gaggtgaagt tcaactggtc cgtggacggc 840  
gtggaggtgc acaacgcaa gaccaagccc agagaggagc agtacaacag cacctacagg 900  
gtggtgtccg tgctgacctg gctgcaccag gactggctga acggcaagga gtacaagtgt 960  
aaggtgtcca acaaggccct gccagccca atcgaaaaga ccatcagcaa ggccaagggc 1020  
cagccaagag agccccaggt gtacaccctg ccaccagca gggaggagat gaccaagaac 1080  
caggtgtccc tgacctgtct ggtgaagggc ttctaccaa gcgacatgc cgtggagtgg 1140  
gagageaacg gccagcccga gaacaactac aagaccacc cccagtgtct ggacagcgac 1200  
ggcagcttct tctgtacag caagctgacc gtggacaaga gcagatggca gcagggaac 1260  
gtgttcagct gctccgtgat gcacgaggcc ctgcacaacc actacacca gaagagcctg 1320

agcctgtccc caggcaagtg a

1341

<210> 43

<211> 657

<212> DNA

<213> mus musculus

<400> 43

gacattgtgc tgaccagtc tccgatagc ctggccgtgt ccctgggcga gagggtacc 60

atcaactgta aaagctccga atctgtggac tcttacgcaa acagctttat gactggtat 120

cagcaaaagc caggccaacc tccaaagctg ctgatttaca gggcttctac caggagagc 180

ggcgtgcccc ataggttcag cggatctggc agcaggaccg actttacact gaccatctcc 240

agcctgcagg ccgaagatgt ggcagtctat tactgccagc agtccaagga ggacccctg 300

actttcgggg gtggtactaa agtggagatc aagcgtacgg tggccgctcc cagcgtgttc 360

atcttcccc caagcgacga gcagctgaag agcggcaccg ccagcgtggt gtgtctgctg 420

aacaacttct accccaggga ggccaaggtg cagtggaagg tggacaacgc cctgcagagc 480

ggcaacagcc aggagagcgt caccgagcag gacagcaagg actccaccta cagcctgagc 540

agcacctga cctgagcaa ggccgactac gagaagcaca aggtgtacgc ctgtgaggtg 600

accaccagg gcctgtccag cccctgacc aagagcttca acaggggcga gtgtga 657

<210> 44

<211> 657

<212> DNA

<213> mus musculus

<400> 44

gacattgtga tgaccagtc tccgatagc ctggccgtgt ccctgggcga gagggtacc 60

atcaactgta aaagctccga atctgtggac tcttacgcaa acagctttat gactggtat 120

cagcaaaagc caggccaacc tccaaagctg ctgatttaca gggcttctac caggagagc 180

ggcgtgcccc ataggttcag cggatctggc agcggcaccg actttacact gaccatctcc 240

agcctgcagg ccgaagatgt ggcagtctat tactgccagc agtccaagga ggacccctg 300

actttcgggg gtggtactaa agtggagatc aagcgtacgg tggccgctcc cagcgtgttc 360

atcttcccc caagcgacga gcagctgaag agcggcaccg ccagcgtggt gtgtctgctg 420

aacaacttct accccaggga ggccaaggtg cagtggaagg tggacaacgc cctgcagagc 480

ggcaacagcc aggagagcgt caccgagcag gacagcaagg actccaccta cagcctgagc 540  
agcacctga cctgagcaa ggccgactac gagaagcaca aggtgtacgc ctgtgaggtg 600  
accaccagg gcctgtccag ccccgtgacc aagagettca acaggggcca gtgctga 657

<210> 45  
<211> 657  
<212> DNA  
<213> mus musculus

<400> 45  
gacattgtga tgaccagtc tccgatagc ctggccgtgt ccctgggcca gagggctacc 60  
atcaactgta aaagctcca atctgtggac tcttacgcaa acagtttct gcaactgtat 120  
cagcaaaagc caggccaacc tccaaagctg ctgatttaca gggcttctac caggagagc 180  
ggcgtgccc ataggttcag cggatctggc agcggcaccg actttacact gaccatctcc 240  
agcctgcagg ccgaagatgt ggcagtctat tactgccagc agtccaagga ggacccctg 300  
actttcgggg gtggtactaa agtggagatc aagcgtacgg tggccgctcc cagcgtgttc 360  
atcttcccc caagcgacga gcagctgaag agcggcaccg ccagcgtggt gtgtctgctg 420  
aacaacttct acccaggga ggccaaggtg cagtggaagg tggacaacgc cctgcagagc 480  
ggcaacagcc aggagagcgt caccgagcag gacagcaagg actccaccta cagcctgagc 540  
agcacctga cctgagcaa ggccgactac gagaagcaca aggtgtacgc ctgtgaggtg 600  
accaccagg gcctgtccag ccccgtgacc aagagettca acaggggcca gtgctga 657

<210> 46  
<211> 118  
<212> PRT  
<213> mus musculus

<400> 46

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala  
1 5 10 15

Ser Val Lys Ile Ser Cys Lys Thr Ser Gly Tyr Ile Phe Thr Ala Tyr  
20 25 30

Thr Met His Trp Val Arg Gln Ser Leu Gly Glu Ser Leu Asp Trp Ile  
35 40 45

Gly Gly Ile Lys Pro Asn Asn Gly Leu Ala Asn Tyr Asn Gln Lys Phe  
50 55 60

Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr  
65 70 75 80

Met Asp Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys  
85 90 95

Ala Arg Ser Glu Ile Thr Thr Glu Phe Asp Tyr Trp Gly Gln Gly Thr  
100 105 110

Ala Leu Thr Val Ser Ser  
115

<210> 47

<211> 112

<212> PRT

<213> mus musculus

<400> 47

Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly  
1 5 10 15

Gln Arg Ala Thr Ile Ser Cys Arg Ala Ser Glu Ser Val Asp Ser Tyr  
20 25 30

Ala Asn Ser Phe Met His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro  
35 40 45

Lys Leu Leu Ile Tyr Arg Ala Ser Asn Leu Glu Ser Gly Ile Pro Ala  
50 55 60

Arg Phe Ser Gly Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Asn  
65 70 75 80

Pro Val Glu Ala Asp Asp Val Ala Thr Tyr Tyr Cys Gln Gln Ser Lys  
85 90 95

Glu Asp Pro Leu Thr Phe Gly Ser Gly Thr Lys Leu Glu Met Lys Arg  
100 105 110

<210> 48  
<211> 354  
<212> DNA  
<213> mus musculus

<400> 48  
gaggtccagc tgcagcagag cgggccagaa ctggttaaac ctggcggcag cgtgaagatt 60  
agctgtaaga ccagcgggta catctttaca gcatatacca tgcactgggt gaggcagagt 120  
ctggggggaat ctctggactg gatcggaggt attaagccca acaatggcct ggetaactat 180  
aatcaaaaat tcaagggcaa agccacactg accgtcgata agtctcttc cacagcttac 240  
atggatctga gaagcctgac atccgaggac agtgcagtgt actactgcgc ccggtctgag 300  
atcaactaccg agttegacta ttggggacag ggcactgcac tgaccgtctc ctcc 354

<210> 49  
<211> 333  
<212> DNA  
<213> mus musculus

<400> 49  
gatattgtgc tgacacagtc tcccgcaage ctcgctgta gcttgggtca acgggccaca 60  
attagttgtc gcgctctga atccgtcgac tettatgcta acagctttat gcaactggat 120  
caacagaagc ccgggcagcc acctaaactg ctgatctaca gggccagcaa tctggagagt 180  
ggcatcccag ctagatttag cgggtccggg tccaggaccg acttactct gaccatcaac 240  
cccgtggagg cagacgacgt ggccacttac tactgccagc agtctaaaga ggatcccctc 300  
acattcggct ccggaacaaa gctggaaatg aag 333

<210> 50  
<211> 443  
<212> PRT  
<213> mus musculus

<400> 50

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala  
1 5 10 15

Ser Val Lys Ile Ser Cys Lys Thr Ser Gly Tyr Ile Phe Thr Ala Tyr  
20 25 30

Thr Met His Trp Val Arg Gln Ser Leu Gly Glu Ser Leu Asp Trp Ile  
35 40 45

Gly Gly Ile Lys Pro Asn Asn Gly Leu Ala Asn Tyr Asn Gln Lys Phe  
50 55 60

Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr  
65 70 75 80

Met Asp Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys  
85 90 95

Ala Arg Ser Glu Ile Thr Thr Glu Phe Asp Tyr Trp Gly Gln Gly Thr  
100 105 110

Ala Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro  
115 120 125

Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly  
130 135 140

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn  
145 150 155 160

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln  
165 170 175

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser  
180 185 190

Asn Phe Gly Thr Gln Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser  
195 200 205

Asn Thr Lys Val Asp Lys Thr Val Glu Arg Lys Cys Cys Val Glu Cys  
210 215 220

Pro Pro Cys Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe  
225 230 235 240

Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val



245            250            255

Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Gln Phe  
260            265            270

Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro  
275            280            285

Arg Glu Glu Gln Phe Asn Ser Thr Phe Arg Val Val Ser Val Leu Thr  
290            295            300

Val Val His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val  
305            310            315            320

Ser Asn Lys Gly Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr  
325            330            335

Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg  
340            345            350

Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly  
355            360            365

Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro  
370            375            380

Glu Asn Asn Tyr Lys Thr Thr Pro Pro Met Leu Asp Ser Asp Gly Ser  
385            390            395            400

Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln  
405            410            415

Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His  
420            425            430

Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly  
435            440

<210> 51  
<211> 446  
<212> PRT

<213> mus musculus

<400> 51

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala  
1 5 10 15

Ser Val Lys Ile Ser Cys Lys Thr Ser Gly Tyr Ile Phe Thr Ala Tyr  
20 25 30

Thr Met His Trp Val Arg Gln Ser Leu Gly Glu Ser Leu Asp Trp Ile  
35 40 45

Gly Gly Ile Lys Pro Asn Asn Gly Leu Ala Asn Tyr Asn Gln Lys Phe  
50 55 60

Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr  
65 70 75 80

Met Asp Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys  
85 90 95

Ala Arg Ser Glu Ile Thr Thr Glu Phe Asp Tyr Trp Gly Gln Gly Thr  
100 105 110

Ala Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro  
115 120 125

Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly  
130 135 140

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn  
145 150 155 160

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln  
165 170 175

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser  
180 185 190

Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser  
195 200 205

Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Cys His  
210 215 220

Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe  
225 230 235 240

Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro  
245 250 255

Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val  
260 265 270

Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr  
275 280 285

Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val  
290 295 300

Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys  
305 310 315 320

Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser  
325 330 335

Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro  
340 345 350

Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val  
355 360 365

Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly  
370 375 380

Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp  
385 390 395 400

Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp  
405 410 415

Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His  
420 425 430

Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
435 440 445

<210> 52  
<211> 218  
<212> PRT  
<213> mus musculus

<400> 52

Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly  
1 5 10 15

Gln Arg Ala Thr Ile Ser Cys Arg Ala Ser Glu Ser Val Asp Ser Tyr  
20 25 30

Ala Asn Ser Phe Met His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro  
35 40 45

Lys Leu Leu Ile Tyr Arg Ala Ser Asn Leu Glu Ser Gly Ile Pro Ala  
50 55 60

Arg Phe Ser Gly Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Asn  
65 70 75 80

Pro Val Glu Ala Asp Asp Val Ala Thr Tyr Tyr Cys Gln Gln Ser Lys  
85 90 95

Glu Asp Pro Leu Thr Phe Gly Ser Gly Thr Lys Leu Glu Met Lys Arg  
100 105 110

Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln  
115 120 125

Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr  
130 135 140

Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser  
145 150 155 160

Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr  
165 170 175

Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys  
180 185 190

His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro  
195 200 205

Val Thr Lys Ser Phe Asn Arg Gly Glu Cys  
210 215

<210> 53  
<211> 1332  
<212> DNA  
<213> mus musculus

<400> 53  
gagggtccagc tgcagcagag cgggccagaa ctggttaaac ctggcgccag cgtgaagatt 60  
agctgtaaga ccagcgggta catctttaca gcatatacca tgcactgggt gaggcagagt 120  
ctggggggaat ctctggactg gatcggaggt attaagccca acaatggcct ggetaactat 180  
aatcaaaaat tcaagggeaa agccacactg accgtcgata agtctcttc cacagcttac 240  
atggatctga gaagcctgac atccgaggac agtgcagtgt actactgcgc ccggtctgag 300  
atactaccg agttcgacta ttggggacag ggcactgcac tgaccgtctc ctccgccage 360  
accaagggcc caagcgtgtt cccgctagcg cctgtctcca gaagcaccag cgagagcaca 420  
gccgcctgg gctgectggt gaaggactac ttccccgagc ccgtgaccgt gtctctggaac 480  
agcggagccc tgaccagegg cgtgcacacc ttccccgccc tgctgcagag cageggcctg 540  
tacagcctga gcagcgtggt gaccgtgcca agcagcaact tcggcaccca gacctacacc 600  
tgtaacgtgg accacaagcc cagcaacacc aaggtggaca agaccgtgga gaggaagtgc 660  
tgtgtggagt gccccctg cccagccccc ccagtggccc gaccagegt gttcctgttc 720  
cccccaage ccaaggacac cctgatgac agcagaacct ccgaggtgac ctgtgtggtg 780  
gtggacgtgt cccacgagga ccccaggtg cagtcaact ggtacgtgga cggcgtggag 840  
gtgcacaacg ccaagaccaa gccagagag gagcagtta acagcacctt ccgggtggtg 900  
tccgtctga ccgtggtgca ccaggactgg ctgaacggca aggagtataa gtgtaaggtc 960

tccaacaagg gcctgccagc ccccatcgaa aagaccatca gcaagaccaa gggacagcca 1020  
agagagccac aggtctacac cctgcccccc agcaggagg agatgaccaa gaaccaggtg 1080  
tccctgacct gtctggtgaa gggcttctac ccaagegaca tcgccgtgga gtgggagagc 1140  
aacggccagc ccgagaacaa ctacaagacc acccccccaa tgctggacag cgacggcagc 1200  
ttcttctgt acagcaagct gacagtggac aagagcagat ggcagcaggg caactgttc 1260  
agctgtccg tgatgcacga ggccctgcac aaccactaca cccagaagag cctgagcctg 1320  
tcccaggt ga 1332

<210> 54  
<211> 1341  
<212> DNA  
<213> mus musculus

<400> 54  
gaggtccagc tgcagcagag cgggccagaa ctggttaaac ctggcggcag cgtgaagatt 60  
agctgtaaga ccagcggfta catctttaca geatatacca tgcaactgggt gaggcagagt 120  
ctgggggaat ctctggactg gatcggaggt attaagccca acaatggcct ggctaactat 180  
aatcaaaaat tcaagggcaa agccacactg accgtcgata agtcctcttc cacagcttac 240  
atggatctga gaagcctgac atccgaggac agtgcagtgt actactgcgc ccggtctgag 300  
atactaccg agttcgacta ttggggacag ggcactgcac tgaccgtctc ctccgccagc 360  
accaagggcc caagegtgtt cccgctagcc ccgtcttcaa agagtacctc aggcggaact 420  
gccgctcttg gttgccttgt gaaggactat tttctgagc ccgtgacggt cagctggaac 480  
tctggcgcac tgactagegg cgtacacaca ttcctgcgg tgctccaaag ttccgggctg 540  
tactcaactg cctccgttgt gaccgttcca tctagttccc tcgggacaca gacatacatc 600  
tgtaatgtga atcataagcc ttcaaacacc aaggtcgaca aacgggtcga gcccaaaagc 660  
tgcgattgcc actgtctctc atgtccagcc cccgaactgt tgggcggacc gagegtgttc 720  
ttgtttccac ccaagcccaa agataccctc atgatcagca gaacccccga ggtgacctgt 780  
gtggtggtgg acgtgtccca cgaggacca gaggtgaagt tcaactggta cgtggacggc 840  
gtggaggtgc acaacgcaa gaccaagccc agagaggagc agtacaacag cacctacagg 900  
gtggtgtccg tgctgaccgt gctgcaccag gactggctga acggcaagga gtacaagtgt 960

aaggtgtcca acaaggccct gccagcccca atcgaaaaga ccatcagcaa ggccaagggc 1020  
cagccaagag agccccaggt gtacaccctg ccaccagca gggaggagat gaccaagaac 1080  
caggtgtccc tgacctgtct ggtgaagggc ttctaccaa gcgacatgc cgtggagtgg 1140  
gagageaacg gccagcccga gaacaactac aagaccacc cccagtgtt ggacagcgac 1200  
ggcagcttct tctgtacag caagctgacc gtggacaaga gcagatggca gcagggcaac 1260  
gtgttcagct gtccgtgat gcagaggcc ctgcacaacc actacacca gaagagcctg 1320  
agcctgtccc caggcaagtg a 1341

<210> 55  
<211> 657  
<212> DNA  
<213> mus musculus

<400> 55  
gatattgtgc tgacacagtc tcccgcaagc ctcgctgtta gcttgggtca acgggccaca 60  
attagttgtc gcgectctga atccgtcgac tcttatgcta acagcttat gcaactggtat 120  
caacagaagc ccgggcagcc acctaaactg ctgatctaca gggccagcaa tctggagagt 180  
ggcateccag ctagatttag cggttccggg tccaggaccg acttcaactet gaccatcaac 240  
cccgtggagg cagacgacgt ggccacttac tactgccagc agtctaaaga ggatcccctc 300  
acattcggct ccggaacaaa gctggaaatg aagcgtacgg tggccgctcc cagcgtgttc 360  
atcttcccc caagcgacga gcagctgaag agcggcaccg ccagcgtggt gtgtctgctg 420  
aacaacttct accccaggga ggccaaggtg cagtggaagg tggacaacgc cctgcagagc 480  
ggcaacagcc aggagagcgt caccgagcag gacagcaagg actccaccta cagcctgagc 540  
agcacctga cctgagcaa ggccgactac gagaagcaca aggtgtacgc ctgtgaggtg 600  
accaccagg gcctgtccag cccctgacc aagagcttca acaggggcga gtgctga 657

<210> 56  
<211> 14  
<212> PRT  
<213> artificial

<220>  
<223> consensus artificial hinge proteic sequence

<220>

<221> mutagen  
<222> (1)..(1)  
<223> X is P, R, C or nothing

<220>  
<221> mutagen  
<222> (2)..(2)  
<223> X is K, C, R or nothing

<220>  
<221> mutagen  
<222> (3)..(3)  
<223> X is S, C, D or nothing

<220>  
<221> mutagen  
<222> (5)..(5)  
<223> X is D, C, G or nothing

<220>  
<221> mutagen  
<222> (6)..(6)  
<223> X is K, C or nothing

<220>  
<221> mutagen  
<222> (7)..(7)  
<223> X is T, C or nothing

<220>  
<221> mutagen  
<222> (8)..(8)  
<223> X is H, V, K or nothing

<220>  
<221> mutagen  
<222> (9)..(9)  
<223> X is T, C, E, P or nothing

<220>  
<221> mutagen  
<222> (11)..(11)  
<223> X is P or I

<220>  
<221> mutagen  
<222> (12)..(12)  
<223> X is P or nothing

<220>  
<221> mutagen  
<222> (14)..(14)  
<223> X is P or T



<400> 56

Xaa Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa Cys Xaa Xaa Cys Xaa  
1 5 10

<210> 57

<211> 14

<212> PRT

<213> artificial

<220>

<223> consensus artificial hinge proteic sequence

<220>

<221> mutagen

<222> (1)..(1)

<223> X is P, R, C or nothing

<220>

<221> mutagen

<222> (2)..(2)

<223> X is K, C, R or nothing

<220>

<221> mutagen

<222> (3)..(3)

<223> X is S, C, D or nothing

<220>

<221> mutagen

<222> (5)..(5)

<223> X is D, C, G or nothing

<220>

<221> mutagen

<222> (6)..(6)

<223> X is K, C or nothing

<220>

<221> mutagen

<222> (7)..(7)

<223> X is T, C or nothing

<220>

<221> mutagen

<222> (8)..(8)

<223> X is H, V, K or nothing

<220>

<221> mutagen

<222> (9)..(9)

<223> X is T, C, E, P or nothing

<400> 57

Xaa Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa Cys Pro Pro Cys Pro  
1 5 10

<210> 58

<211> 14

<212> PRT

<213> artificial

<220>

<223> artificial hinge region

<400> 58

Cys Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro  
1 5 10

<210> 59

<211> 14

<212> PRT

<213> artificial

<220>

<223> artificial hinge region

<400> 59

Pro Cys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro  
1 5 10

<210> 60

<211> 14

<212> PRT

<213> artificial

<220>

<223> artificial hinge region

<400> 60

Pro Lys Cys Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro  
1 5 10

<210> 61

<211> 14

<212> PRT

<213> artificial

<220>

<223> artificial hinge region

<400> 61

Pro Lys Ser Cys Cys Lys Thr His Thr Cys Pro Pro Cys Pro  
1 5 10

<210> 62

<211> 14

<212> PRT

<213> artificial

<220>

<223> artificial hinge region

<400> 62

Pro Lys Ser Cys Asp Cys Thr His Thr Cys Pro Pro Cys Pro  
1 5 10

<210> 63

<211> 14

<212> PRT

<213> artificial

<220>

<223> artificial hinge region

<400> 63

Pro Lys Ser Cys Asp Lys Cys His Thr Cys Pro Pro Cys Pro  
1 5 10

<210> 64

<211> 14

<212> PRT

<213> artificial

<220>

<223> artificial hinge region

<400> 64

Pro Lys Ser Cys Asp Lys Thr His Cys Cys Pro Pro Cys Pro  
1 5 10

<210> 65

<211> 12

<212> PRT

<213> artificial

<220>

<223> artificial hinge region

<400> 65

Lys Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro  
1 5 10

<210> 66

<211> 13

<212> PRT

<213> artificial

<220>

<223> artificial hinge region

<400> 66

Pro Lys Ser Cys Asp Cys His Thr Cys Pro Pro Cys Pro  
1 5 10

<210> 67

<211> 13

<212> PRT

<213> artificial

<220>

<223> artificial hinge region

<400> 67

Pro Lys Ser Cys Asp Cys Thr His Cys Pro Pro Cys Pro  
1 5 10

<210> 68

<211> 12

<212> PRT

<213> artificial

<220>

<223> artificial hinge region

<400> 68

Pro Cys Ser Cys Lys His Thr Cys Pro Pro Cys Pro  
1 5 10

<210> 69

<211> 12  
<212> PRT  
<213> artificial

<220>  
<223> artificial hinge region

<400> 69

Pro Ser Cys Cys Thr His Thr Cys Pro Pro Cys Pro  
1 5 10

<210> 70  
<211> 12  
<212> PRT  
<213> artificial

<220>  
<223> artificial hinge region

<400> 70

Pro Ser Cys Asp Lys His Cys Cys Pro Pro Cys Pro  
1 5 10

<210> 71  
<211> 10  
<212> PRT  
<213> artificial

<220>  
<223> artificial hinge region

<400> 71

Pro Lys Ser Cys Thr Cys Pro Pro Cys Pro  
1 5 10

<210> 72  
<211> 14  
<212> PRT  
<213> artificial

<220>  
<223> artificial hinge region

<400> 72

Pro Lys Ser Cys Asp Lys Cys Val Glu Cys Pro Pro Cys Pro  
1 5 10

<210> 73  
<211> 42  
<212> DNA  
<213> artificial

<220>  
<223> artificial hinge region

<400> 73  
tgcaagagct gcgacaagac ccacacctgt cccccctgcc ct 42

<210> 74  
<211> 42  
<212> DNA  
<213> artificial

<220>  
<223> artificial hinge region

<400> 74  
ccctgcagct gcgacaagac ccacacctgt cccccctgcc ct 42

<210> 75  
<211> 42  
<212> DNA  
<213> artificial

<220>  
<223> artificial hinge region

<400> 75  
cccaagtgct gcgacaagac ccacacctgt cccccctgcc ct 42

<210> 76  
<211> 42  
<212> DNA  
<213> artificial

<220>  
<223> artificial hinge region

<400> 76  
cctaagagct gttgcaagac ccacacctgt cccccctgcc ct 42

<210> 77  
<211> 42  
<212> DNA  
<213> artificial

<220>  
 <223> artificial hinge region

<400> 77  
 cccaagagct gcgactgcac ccacacctgt cccccctgcc ct 42

<210> 78  
 <211> 42  
 <212> DNA  
 <213> artificial

<220>  
 <223> artificial hinge region

<400> 78  
 cccaagagct ggcacaagtg ccacacctgt cccccctgcc ct 42

<210> 79  
 <211> 42  
 <212> DNA  
 <213> artificial

<220>  
 <223> artificial hinge region

<400> 79  
 cccaagagct ggcacaagac ccactgetgt cccccctgcc ct 42

<210> 80  
 <211> 36  
 <212> DNA  
 <213> artificial

<220>  
 <223> artificial hinge region

<400> 80  
 aagtgcgaca agaccacac ctgtcccccc tgcct 36

<210> 81  
 <211> 39  
 <212> DNA  
 <213> artificial

<220>  
 <223> artificial hinge region

<400> 81  
 cccaagagct gcgactgcca cacctgtccc cctgcct 39

<210> 82  
<211> 39  
<212> DNA  
<213> artificial  
  
<220>  
<223> artificial hinge region  
  
<400> 82  
cccaagagct gegactgcac ccaactgcccc ccctgcct 39

<210> 83  
<211> 36  
<212> DNA  
<213> artificial  
  
<220>  
<223> artificial hinge domain  
  
<400> 83  
ccctgcagct gcaagecac ctgtcccccc tgcct 36

<210> 84  
<211> 36  
<212> DNA  
<213> artificial  
  
<220>  
<223> artificial hinge region  
  
<400> 84  
cctagctgct gcacccacac ctgtcccccc tgcct 36

<210> 85  
<211> 36  
<212> DNA  
<213> artificial  
  
<220>  
<223> artificial hinge region  
  
<400> 85  
cccagctgcg acaageactg ctgtcccccc tgcct 36

<210> 86  
<211> 30  
<212> DNA  
<213> artificial



<220>

<223> artificial hinge region

<400> 86

cccaagagct gcacctgtcc ccctgtcct 30

<210> 87

<211> 42

<212> DNA

<213> artificial

<220>

<223> artificial hinge region

<400> 87

cccaagagct gegataagtg cgtggagtgc cccccctgtc ct 42

<210> 88

<211> 448

<212> PRT

<213> mus musculus

<400> 88

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala  
1 5 10 15

Ser Val Lys Ile Ser Cys Lys Thr Ser Gly Tyr Ile Phe Thr Ala Tyr  
20 25 30

Thr Met His Trp Val Arg Gln Ser Leu Gly Glu Ser Leu Asp Trp Ile  
35 40 45

Gly Gly Ile Lys Pro Asn Asn Gly Leu Ala Asn Tyr Asn Gln Lys Phe  
50 55 60

Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr  
65 70 75 80

Met Asp Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys  
85 90 95

Ala Arg Ser Glu Ile Thr Thr Glu Phe Asp Tyr Trp Gly Gln Gly Thr  
100 105 110

Ala Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro  
115 120 125

Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly  
130 135 140

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn  
145 150 155 160

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln  
165 170 175

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser  
180 185 190

Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser  
195 200 205

Asn Thr Lys Val Asp Lys Arg Val Glu Cys Lys Ser Cys Asp Lys Thr  
210 215 220

His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser  
225 230 235 240

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg  
245 250 255

Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro  
260 265 270

Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala  
275 280 285

Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val  
290 295 300

Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr  
305 310 315 320

Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr  
325 330 335

Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu  
340 345 350

Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys  
355 360 365

Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser  
370 375 380

Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp  
385 390 395 400

Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser  
405 410 415

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala  
420 425 430

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
435 440 445

<210> 89  
<211> 448  
<212> PRT  
<213> mus musculus

<400> 89

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala  
1 5 10 15

Ser Val Lys Ile Ser Cys Lys Thr Ser Gly Tyr Ile Phe Thr Ala Tyr  
20 25 30

Thr Met His Trp Val Arg Gln Ser Leu Gly Glu Ser Leu Asp Trp Ile  
35 40 45

Gly Gly Ile Lys Pro Asn Asn Gly Leu Ala Asn Tyr Asn Gln Lys Phe  
50 55 60

Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr

65            70            75            80

Met Asp Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys  
          85            90            95

Ala Arg Ser Glu Ile Thr Thr Glu Phe Asp Tyr Trp Gly Gln Gly Thr  
          100            105            110

Ala Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro  
          115            120            125

Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly  
          130            135            140

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn  
          145            150            155            160

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln  
          165            170            175

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser  
          180            185            190

Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser  
          195            200            205

Asn Thr Lys Val Asp Lys Arg Val Glu Pro Cys Ser Cys Asp Lys Thr  
          210            215            220

His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser  
          225            230            235            240

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg  
          245            250            255

Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro  
          260            265            270

Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala  
          275            280            285

Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val  
290 295 300

Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr  
305 310 315 320

Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr  
325 330 335

Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu  
340 345 350

Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys  
355 360 365

Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser  
370 375 380

Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp  
385 390 395 400

Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser  
405 410 415

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala  
420 425 430

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
435 440 445

<210> 90

<211> 448

<212> PRT

<213> mus musculus

<400> 90

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala  
1 5 10 15

Ser Val Lys Ile Ser Cys Lys Thr Ser Gly Tyr Ile Phe Thr Ala Tyr  
20 25 30

Thr Met His Trp Val Arg Gln Ser Leu Gly Glu Ser Leu Asp Trp Ile  
35 40 45

Gly Gly Ile Lys Pro Asn Asn Gly Leu Ala Asn Tyr Asn Gln Lys Phe  
50 55 60

Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr  
65 70 75 80

Met Asp Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys  
85 90 95

Ala Arg Ser Glu Ile Thr Thr Glu Phe Asp Tyr Trp Gly Gln Gly Thr  
100 105 110

Ala Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro  
115 120 125

Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly  
130 135 140

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn  
145 150 155 160

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln  
165 170 175

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser  
180 185 190

Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser  
195 200 205

Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Cys Cys Asp Lys Thr  
210 215 220

His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser  
225 230 235 240

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg  
245 250 255

Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro  
260 265 270

Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala  
275 280 285

Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val  
290 295 300

Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr  
305 310 315 320

Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr  
325 330 335

Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu  
340 345 350

Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys  
355 360 365

Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser  
370 375 380

Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp  
385 390 395 400

Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser  
405 410 415

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala  
420 425 430

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
435 440 445

<210> 91

<211> 448

<212> PRT

<213> mus musculus

<400> 91

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala  
1           5           10           15

Ser Val Lys Ile Ser Cys Lys Thr Ser Gly Tyr Ile Phe Thr Ala Tyr  
20           25           30

Thr Met His Trp Val Arg Gln Ser Leu Gly Glu Ser Leu Asp Trp Ile  
35           40           45

Gly Gly Ile Lys Pro Asn Asn Gly Leu Ala Asn Tyr Asn Gln Lys Phe  
50           55           60

Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr  
65           70           75           80

Met Asp Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys  
85           90           95

Ala Arg Ser Glu Ile Thr Thr Glu Phe Asp Tyr Trp Gly Gln Gly Thr  
100           105           110

Ala Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro  
115           120           125

Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly  
130           135           140

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn  
145           150           155           160

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln  
165           170           175

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser  
180           185           190

Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser



195            200            205

Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Cys Lys Thr  
210            215            220

His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser  
225            230            235            240

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg  
245            250            255

Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro  
260            265            270

Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala  
275            280            285

Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val  
290            295            300

Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr  
305            310            315            320

Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr  
325            330            335

Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu  
340            345            350

Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys  
355            360            365

Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser  
370            375            380

Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp  
385            390            395            400

Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser  
405            410            415

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala  
420 425 430

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
435 440 445

<210> 92

<211> 448

<212> PRT

<213> mus musculus

<400> 92

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala  
1 5 10 15

Ser Val Lys Ile Ser Cys Lys Thr Ser Gly Tyr Ile Phe Thr Ala Tyr  
20 25 30

Thr Met His Trp Val Arg Gln Ser Leu Gly Glu Ser Leu Asp Trp Ile  
35 40 45

Gly Gly Ile Lys Pro Asn Asn Gly Leu Ala Asn Tyr Asn Gln Lys Phe  
50 55 60

Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr  
65 70 75 80

Met Asp Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys  
85 90 95

Ala Arg Ser Glu Ile Thr Thr Glu Phe Asp Tyr Trp Gly Gln Gly Thr  
100 105 110

Ala Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro  
115 120 125

Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly  
130 135 140

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn  
145 150 155 160

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln  
165 170 175

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser  
180 185 190

Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser  
195 200 205

Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Cys Thr  
210 215 220

His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser  
225 230 235 240

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg  
245 250 255

Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro  
260 265 270

Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala  
275 280 285

Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val  
290 295 300

Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr  
305 310 315 320

Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr  
325 330 335

Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu  
340 345 350

Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys  
355 360 365

Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser  
370 375 380

Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp  
385 390 395 400

Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser  
405 410 415

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala  
420 425 430

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
435 440 445

<210> 93

<211> 448

<212> PRT

<213> mus musculus

<400> 93

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala  
1 5 10 15

Ser Val Lys Ile Ser Cys Lys Thr Ser Gly Tyr Ile Phe Thr Ala Tyr  
20 25 30

Thr Met His Trp Val Arg Gln Ser Leu Gly Glu Ser Leu Asp Trp Ile  
35 40 45

Gly Gly Ile Lys Pro Asn Asn Gly Leu Ala Asn Tyr Asn Gln Lys Phe  
50 55 60

Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr  
65 70 75 80

Met Asp Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys  
85 90 95

Ala Arg Ser Glu Ile Thr Thr Glu Phe Asp Tyr Trp Gly Gln Gly Thr  
100 105 110

Ala Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro  
115 120 125

Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly  
130 135 140

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn  
145 150 155 160

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln  
165 170 175

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser  
180 185 190

Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser  
195 200 205

Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Lys Cys  
210 215 220

His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser  
225 230 235 240

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg  
245 250 255

Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro  
260 265 270

Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala  
275 280 285

Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val  
290 295 300

Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr  
305 310 315 320

Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr

325            330            335

Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu  
340            345            350

Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys  
355            360            365

Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser  
370            375            380

Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp  
385            390            395            400

Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser  
405            410            415

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala  
420            425            430

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
435            440            445

<210> 94  
<211> 448  
<212> PRT  
<213> mus musculus

<400> 94

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala  
1            5            10            15

Ser Val Lys Ile Ser Cys Lys Thr Ser Gly Tyr Ile Phe Thr Ala Tyr  
20            25            30

Thr Met His Trp Val Arg Gln Ser Leu Gly Glu Ser Leu Asp Trp Ile  
35            40            45

Gly Gly Ile Lys Pro Asn Asn Gly Leu Ala Asn Tyr Asn Gln Lys Phe  
50            55            60

Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr  
65            70            75            80

Met Asp Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys  
          85            90            95

Ala Arg Ser Glu Ile Thr Thr Glu Phe Asp Tyr Trp Gly Gln Gly Thr  
      100            105            110

Ala Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro  
      115            120            125

Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly  
      130            135            140

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn  
      145            150            155            160

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln  
      165            170            175

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser  
      180            185            190

Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser  
      195            200            205

Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Lys Thr  
      210            215            220

His Cys Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser  
      225            230            235            240

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg  
      245            250            255

Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro  
      260            265            270

Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala  
      275            280            285

Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val  
290 295 300

Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr  
305 310 315 320

Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr  
325 330 335

Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu  
340 345 350

Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys  
355 360 365

Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser  
370 375 380

Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp  
385 390 395 400

Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser  
405 410 415

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala  
420 425 430

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
435 440 445

<210> 95  
<211> 446  
<212> PRT  
<213> mus musculus

<400> 95

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala  
1 5 10 15

Ser Val Lys Ile Ser Cys Lys Thr Ser Gly Tyr Ile Phe Thr Ala Tyr



20            25            30

Thr Met His Trp Val Arg Gln Ser Leu Gly Glu Ser Leu Asp Trp Ile  
35                    40                    45

Gly Gly Ile Lys Pro Asn Asn Gly Leu Ala Asn Tyr Asn Gln Lys Phe  
50                    55                    60

Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr  
65                    70                    75                    80

Met Asp Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys  
85                    90                    95

Ala Arg Ser Glu Ile Thr Thr Glu Phe Asp Tyr Trp Gly Gln Gly Thr  
100                    105                    110

Ala Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro  
115                    120                    125

Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly  
130                    135                    140

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn  
145                    150                    155                    160

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln  
165                    170                    175

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser  
180                    185                    190

Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser  
195                    200                    205

Asn Thr Lys Val Asp Lys Arg Val Glu Lys Cys Asp Lys Thr His Thr  
210                    215                    220

Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe  
225                    230                    235                    240

Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro  
245 250 255

Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val  
260 265 270

Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr  
275 280 285

Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val  
290 295 300

Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys  
305 310 315 320

Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser  
325 330 335

Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro  
340 345 350

Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val  
355 360 365

Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly  
370 375 380

Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp  
385 390 395 400

Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp  
405 410 415

Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His  
420 425 430

Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
435 440 445

<210> 96

<211> 447  
<212> PRT  
<213> mus musculus

<400> 96

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala  
1 5 10 15

Ser Val Lys Ile Ser Cys Lys Thr Ser Gly Tyr Ile Phe Thr Ala Tyr  
20 25 30

Thr Met His Trp Val Arg Gln Ser Leu Gly Glu Ser Leu Asp Trp Ile  
35 40 45

Gly Gly Ile Lys Pro Asn Asn Gly Leu Ala Asn Tyr Asn Gln Lys Phe  
50 55 60

Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr  
65 70 75 80

Met Asp Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys  
85 90 95

Ala Arg Ser Glu Ile Thr Thr Glu Phe Asp Tyr Trp Gly Gln Gly Thr  
100 105 110

Ala Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro  
115 120 125

Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly  
130 135 140

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn  
145 150 155 160

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln  
165 170 175

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser  
180 185 190

Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser  
195 200 205

Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Cys His  
210 215 220

Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val  
225 230 235 240

Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr  
245 250 255

Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu  
260 265 270

Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys  
275 280 285

Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser  
290 295 300

Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys  
305 310 315 320

Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile  
325 330 335

Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro  
340 345 350

Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu  
355 360 365

Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn  
370 375 380

Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser  
385 390 395 400

Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg  
405 410 415

Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu  
420 425 430

His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
435 440 445

<210> 97

<211> 447

<212> PRT

<213> mus musculus

<400> 97

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala  
1 5 10 15

Ser Val Lys Ile Ser Cys Lys Thr Ser Gly Tyr Ile Phe Thr Ala Tyr  
20 25 30

Thr Met His Trp Val Arg Gln Ser Leu Gly Glu Ser Leu Asp Trp Ile  
35 40 45

Gly Gly Ile Lys Pro Asn Asn Gly Leu Ala Asn Tyr Asn Gln Lys Phe  
50 55 60

Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr  
65 70 75 80

Met Asp Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys  
85 90 95

Ala Arg Ser Glu Ile Thr Thr Glu Phe Asp Tyr Trp Gly Gln Gly Thr  
100 105 110

Ala Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro  
115 120 125

Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly  
130 135 140

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn

145            150            155            160  
 Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln  
           165            170            175  
 Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser  
           180            185            190  
 Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser  
           195            200            205  
 Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Cys Thr  
           210            215            220  
 His Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val  
           225            230            235            240  
 Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr  
           245            250            255  
 Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu  
           260            265            270  
 Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys  
           275            280            285  
 Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser  
           290            295            300  
 Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys  
           305            310            315            320  
 Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile  
           325            330            335  
 Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro  
           340            345            350  
 Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu  
           355            360            365

Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn  
370 375 380

Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser  
385 390 395 400

Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg  
405 410 415

Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu  
420 425 430

His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
435 440 445

<210> 98

<211> 446

<212> PRT

<213> mus musculus

<400> 98

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala  
1 5 10 15

Ser Val Lys Ile Ser Cys Lys Thr Ser Gly Tyr Ile Phe Thr Ala Tyr  
20 25 30

Thr Met His Trp Val Arg Gln Ser Leu Gly Glu Ser Leu Asp Trp Ile  
35 40 45

Gly Gly Ile Lys Pro Asn Asn Gly Leu Ala Asn Tyr Asn Gln Lys Phe  
50 55 60

Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr  
65 70 75 80

Met Asp Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys  
85 90 95

Ala Arg Ser Glu Ile Thr Thr Glu Phe Asp Tyr Trp Gly Gln Gly Thr  
100 105 110

Ala Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro  
115 120 125

Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly  
130 135 140

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn  
145 150 155 160

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln  
165 170 175

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser  
180 185 190

Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser  
195 200 205

Asn Thr Lys Val Asp Lys Arg Val Glu Pro Cys Ser Cys Lys His Thr  
210 215 220

Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe  
225 230 235 240

Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro  
245 250 255

Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val  
260 265 270

Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr  
275 280 285

Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val  
290 295 300

Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys  
305 310 315 320



Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser  
325 330 335

Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro  
340 345 350

Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val  
355 360 365

Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly  
370 375 380

Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp  
385 390 395 400

Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp  
405 410 415

Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His  
420 425 430

Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
435 440 445

<210> 99

<211> 446

<212> PRT

<213> mus musculus

<400> 99

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala  
1 5 10 15

Ser Val Lys Ile Ser Cys Lys Thr Ser Gly Tyr Ile Phe Thr Ala Tyr  
20 25 30

Thr Met His Trp Val Arg Gln Ser Leu Gly Glu Ser Leu Asp Trp Ile  
35 40 45

Gly Gly Ile Lys Pro Asn Asn Gly Leu Ala Asn Tyr Asn Gln Lys Phe  
50 55 60

Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr  
65            70            75            80

Met Asp Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys  
          85            90            95

Ala Arg Ser Glu Ile Thr Thr Glu Phe Asp Tyr Trp Gly Gln Gly Thr  
          100            105            110

Ala Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro  
          115            120            125

Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly  
          130            135            140

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn  
145            150            155            160

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln  
          165            170            175

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser  
          180            185            190

Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser  
          195            200            205

Asn Thr Lys Val Asp Lys Arg Val Glu Pro Ser Cys Cys Thr His Thr  
          210            215            220

Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe  
225            230            235            240

Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro  
          245            250            255

Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val  
          260            265            270

Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr

275            280            285

Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val  
290            295            300

Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys  
305            310            315            320

Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser  
325            330            335

Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro  
340            345            350

Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val  
355            360            365

Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly  
370            375            380

Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp  
385            390            395            400

Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp  
405            410            415

Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His  
420            425            430

Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
435            440            445

<210> 100

<211> 446

<212> PRT

<213> mus musculus

<400> 100

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala  
1            5            10            15

Ser Val Lys Ile Ser Cys Lys Thr Ser Gly Tyr Ile Phe Thr Ala Tyr  
20 25 30

Thr Met His Trp Val Arg Gln Ser Leu Gly Glu Ser Leu Asp Trp Ile  
35 40 45

Gly Gly Ile Lys Pro Asn Asn Gly Leu Ala Asn Tyr Asn Gln Lys Phe  
50 55 60

Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr  
65 70 75 80

Met Asp Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys  
85 90 95

Ala Arg Ser Glu Ile Thr Thr Glu Phe Asp Tyr Trp Gly Gln Gly Thr  
100 105 110

Ala Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro  
115 120 125

Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly  
130 135 140

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn  
145 150 155 160

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln  
165 170 175

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser  
180 185 190

Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser  
195 200 205

Asn Thr Lys Val Asp Lys Arg Val Glu Pro Ser Cys Asp Lys His Cys  
210 215 220

Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe  
225 230 235 240

Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro  
245 250 255

Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val  
260 265 270

Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr  
275 280 285

Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val  
290 295 300

Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys  
305 310 315 320

Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser  
325 330 335

Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro  
340 345 350

Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val  
355 360 365

Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly  
370 375 380

Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp  
385 390 395 400

Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp  
405 410 415

Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His  
420 425 430

Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
435 440 445

<210> 101  
<211> 444  
<212> PRT  
<213> mus musculus

<400> 101

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala  
1           5           10           15

Ser Val Lys Ile Ser Cys Lys Thr Ser Gly Tyr Ile Phe Thr Ala Tyr  
20           25           30

Thr Met His Trp Val Arg Gln Ser Leu Gly Glu Ser Leu Asp Trp Ile  
35           40           45

Gly Gly Ile Lys Pro Asn Asn Gly Leu Ala Asn Tyr Asn Gln Lys Phe  
50           55           60

Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr  
65           70           75           80

Met Asp Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys  
85           90           95

Ala Arg Ser Glu Ile Thr Thr Glu Phe Asp Tyr Trp Gly Gln Gly Thr  
100           105           110

Ala Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro  
115           120           125

Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly  
130           135           140

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn  
145           150           155           160

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln  
165           170           175

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser  
180           185           190

Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser  
195 200 205

Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Thr Cys Pro  
210 215 220

Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe  
225 230 235 240

Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val  
245 250 255

Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe  
260 265 270

Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro  
275 280 285

Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr  
290 295 300

Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val  
305 310 315 320

Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala  
325 330 335

Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg  
340 345 350

Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly  
355 360 365

Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro  
370 375 380

Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser  
385 390 395 400

Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln

405 410 415

Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His  
420 425 430

Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
435 440

<210> 102

<211> 448

<212> PRT

<213> mus musculus

<400> 102

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala  
1 5 10 15

Ser Val Lys Ile Ser Cys Lys Thr Ser Gly Tyr Ile Phe Thr Ala Tyr  
20 25 30

Thr Met His Trp Val Arg Gln Ser Leu Gly Glu Ser Leu Asp Trp Ile  
35 40 45

Gly Gly Ile Lys Pro Asn Asn Gly Leu Ala Asn Tyr Asn Gln Lys Phe  
50 55 60

Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr  
65 70 75 80

Met Asp Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys  
85 90 95

Ala Arg Ser Glu Ile Thr Thr Glu Phe Asp Tyr Trp Gly Gln Gly Thr  
100 105 110

Ala Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro  
115 120 125

Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly  
130 135 140



Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn  
145            150            155            160

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln  
165            170            175

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser  
180            185            190

Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser  
195            200            205

Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Lys Cys  
210            215            220

Val Glu Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser  
225            230            235            240

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg  
245            250            255

Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro  
260            265            270

Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala  
275            280            285

Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val  
290            295            300

Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr  
305            310            315            320

Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr  
325            330            335

Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu  
340            345            350

Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys  
355            360            365

Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser  
370 375 380

Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp  
385 390 395 400

Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser  
405 410 415

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala  
420 425 430

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
435 440 445

<210> 103

<211> 1347

<212> DNA

<213> mus musculus

<400> 103

gagggtccagc tgcagcagag cgggccagaa ctgggttaaac ctggcggccag cgtgaagatt 60

agctgtaaga ccagcgggta catctttaca gcataacca tgcactgggt gaggcagagt 120

ctgggggaat ctctggactg gatcggagggt attaagccca acaatggcct ggctaactat 180

aatcaaaaat tcaagggcaa agccacactg accgtcgata agtctcttc cacagettac 240

atggatctga gaagcctgac atccgaggac agtgcagtgt actactgcgc ccggtctgag 300

atactaccg agttcgacta ttggggacag ggcactgcac tgaccgtctc ctccgccagc 360

accaagggcc caagegtgtt cccgctagcc cccagcagca agagcaccag cggcggcaca 420

gccgcctgg gctgectggt gaaggactac ttccccgagc ccgtgaccgt gtcttgaac 480

agcggagccc tgacctccgg cgtgcacaca ttccccgccc tgctgcagag cagcggcctg 540

tacagcctga gcagcgtggt gaccgtgctt agcagcagcc tgggcacca gacctacac 600

tgcaacgtga accacaagcc cagcaacacc aaggtggaca agcgggtgga gtgcaagagc 660

tgcgacaaga cccacacctg tccccctgc cctgccctg aactgctggg cggaccagc 720

gtgttctgt tccccccaa gcccaaggac acctgatga tcagcagaac ccccgaggtg 780

acctgtgtgg tggtagacgt gtcccacgag gaccagagg tgaagttcaa ctggtacgtg 840  
 gacggcgtgg aggtgcacaa cgccaagacc aagcccagag aggagcagta caacagcacc 900  
 tacaggggtgg tgtccgtgct gaccgtgctg caccaggact ggctgaacgg caaggagtac 960  
 aagtgaagg tgtccaacaa ggcctgcca gcccacatcg aaaagaccat cagcaaggcc 1020  
 aagggccagc caagagagcc ccaggtgtac acctgccac ccagcaggga ggagatgacc 1080  
 aagaaccagg tgtccctgac ctgtctggtg aagggttct acccaagcga catgcccgtg 1140  
 gagtgggaga gcaacggcca gcccgagaac aactacaaga ccaccccc agtgctggac 1200  
 agcgacggca gcttctct gtacagcaag ctgaccgtgg acaagagcag atggcagcag 1260  
 ggcaacgtgt tcagctgctc cgtgatgac gaggcctgc acaaccacta caccagaag 1320  
 agcctgagcc tgtcccagg caagtga 1347

<210> 104  
 <211> 1347  
 <212> DNA  
 <213> mus musculus

<400> 104  
 gaggtccagc tgcagcagag cgggccagaa ctggttaaac ctggcggcag cgtgaagatt 60  
 agctgaaga ccagcggfta catctttaca gcatatacca tgcactgggt gaggcagagt 120  
 ctgggggaat ctctggactg gatcggagggt attaagccca acaatggcct ggctaactat 180  
 aatcaaaaat tcaagggcaa agccacactg accgtcgata agtctcttc cacagcttac 240  
 atggatctga gaagcctgac atccaggagc agtgacagtgt actactgcgc ccggtctgag 300  
 atcaactaccg agttcgacta ttggggacag ggcactgcac tgaccgtctc ctccgccagc 360  
 accaagggcc caagcgtgtt cccgctagcc cccagcagca agagcaccag cggcggcaca 420  
 gccgcctgg gctgcctggt gaaggactac ttccccgagc ccgtgaccgt gtcttgaac 480  
 agcggagccc tgacctccgg cgtgcacaca ttccccgccg tgctgcagag cagcggcctg 540  
 tacagcctga gcagcgtggt gaccgtgctc agcagcagcc tgggcacca gacctacatc 600  
 tgcaacgtga accacaagcc cagcaacacc aaggtggaca agagagtgga gccctgcagc 660  
 tgcgacaaga cccacacctg tccccctgc cctgccctg aactgctggg cggaccagc 720  
 gtgttctgt tccccccaa gcccaaggac acctgatga tcagcagaac ccccgaggtg 780  
 acctgtgtgg tggtagacgt gtcccacgag gaccagagg tgaagttcaa ctggtacgtg 840

gacggcgtgg aggtgcacaa cgccaagacc aagcccagag aggagcagta caacagcacc 900  
tacagggtgg tgtccgtgct gaccgtgctg caccaggact ggctgaacgg caaggagtac 960  
aagtgtaaagg tgtccaacaa ggcctgcca gcccacatcg aaaagaccat cagcaaggcc 1020  
aagggccagc caagagagcc ccaggtgtac accctgccac ccagcaggga ggagatgacc 1080  
aagaaccagg tgtccctgac ctgtctggtg aagggttctt acccaagcga catcgccgtg 1140  
gagtgggaga gcaacggcca gcccgagaac aactacaaga ccaccccc agtgetggac 1200  
agcgacggca gcttctct gtacagcaag ctgaccgtgg acaagagcag atggcagcag 1260  
ggcaacgtgt tcagctgctc cgtgatgac gaggcctgc acaaccacta caccagaag 1320  
agcctgagcc tgtcccagg caagtga 1347

<210> 105  
<211> 1347  
<212> DNA  
<213> mus musculus

<400> 105  
gaggtccagc tgcagcagag cgggccagaa ctggftaac ctggcggcag cgtgaagatt 60  
agctgtaaga ccagcggfta catctttaca gcalatacca tgcaactgggt gaggcagagt 120  
ctgggggaat ctctggactg gatcggaggt attaagccca acaatggcct ggctaactat 180  
aatcaaaaat tcaagggcaa agccacactg accgtcgata agtctcttc cacagcttac 240  
atggatctga gaagcctgac atccgaggac agtgacgtgt actactgcgc ccggctgag 300  
atactaccg agttcgacta ttggggacag ggcactgcac tgaccgtctc ctccgccagc 360  
accaagggcc caagegtgtt cccgctagcc cccagcagca agagcaccag cggcggcaca 420  
gccgccctgg gctgectggt gaaggactac tccccgagc ccgtgaccgt gtcttggaac 480  
agcggagccc tgacctcgg cgtgcacaca tccccgccg tgctgcagag cagcggcctg 540  
tacagcctga gcagcgtggt gaccgtgctt agcagcagcc tgggcacca gacctacatc 600  
tgcaacgtga accacaagcc cagcaacacc aagtgggaca agcgcgtgga gcccaagtgc 660  
tgcgacaaga cccacacctg tccccctgc cctgccctg aactgtggg cggaccagc 720  
gtgttctctg tccccccaa gcccaaggac accctgatga tcagcagaac ccccaggtg 780  
acctgtgtgg tggtagcgt gtcccacgag gaccagagg tgaagttcaa ctggtactg 840

gacggcgtgg aggtgcacaa cgccaagacc aagcccagag agggagcagta caacagcacc 900  
tacaggggtgg tgtccgtgct gaccgtgctg caccaggact ggctgaacgg caaggagtac 960  
aagtgtlaagg tgtccaacaa ggcctgcca gcccacatcg aaaagaccat cagcaaggcc 1020  
aagggccagc caagagagcc ccaggtgtac accctgccac ccagcaggga ggagatgacc 1080  
aagaaccagg tgtccctgac ctgtctggtg aagggttctt acccaagcga catcgccgtg 1140  
gagtgaggaga gcaacggcca gcccgagaac aactacaaga ccaccccc agtgetggac 1200  
agcgacggca gcttctct gtacagcaag ctgaccgtgg acaagagcag atggcagcag 1260  
ggcaacgtgt tcagctgctc cgtgatgcac gaggccctgc acaaccacta caccagaag 1320  
agcctgagcc tgtccccagg caagtga 1347

<210> 106  
<211> 1347  
<212> DNA  
<213> mus musculus

<400> 106  
gaggtccagc tgcagcagag cgggccagaa ctggftaac ctggcggcag cgtgaagatt 60  
agctgtaaga ccagcggfta catctttaca gcatatacca tgcactgggt gaggcagagt 120  
ctgggggaat ctctggactg gatcggaggt attaagccca acaatggcct ggctaactat 180  
aatcaaaaat tcaaggcga agccacactg accgtcgata agtctcttc cacagcttac 240  
atggatctga gaagcctgac atccgaggac agtgcagtgt actactgcgc ccggtctgag 300  
atcactaccg agttcgacta ttggggacag ggcactgcac tgaccgtctc ctccgccagc 360  
accaagggcc caagcgtgtt cccgtagecc cccagcagca agagcaccag cggcggcaca 420  
gccgccctgg gctgectggt gaaggactac ttccccgagc cctgaccgt gtcttgaac 480  
agcggagccc tgacctcgg cgtgcacaca ttccccgccg tgctgcagag cagcggcctg 540  
tacagctga gcagcgtggt gaccgtgctc agcagcagcc tgggcacca gacctacac 600  
tgcaacgtga accacaagcc cagcaacacc aaggtggaca agagagtga gcctaagagc 660  
tgttgaaga cccacactg tccccctgc cctgcccctg aactgctggg cggaccagc 720  
gtgttctgt tccccccaa gcccaggac accctgatga tcagcagaac ccccaggtg 780  
acctgtgtgg tggtagcgt gtcccacgag gaccagagg tgaagtcaa ctggtactg 840  
gacggcgtgg aggtgcacaa cgccaagacc aagcccagag agggagcagta caacagcacc 900

tacagggtgg tgtccgtgct gaccgtgctg caccaggact ggctgaacgg caaggagtac 960  
aagtgtaaagg tgtccaacaa ggccctgccca gcccgaatcg aaaagaccat cagcaaggcc 1020  
aagggccagc caagagagcc ccaggtgtac accctgccac ccagcaggga ggagatgacc 1080  
aagaaccagg tgtccctgac ctgtctggtg aagggttctt acccaagcga catcgccgtg 1140  
gagtgggaga gcaacggcca gcccgagaac aactacaaga ccaccccc agtgctggac 1200  
agcgacggca gcttcttct gtacagcaag ctgaccgtgg acaagagcag atggcagcag 1260  
ggcaacgtgt tcagctgctc cgtgatgcac gaggccctgc acaaccacta caccagaag 1320  
agcctgagcc tgtccccagg caagtga 1347

<210> 107  
<211> 1346  
<212> DNA  
<213> mus musculus

<400> 107  
gaggtccagc tgcagcagag cgggccagaa ctggttaaac ctggcggcag cgtgaagatt 60  
agctgtaaaga ccagcgggta catctttaca gcatatacca tgcactgggt gaggcagagt 120  
ctggggggaat ctctggactg gatcggaggt attaagccca acaatggcct ggetaactat 180  
aatcaaaaat tcaagggcaa agccacactg accgtcgata agtctcttc cacagcttac 240  
atggatctga gaagcctgac atccgaggac agtgcagtgt actactgcgc ccggtctgag 300  
atactaccg agttcgaacta ttggggacag ggcactgcac tgaccgtctc ctccgccage 360  
accaagggcc caagcgtgtt cccgctagcc cccagcagca agagcaccag cggcggcaca 420  
gccgcctgg gctgcctggt gaaggactac tccccgagc cctgaccgt gtcttgaac 480  
agcggagccc tgacctcgg cgtgcacaca tccccgccg tgctgcagag cagcggcctg 540  
tacagcctga gcagcgtggt gaccgtgctc agcagcagcc tgggcacca gacctacac 600  
tgcaacgtga accacaagcc cagcaacacc aaggtggaca agagagtgga gcccaagagc 660  
tgcgactgca cccacactg tccccctgc cctgccctg aactgctggg cggaccagc 720  
gtgttctgt tccccccaa gcccaaggac accctgatga tcagcagaac ccccagagtg 780  
acctgtgtgg tgggtggacgt gtcccacgag gaccagagg tgaagttcaa ctggtactg 840  
gacggcgtgg aggtgcacaa cgccaagacc aagcccagag aggagcagta caacagcacc 900

tacagggtgg tgtccgtgct gaccgtgctg caccaggact ggctgaacgg caaggagtac 960  
aagtgtaaagg tgtccaacaa ggcctgcca gcccacatcg aaaagaccat cagcaaggcc 1020  
aagggccagc caagagagcc ccaggtgtac accctgccac ccagcagga ggagatgacc 1080  
aagaaccagg tgtccctgac ctgtctggtg aagggttctt acccaagcga catcgccgtg 1140  
gagtgggaga gcaacggcca gcccgagaac aactacaaga ccaccccc agtctggac 1200  
agcgacggca gcttctct gtacagcaag ctgaccgtgg acaagagcag atggcagcag 1260  
ggcaacgtgt tcagctgctc cgtgatgac gaggcctgc acaaccacta caccagaag 1320  
agcctgagcc tgtcccagg caagtg 1346

<210> 108  
<211> 1347  
<212> DNA  
<213> mus musculus

<400> 108  
gaggtccagc tgcagcagag cgggccagaa ctggftaac ctggcggcag cgtgaagatt 60  
agctgtaaga ccagcggfta catctttaca gcatatacca tgcactgggt gaggcagagt 120  
ctgggggaat ctctgactg gatcggaggt attaagccca acaatggcct ggctaactat 180  
aatcaaaaat tcaagggcaa agccacactg accgtcgata agtctcttc cacagcttac 240  
atggatctga gaagcctgac atccgaggac agtgcagtgt actactgcgc ccggtctgag 300  
atactaccg agttcgacta ttggggacag ggcactgac tgaccgtctc ctccgccagc 360  
accaagggcc caagcgtgtt cccgtagcc cccagcagca agagcaccag cggcggcaca 420  
gccgcctgg gctgctggt gaaggactac tccccgagc ccgtgaccgt gtctggaac 480  
agcggagccc tgacctccgg cgtgcacaca tccccgccg tgctgcagag cagcggcctg 540  
tacagcctga gcagcgtggt gaccgtgct agcagcagcc tgggcacca gacctacac 600  
tgcaacgtga accacaagcc cagcaacacc aaggtggaca agagagtga gcccaagagc 660  
tgcgacaagt gccacactg tccccctgc cctgcccctg aactgctggg cggaccagc 720  
gtgttctgt tcccccaa gcccaaggac accctgatga tcagcagaac ccccagagtg 780  
acctgtgtgg tgggtggact gtcccacgag gaccagagg tgaagttcaa ctggtactg 840  
gacggcgtgg aggtgcacaa cgccaagacc aagcccagag aggagcagta caacagcacc 900  
tacagggtgg tgtccgtgct gaccgtgctg caccaggact ggctgaacgg caaggagtac 960

aagtgttaagg tgtccaacaa ggccctgcca gcccacatcg aaaagaccat cagcaaggcc 1020  
aagggccagc caagagagcc ccaggtgtac accctgccac ccagcaggga ggagatgacc 1080  
aagaaccagg tgtccctgac ctgtctggtg aagggttct acccaagega catcgccgtg 1140  
gagtgggaga gcaacggcca gcccgagaac aactacaaga ccaccccccc agtgctggac 1200  
agcgacggca gcttcttct gtacagcaag ctgaccgtgg acaagagcag atggcagcag 1260  
ggcaacgtgt tcagctgctc cgtgatgca c gaggccctgc acaaccacta caccagaag 1320  
agcctgagcc tgtccccagg caagtga 1347

<210> 109  
<211> 1347  
<212> DNA  
<213> mus musculus

<400> 109  
gaggtccagc tgcagcagag cgggccagaa ctggttaaac ctggcggccag cgtgaagatt 60  
agctgtaaga ccagcggta catctttaca gcatatacca tgcactgggt gaggcagagt 120  
ctgggggaat ctctggactg gatcggaggt attaagccca acaatggcct ggctaactat 180  
aatcaaaaat tcaagggcaa agccacactg accgtcgata agtcctcttc cacagettac 240  
atggatctga gaagcctgac atccgaggac agtgcagtgt actactgcgc ccggtctgag 300  
atactaccg agttcgacta ttggggacag ggcactgcac tgaccgtctc ctccgccagc 360  
accaagggcc caagcgtgtt cccgctagcc cccagcagca agagcaccag cggcggcaca 420  
gccgcctgg gctgcctggt gaaggactac tccccgagc ccgtgaccgt gtcctggaac 480  
agcggagccc tgacctccgg cgtgcacacc tccccgccg tgcctgcagag cagcggcctg 540  
tacagcctga gcagcgtggt gaccgtgctt agcagcagcc tgggcacca gacctacatc 600  
tgcaacgtga accacaagcc cagcaacacc aaggtggaca agagagtga gcccaagagc 660  
tgcgacaaga cccactgtg tccccctgc cctgcccctg aactgctggg cggaccagc 720  
gtgttctgt tccccccaa gcccaaggac accctgatga tcagcagaac ccccagagtg 780  
acctgtgtgg tgggtggactg gtcccacgag gaccagagg tgaagtcaa ctggtactgt 840  
gacggcgtgg aggtgcacaa cgccaagacc aagcccagag aggagcagta caacagcacc 900  
tacagggtgg tgtccgtgct gaccgtgctg caccaggact ggctgaacgg caaggagtac 960



aagtgttaagg tgtccaacaa ggccctgccca gcccacatcg aaaagacat cagcaaggcc 1020  
aagggccagc caagagagcc ccaggtgtac accctgccac ccagcaggga ggagatgacc 1080  
aagaaccagg tgtccctgac ctgtctggtg aagggttct acccaagega catcgccgtg 1140  
gagtgggaga gcaacggcca gcccgagaac aactacaaga ccaccccc agtgctggac 1200  
agcgacggca gcttcttct gtacagcaag ctgaccgtgg acaagagcag atggcagcag 1260  
ggcaacgtgt tcagctgctc cgtgatgca gaggccctgc acaaccacta caccagaag 1320  
agcctgagcc tgtccccagg caagtga 1347

<210> 110  
<211> 1341  
<212> DNA  
<213> mus musculus

<400> 110  
gaggtccagc tgcagcagag cgggccagaa ctggttaaac ctggcggcag cgtgaagatt 60  
agctgtaaga ccagcgggta catctttaca gcatatacca tgcactgggt gaggcagagt 120  
ctggggggaat ctctggactg gatcggaggt attaagccca acaatggcct ggctaactat 180  
aatcaaaaat tcaagggcaa agccacactg accgtcgata agtctcttc cacagcttac 240  
atggatctga gaagcctgac atccgaggac agtgcagtgt actactgcgc ccggtctgag 300  
atactaccg agttegacta ttggggacag ggcactgcac tgaccgtctc ctccgccagc 360  
accaagggcc caagcgtgtt cccgctagcc cccagcagca agagcaccag cggcggcaca 420  
gccgcctgg gctgcctggt gaaggactac ttccccgagc ccgtgaccgt gtcctggaac 480  
agcggagccc tgacctcgg cgtgcacaca ttccccgccc tgctgcagag cagcggcctg 540  
tacagcctga gcagcgtggt gaccgtgctt agcagcagcc tgggcacca gacctacatc 600  
tgcaacgtga accacaagcc cagcaacacc aagtgggaca agcgggtgga gaagtgcgac 660  
aagaccaca cctgtcccc ctgccctgcc cctgaactgc tggcgggacc cagcgtgttc 720  
ctgttcccc ccaagcccaa ggacacctg atgatcagca gaacccccga ggtgacctgt 780  
gtggtggtgg acgtgtcca cgaggacca gaggtgaagt tcaactggtc cgtggacggc 840  
gtggaggtgc acaacgcaa gaccaagccc agagaggagc agtacaacag cacctacagg 900  
gtggtgtccg tgctgaccgt gctgcaccag gactggctga acggcaagga gtacaagtgt 960  
aaggtgtcca acaaggcct gccagccca atcgaaaaga ccatcagcaa ggccaagggc 1020

cagccaagag agccccaggt gtacacctg ccaccagca gggaggagat gaccaagaac 1080  
caggtgtccc tgacctgtct ggtgaagggc ttctaccaa gcgacatgc cgtggagtgg 1140  
gagagcaacg gccagcccga gaacaactac aagaccacce cccagtgct ggacagcgac 1200  
ggcagettct tctgtacag caagetgacc gtggacaaga gcagatggca gcagggaac 1260  
gtgttcagct gctccgtgat gcacgagcc ctgcacaacc actacacca gaagagcctg 1320  
agcctgtccc caggcaagtg a 1341

<210> 111  
<211> 1344  
<212> DNA  
<213> mus musculus

<400> 111  
gaggtccagc tgcagcagag cgggccagaa ctggttaaac ctggcggcag cgtgaagatt 60  
agctgtaaga ccagcggta catctttaca gcatatacca tgactgggt gaggcagagt 120  
ctgggggaat ctctggactg gatcggaggt attaagccca acaatggcct ggctaactat 180  
aatcaaaaat tcaagggcaa agccacactg accgtcgata agtctcttc cacagcttac 240  
atggatctga gaagcctgac atccgaggac agtgcagtgt actactgcgc ccggtctgag 300  
atcactaccg agttcgacta ttggggacag ggcactgcac tgaccgtctc ctccgccagc 360  
accaagggcc caagcgtgtt cccgctagcc cccagcagca agagcaccag cggcggcaca 420  
gccgcctgg gctgectggt gaaggactac ttccccgagc ccgtgaccgt gtcttgaac 480  
agcggagccc tgacctcgg cgtgcacaca ttccccgccg tgctgcagag cagcggcctg 540  
tacagcctga gcagcgtggt gaccgtgect agcagcagcc tgggcacca gacctacatc 600  
tgcaacgtga accacaagcc cagcaacacc aaggtggaca agagagtga gcccaagagc 660  
tgcgactgcc acacctgtcc cccctgccct gccctgaac tgetgggcgg acccagcgtg 720  
ttctgttcc ccccaagcc caaggacacc ctgatgatca gcagaacccc cgaggtgacc 780  
tgtgtggtgg tggacgtgtc ccacgaggac ccagaggtga agttcaactg gtacgtggac 840  
ggcgtggagg tgcacaacgc caagaccaag cccagagagg agcagtacaa cagcacctac 900  
aggggtgtgt ccgtgtgac cgtgtgcac caggactggc tgaacggcaa ggagtacaag 960  
tgtaaggtgt ccaacaagcc cctgccagcc ccaatcgaag agaccatcag caaggccaag 1020

ggccagccaa gagagcccca ggtgtacacc ctgccacca gcagggagga gatgaccaag 1080  
aaccaggtgt ccctgacctg tctggtgaag ggcttctacc caagcgacat cgccgtggag 1140  
tgggagagca acggccagcc cgagaacaac tacaagacca cccccccagt gctggacagc 1200  
gacggcagct tcttctgta cagcaagctg accgtggaca agagcagatg gcagcagggc 1260  
aacgtgtca gctgctccgt gatgcacgag gccctgcaca accactacac ccagaagagc 1320  
ctgagcctgt ccccaggcaa gtga 1344

<210> 112  
<211> 1344  
<212> DNA  
<213> mus musculus

<400> 112  
gaggtccage tgcagcagag cgggccagaa ctggttaaac ctggcggcag cgtgaagatt 60  
agctgtaaga ccagcgggta catctttaca gcatatacca tgcaactggg gaggcagagt 120  
ctggggggaat ctctggactg gatcggaggt ataaagccca acaatggcct ggctaactat 180  
aatcaaaaat tcaagggcaa agccacactg accgtcgata agtctcttc cacagcttac 240  
atggatctga gaagcctgac atccgaggac agtgcagtgt actactgcgc ccggtctgag 300  
atcactaccg agttcgacta ttggggacag ggcactgcac tgaccgtctc ctccgccagc 360  
accaagggcc caagcgtgtt cccgctagcc cccagcagca agagcaccag cggcggcaca 420  
gccgcctgg gctgcctggt gaaggactac ttccccgagc ccgtgaccgt gtcttggaac 480  
agcggagccc tgacctcgg cgtgcacacc ttccccgccg tgctgcagag cagcggcctg 540  
tacagcctga gcagcgtggt gaccgtgect agcagcagcc tgggcacca gacctacatc 600  
tgcaactgta accacaagcc cagcaacacc aaggtggaca agagagtgga gcccaagagc 660  
tgcgactgca cccactgccc ccctgcctt gccctgaac tgctgggcgg acccagcgtg 720  
ttctgttcc ccccaagcc caaggacacc ctgatgatca gcagaacccc cgaggtgacc 780  
tgtgtggtgg tggacgtgtc ccacgaggac ccagaggtga agttcaactg gtacgtggac 840  
ggcgtggagg tgcacaacgc caagaccaag cccagagagg agcagtacaa cagcacctac 900  
aggggtgtgt ccgtgctgac cgtgctgcac caggactggc tgaacggcaa ggagtacaag 960  
tgtaaggtgt ccaacaagcc cctgccagcc ccaatcgaag agaccatcag caaggccaag 1020  
ggccagccaa gagagcccca ggtgtacacc ctgccacca gcagggagga gatgaccaag 1080

aaccaggtgt ccctgacctg tctggtgaag ggcttctacc caagegacat cgccgtggag 1140  
tgggagagca acggccagcc cgagaacaac tacaagacca cccccccagt gctggacagc 1200  
gacggcagct tcttctgta cagcaagctg accgtggaca agagcagatg gcagcagggc 1260  
aacgtgttca gctgctccgt gatgcacgag gccctgcaca accactacac ccagaagagc 1320  
ctgagcctgt ccccaggcaa gtga 1344

<210> 113  
<211> 1341  
<212> DNA  
<213> mus musculus

<400> 113  
gaggtccagc tgcagcagag cgggccagaa ctggttaaac ctggcggcag cgtgaagatt 60  
agctgtaaga ccagcggfta catctttaca gcatatacca tgcactgggt gaggcagagt 120  
ctggggggaat ctctggactg gatcggaggt attaagecca acaatggcct ggctaactat 180  
aatcaaaaat tcaagggcaa agccacactg accgtcgata agtctcttc cacagcttac 240  
atggatctga gaagcctgac atccgaggac agtgcagtgt actactgcgc ccggtctgag 300  
atactaccg agttcgacla ttggggacag ggcactgcac tgaccgtctc ctccgccagc 360  
accaagggcc caagegtgtt cccgctagcc cccagcagca agagcaccag cggcggcaca 420  
gccgccctgg gctgctggt gaaggactac tccccgagc ccgtgaccgt gtcttggaac 480  
agcggagccc tgacctccgg cgtgcacaca tccccgccg tgctgcagag cagcggcctg 540  
tacagcctga gcagcgtggt gaccgtgctc agcagcagcc tgggcacca gacctacatc 600  
tgcaactgta accacaagcc cagcaacacc aaggtggaca agagagtga gcctgcagc 660  
tgcaagcaca cctgtcccc ctgcctgcc cctgaactgc tggcgggacc cagcgtgttc 720  
ctgttcccc ccaagcccaa ggacaccctg atgacagca gaacccccga ggtgacctgt 780  
gtggtggtgg acgtgtccca cgaggacca gaggtgaagt tcaactgta cgtggacggc 840  
gtggaggtgc acaacgcaa gaccaagccc agagaggagc agtacaacag cacctacagg 900  
gtggtgtccg tgctgacctg gctgcaccag gactggctga acggcaagga gtacaagtgt 960  
aaggtgtcca acaaggccct gccagccca atcgaaaaga ccatcagcaa ggccaagggc 1020  
cagccaagag agccccaggt gtacaccctg ccaccagca gggaggagat gaccaagaac 1080

caggtgtccc tgacctgtct ggtgaagggc ttctacccaa gcgacatcgc cgtggagtgg 1140  
gagagcaacg gccagcccga gaacaactac aagaccacc cccagtgct ggacagcgac 1200  
ggcagcttct tcctgtacag caagctgacc gtggacaaga gcagatggca gcagggcaac 1260  
gtgttcagct gctccgtgat gcacgaggcc ctgcacaacc actacacca gaagagcctg 1320  
agcctgtccc caggcaagtg a 1341

<210> 114  
<211> 1341  
<212> DNA  
<213> mus musculus

<400> 114  
gaggtccagc tgcagcagag cgggccagaa ctggttaaac ctggcggcag cgtgaagatt 60  
agctgtaaga ccagcggfta catctttaca gcatatacca tgcactgggt gaggcagagt 120  
ctggggggaat ctctggactg gatcggaggt attaagccca acaatggcct ggctaactat 180  
aatcaaaaat tcaagggcaa agccacactg accgtcgata agtctcttc cacagcttac 240  
atggatctga gaagcctgac atccgaggac agtgcagtgt actactgcgc ccggtctgag 300  
atactaccg agttcgacta ttggggacag ggcactgcac tgaccgtctc ctccgccagc 360  
accaagggcc caagcgtgtt cccgctagcc cccagcagca agagcaccag cggcggcaca 420  
gccgccctgg gctgectggt gaaggactac ttccccgagc ccgtgaccgt gtcttggaac 480  
agcggagccc tgacctccgg cgtgcacaca ttccccgccg tgctgcagag cagcggcctg 540  
tacagcctga gcagcgtggt gaccgtgcct agcagcagcc tgggcacca gacctacatc 600  
tgcaacgtga accacaagcc cagcaacacc aaggtggaca agagagtga gcctagctgc 660  
tgcaccaca cctgtcccc ctgccctgcc cctgaactgc tgggeggacc cagcgtgttc 720  
ctgttcccc ccaagcccaa ggacaccctg atgacagca gaacccccga ggtgacctgt 780  
gtggtggtgg acgtgtccca cgaggacca gaggtgaagt tcaactgta cgtggacggc 840  
gtggaggtgc acaacgcaa gaccaagccc agagaggagc agtacaacag cacctacagg 900  
gtggtgtccg tgctgaccgt gctgcaccag gactggctga acggcaagga gtacaagtgt 960  
aaggtgtcca acaaggccct gccagccca atcgaaaaga ccatcagcaa ggccaagggc 1020  
cagccaagag agccccaggt gtacaccctg ccaccagca gggaggagat gaccaagaac 1080  
caggtgtccc tgacctgtct ggtgaagggc ttctacccaa gcgacatcgc cgtggagtgg 1140

gagagcaacg gccagcccga gaacaactac aagaccacce cccagtgct ggacagcgac 1200

ggcagcttct tctgtacag caagctgacc gtggacaaga gcagatggca gcagggaac 1260

gtgttcagct gctccgtgat gcacgaggec ctgcacaacc actacacca gaagagcctg 1320

agcctgtccc caggcaagtg a 1341

<210> 115

<211> 1341

<212> DNA

<213> mus musculus

<400> 115

gaggtccagc tgcagcagag cgggccagaa ctggttaaac ctggcggcag cgtgaagatt 60

agctgtaaga ccagcgggta catctttaca geatatacca tgcactgggt gaggcagagt 120

ctgggggaat ctctggactg gatcggaggt attaagccca acaatggcct ggctaactat 180

aatcaaaaat tcaagggcaa agccacactg accgtcgata agtctcttc cacagcttac 240

atggatctga gaagcctgac atccgaggac agtgcagtgt actactgcgc ccggtctgag 300

atactaccg agttcgacta ttggggacag ggcaactgcac tgaccgtctc ctccgccagc 360

accaagggcc caagegtgtt cccgctagcc cccagcagca agagcaccag cggcggcaca 420

gccgccctgg gctgcctggt gaaggactac ttccccgagc cctgaccgt gtcctggaac 480

agcggagccc tgacctccgg cgtgcacacc ttccccgccc tgctgcagag cagcggcctg 540

tacagcctga gcagcgtggt gaccgtgcct agcagcagcc tgggcacca gacctacatc 600

tgcaacgtga accacaagcc cagcaacacc aagtgggaca agagagtgga gccagctgc 660

gacaagcaet getgecccc ctgcctgcc cctgaactgc tgggcggacc cagcgtgttc 720

ctgttcccc ccaagcccaa ggacacctg atgatcagca gaacccccga ggtgacctgt 780

gtggtggtgg acgtgtccca cgaggacca gaggtgaagt tcaactgta cgtggacggc 840

gtggaggtgc acaacgcaa gaccaagccc agagaggagc agtacaacag cacctacagg 900

gtggtgtccc tgctgaccgt gctgcaccag gactggctga acggcaagga gtacaagtgt 960

aaggtgtcca acaaggcct gccagccca atcgaaaaga ccatcagcaa ggccaagggc 1020

cagccaagag agccccaggt gtacacctg ccaccagca gggaggagat gaccaagaac 1080

caggtgtccc tgacctgtct ggtgaagggc ttctaccaa gcgacatgc cgtggagtgg 1140

gagagcaacg gccagcccga gaacaactac aagaccaccc ccccagtgct ggacagcgac 1200  
ggcagcttct tctgtacag caagctgacc gtggacaaga gcagatggca gcagggaac 1260  
gtgttcagct gctccgtgat gcacgagggc ctgcacaacc actacacca gaagagcctg 1320  
agcctgtccc caggcaagtg a 1341

<210> 116  
<211> 1335  
<212> DNA  
<213> mus musculus

<400> 116  
gaggtccagc tgcagcagag cgggccagaa ctggttaaac ctggcggcag cgtgaagatt 60  
agctgtaaga ccagcgggta catctttaca geatatacca tgcactgggt gaggcagagt 120  
ctgggggaat ctctggactg gatcggaggt attaagccca acaatggcct ggctaactat 180  
aatcaaaaat tcaagggcaa agccacactg accgtcgata agtctcttc cacagcttac 240  
atggatctga gaagcctgac atccaggagc agtgcagtgt actactgcgc ccggtctgag 300  
atactaccg agttcgacta ttggggacag ggcactgcac tgaccgtctc ctccgccagc 360  
accaagggcc caagegtgtt cccgctagcc cccagcagca agagcacaag cggcgggaaca 420  
gccgccctgg gctgcctggt gaaggactac ttccccgagc ccgtgaccgt gtcttggaac 480  
agcggagccc tgaccagcgg cgtgcacacc ttccagccg tgctgcagag cagcggcctg 540  
tacagcctga gcagcgtggt gaccgtgcct agcagcagcc tgggcacca gacctacatc 600  
tgcaacgtga accacaagcc cagcaacacc aaggtggaca agagagtgga gcccaagagc 660  
tgcacctgct cccctgtcc tgcccctgag ctgctgggcg gaccagcgt gttctgttc 720  
cccccaagc ccaaggacac cctgatgac agcagaaccc ccgaggtgac ctgtgtggtg 780  
gtggacgtgt cccacgagga cccagaggtg aagttcaact ggtacgtgga cggcgtggag 840  
gtgcacaacg ccaagaccaa gccagagag gagcagtaca acagcaccta cagggtggtg 900  
tccgtgctga ccgtgctgca ccaggactgg ctgaacggca aggagtataa gtgtaaggtg 960  
tccaacaagg cctgccagc cccaatcgaa aagaccatca gcaaggccaa gggccagcca 1020  
agagagcccc aggtgtacac cctgccacce agcagggagg agatgaccaa gaaccaggtg 1080  
tcctgacct gtctggtgaa gggcttctac ccaagcgaca tcgccgtgga gtgggagagc 1140  
aacggccagc ccgagaacaa ctacaagacc accccccag tgctggacag cgacggcagc 1200

ttcttctgt acagcaagct gaccgtggac aagagcagat ggcagcaggg caacgtgttc 1260  
agctgctccg tgatgcacga ggcctgcac aacctatacccagaagag cctgagcctg 1320  
tccccaggca agtga 1335

<210> 117  
<211> 1347  
<212> DNA  
<213> mus musculus

<400> 117  
gaggtccagc tgcagcagag cgggccagaa ctggttaaac ctggcggcag cgtgaagatt 60  
agctgtaaga ccagcgggta catctttaca gcatatacca tgcactgggt gaggcagagt 120  
ctgggggaat ctctggactg gatcggaggt ataaagccca acaatggcct ggctaactat 180  
aatcaaaaat tcaagggcaa agccacactg accgtcgata agtcctcttc cacagcttac 240  
atggatctga gaagcctgac atccgaggac agtgcagtgt actactgcgc ccggtctgag 300  
atcactaccg agtgcacta ttggggacag ggcactgcac tgaccgtctc ctccgccagc 360  
accaagggcc caagcgtgtt cccgctagcc cccagcagca agagcacaag cggcgggaaca 420  
gccgccctgg gctgcctggt gaaggactac ttccccgagc cctgaccgt gtcttggaaac 480  
agcggagccc tgaccagcgg cgtgcacacc ttccagccg tgctgcagag cagcggcctg 540  
tacagcctga gcagcgtggt gaccgtgect agcagcagcc tgggcacca gacctacac 600  
tgcaacgtga accacaagcc cagcaacacc aaggtggaca agagagtgga gcccaagagc 660  
tgcgataagt gcgtggagtg cccccctgt cctgccctg agctgctggg cggaccagc 720  
gtgttctgt tccccccaa gcccaaggac acctgatga tcagcagaac ccccagagtg 780  
acctgtgtgg tgggtggacgt gtcccacgag gaccagagg tgaagtcaa ctggtactg 840  
gacggcgtgg aggtgcacaa cgccaagacc aagcccagag aggagcagta caacagcacc 900  
tacagggtgg tgtccgtgct gaccgtgctg caccaggact ggctgaacgg caaggagtac 960  
aagtgtlaagg tgtccaacaa ggcctgcca gcccacatc aaaagaccat cagcaaggcc 1020  
aagggccagc caagagagcc ccaggtgtac acctgccac ccagcagga ggagatgacc 1080  
aagaaccagg tgtccctgac ctgtctggtg aagggttct acccaagcga catcggcctg 1140  
gagtgggaga gcaacggcca gcccgagaac aactacaaga ccaccccc agtgctggac 1200



agcgacggca gcttcttct gtacagcaag ctgaccgtgg acaagagcag atggcagcag 1260

ggcaacgtgt tcagctgctc cgtgatgcac gaggcctgc acaaccacta caccagaag 1320

agcctgagec tgtccccagg caagtga 1347

1