



US 20020100072A1

(19) **United States**

(12) **Patent Application Publication**
Kikuchi et al.

(10) **Pub. No.: US 2002/0100072 A1**

(43) **Pub. Date: Jul. 25, 2002**

(54) **NOVEL PLANT GENES**

(76) Inventors: **Yasuhiro Kikuchi**, Tsukuba-shi (JP);
Shigeto Kiyokawa, Aomori-shi (JP);
Yukihisa Shimada, Tokyo (JP);
Masaya Ohbayashi, Tsukuba-shi (JP);
Ritsuko Shimada, Tokyo (JP); **Yasushi**
Okinaka, Sapporo-shi (JP)

now Pat. No. 6,114,601, which is a continuation of application No. 08/295,746, filed on Aug. 30, 1994, now abandoned, which is a continuation of application No. PCT/JP92/01520, filed on Nov. 20, 1992.

(30) **Foreign Application Priority Data**

Mar. 2, 1992 (JP) 44963/92

Publication Classification

(51) **Int. Cl.⁷** **A01H 5/00**; C07H 21/04;
C12N 9/02; C12N 5/04
(52) **U.S. Cl.** **800/278**; 435/410; 435/189;
536/23.2

Correspondence Address:

FITZPATRICK CELLA HARPER & SCINTO
30 ROCKEFELLER PLAZA
NEW YORK, NY 10112 (US)

(21) Appl. No.: **09/814,786**

(22) Filed: **Mar. 23, 2001**


(57) **ABSTRACT**

The present invention relates to a DNA which encodes a polypeptide having flavonoid-3',5'-hydroxylase activity, a recombinant DNA containing said DNA, and a plant having a pigment pattern which the plant does not originally have and which is acquired by transformation with said recombinant DNA.

Related U.S. Application Data

(60) Division of application No. 09/616,990, filed on Jul. 14, 2000, now Pat. No. 6,232,109, which is a division of application No. 08/606,505, filed on Feb. 23, 1996,

FIG. 1



※ () may be any amino acid.

NOVEL PLANT GENES

TECHNICAL FIELD

[0001] The present invention relates to a technique to breed plants or plant cells using recombinant DNA technology. More particularly, the present invention relates to a technique to breed novel plant cells or novel plants which show exogenous pigment patterns by transforming plant cells or plants with a recombinant DNA containing a DNA which encodes a polypeptide having flavonoid-3',5'-hydroxylase activity (hereinafter also referred to as the DNA encoding flavonoid-3',5'-hydroxylase).

BACKGROUND ART

[0002] Crossing between varieties has been conventionally employed as a method for altering the color of flowers and fruits of plants. However, crossing is carried out between varieties of the same genus, and usually of the same species, and therefore, it is extremely difficult to give specific colors to certain plant species. For example, in spite of longtime efforts of breeders, no one has yet successfully bred a blue rose or a blue carnation.

[0003] In recent years, recombinant DNA technology has enabled plant breeding between different species or genus, and it is expected to breed new plants having unprecedented pigment patterns which can not be obtained by the conventional breeding methods by crossing (Plant Molecular Biology, vol.13, p.287-294, 1989). For example, it is reported that petunia showing unprecedented brick-red color on flowers was bred by cloning a gene encoding dihydroflavonol-4-reductase, which is an enzyme participating in pigment biosynthetic pathway, from maize and introducing it into petunia (Japanese Published Unexamined Patent Application No. 2305/90; Nature, vol.330, p.677-678, 1987). Further, a report has been made of a case in which new pigment patterns were produced by introducing the chalcone synthase gene of petunia at the sense or anti-sense orientation to partially inhibit the expression of the gene (Nature, vol.333, p.866-869, 1988; The Plant Cell, vol.2, p.279-289, 1990; The Plant Cell, vol.2, p.291-299, 1990).

[0004] Biosynthetic pathways for anthocyanins, which contribute to blue or red color of flowers, have been studied genetically and biochemically in detail using petunia and others (Petunia, Edited by K. C. Sink, Springer Verlag, p.49-76, 1984; The Flavonoids, Edited by J. B. Harborne, Chapman and Hall, p.399-425, 1988; Molecular Approaches to Crop Improvement, Edited by E. S. Denis and D. J. Rewerin, Springer Verlag, p.127-148, 1991). As a result of these studies, it is shown that the presence/absence of hydroxyl group at the 3'- and 5'-positions of the B ring of anthocyanin greatly affects the color of flowers, and also it is shown that, generally, the blue color of flowers is intensified as the B ring is hydroxylated in a higher degree. The hydroxylation of the B ring of anthocyanins occurs at the stage of their precursors, flavanones or dihydroflavonols. As enzymes which catalyze this hydroxylation, two types of enzyme have been known; flavonoid-3'-hydroxylase which hydroxylates only the 3'-position of the B ring, and flavonoid-3',5'-hydroxylase which hydroxylates both the 3'- and 5'-positions. Petunia with blue flowers has both the enzymes, but that with red flowers has only the former one. Plants like roses, carnations, and chrysanthemums do not

have anthocyanins which have B ring hydroxylated at both the 3'- and 5'-positions, and therefore are considered not to have the latter type of enzyme.

[0005] These hydroxylases are localized in the microsomal membrane and require NADPH as a coenzyme. They are presumed to be members of the cytochrome P450 enzyme group on the basis of their behavior against various inhibitors (The Flavonoids, Edited by J. B. Harborne, Chapman and Hall, p.399-425, 1988; Molecular Approaches to Crop Improvement, Edited by E. S. Denis and D. J. Rewerin, Springer Verlag, p.127-148, 1991).

[0006] Cytochrome P450 is an enzyme group which is widely distributed among eucaryotes and procaryotes and which is involved in the biosynthesis of important lipids such as steroids and in the oxidative metabolism of lipophilic substances. In higher animals, it forms a super family consisting of one hundred or more molecular species (J. Biol. Chem., vol.266, p.13469-13472, 1991; Pharmacol. Rev., vol.40, p.243-288, 1988). In plants, cinnamic acid-4-hydroxylase and kaurene oxidase are considered to belong to the cytochrome P450 group (Plant Physiol., vol.96, p.669-674, 1991). Further, a gene encoding a cytochrome P450 enzyme whose function is unknown has been cloned from avocado (Proc. Natl. Acad. Sci. USA, vol.87, p.3904-3908, 1990). As a result of the comparison of the amino acid sequences of various types of cytochrome P450 enzymes, it is known that the sequence of the heme-binding site is conserved (Proc. Natl. Acad. Sci. USA, vol.85, p.7221-7225, 1988; Pharmacol. Rev. vol.40, p.243-288, 1988).

[0007] In petunia, flavonoid-3',5'-hydroxylase is encoded by two dominant genes called Hf-1 and Hf-2. The enzymes encoded by the genes are isozymes, and the degree of expression of Hf-1 is higher (Petunia, Edited by K. C. Sink, Springer Verlag, p.49-76, 1984). Further, characteristics of said enzyme of Verbena have been reported (Z. Naturforschung, vol.37c, p.19-23, 1982).

[0008] It is also reported that 3',5'-hydroxylase, a key enzyme in the biosynthesis of delphinidin, which is a blue pigment in petunia, has been successfully cloned (Nikkei Biotech, Aug. 26, 1991). However, no report has been made yet of a case in which the cloned gene of said enzyme is allowed to express in a plant to alter pigments in the plant.

DISCLOSURE OF THE INVENTION

[0009] The present invention provides a DNA which encodes a polypeptide having flavonoid-3',5'-hydroxylase activity which is represented by the amino acid sequence shown by SEQ ID NO: 1, 63 or 64, a DNA which hybridizes with said DNA, a recombinant DNA constructed by incorporating any of these DNAs or a part of their sequences into a vector DNA, and plant cells or plants which carry said recombinant DNA.

[0010] It is possible to breed plants having novel pigment patterns by introducing said DNA, i.e., a DNA that encodes a polypeptide having flavonoid-3',5'-hydroxylase activity, into plant cells or plants by the use of recombinant DNA technology.

[0011] The DNA of the present invention may be any DNA which encodes a polypeptide having flavonoid-3',5'-hydroxylase activity, i.e., a DNA which encodes a polypeptide represented by the amino acid sequence shown by SEQ

ID NO: 1, 63 or 64, or a DNA which hybridizes with said DNA (hereinafter referred to as hDNA). The hDNA may be any DNA which hybridizes with the DNA encoding the polypeptide represented by the amino acid sequence shown by SEQ ID NO: 1, 63 or 64 in 2×SSC (0.3 M sodium chloride, 0.03 M sodium citrate, pH 7.0) at 50° C.

[0012] The DNA which encodes the polypeptide represented by the amino acid sequence shown by SEQ ID NO: 63 or 64 hybridizes with the DNA which encodes the polypeptide represented by the amino acid sequence shown by SEQ ID NO: 1 under the above-mentioned conditions.

[0013] Other examples of the DNAs of the present invention include DNAs wherein a part of the nucleotide sequence of the above-mentioned DNAs is deleted or replaced with other nucleotide sequences, as far as such DNAs encode a polypeptide having flavonoid-3',5'-hydroxylase activity.

[0014] Examples of the DNA sources include a genomic DNA of plants which have flavonoid-3',5'-hydroxylase, and a cDNA which is synthesized from an mRNA extracted from the expression sites of said enzyme using a reverse transcriptase. Examples of the plants having said enzyme include petunia (Solanaceae), pansy (Violaceae), primrose (Primulaceae), delphinium (Ranunculaceae), sweet pea (Leguminosae), Japanese gentian (Gentianaceae), balloon flower (Campanulaceae), forget-me-not (Boraginaceae), hydrangea (Saxifragaceae), verbena (Verbenaceae), day-flower (Commelinaceae), iris (Iridaceae), hyacinth (Liliaceae), Russell prairie gentian (Gentianaceae), and campanula (Campanulaceae).

[0015] In the present invention, on the basis of the presumption that flavonoid-3',5'-hydroxylase is a member of the cytochrome P450 family, DNA sequences encoding the amino acid sequence of the heme-binding site of cytochrome P450 (hereinafter referred to as the core sequence) are amplified and isolated using the PCR method.

[0016] The core sequence is the region that shows high homology among different molecular species of cytochrome P450 and among those of different organisms. More than 80% of the sequences for cytochrome P450 which have been ever isolated have the core sequence shown in FIG. 1 (DNASIS™ Data Base CD, 009-1 and 2, Hitachi Software Engineering Co., Ltd., 1990). DNA sequences which can encode the amino acid sequence of the region indicated by arrows are hypothesized. Then, in order to amplify and isolate the DNA sequences encoding this region by the PCR method, 16 types of sense primers shown by SEQ ID NO: 2 to 17, and 12 types of antisense primers shown by SEQ ID NO: 18 to 29 are chemically synthesized. The sense primers are synthetic DNA primers each consisting of 18 bases, and each sense primer has, at the 3' end side, one of the 16 types of 8-base DNA sequences at the 3' end encoding Pro-Phe-Gly or Pro-Phe-Ser, and has, at the 5' end side, a 10-base DNA sequence which includes a recognition site for a restriction enzyme, EcoRI. The antisense primers are synthetic DNA primers each consisting of 18 bases, and each primer has, at the 3' end side, one of the 12 types of sequences which are inversely linked to the 8-base DNA sequence at the 3' end encoding Cys-Xxx-Gly (wherein Xxx represents Ile, Leu, Val, Ala, or Pro), and has, at the 5' end side, a 10-base DNA sequence including a recognition site for a restriction enzyme, BamHI.

[0017] By the use of PCR in which these synthetic DNA primers are employed in combination, various DNA frag-

ments which encode the core sequence can be amplified and isolated, and their DNA sequences can be determined. As cytochrome P450 forms a super family which consists of various molecular species, it is expected that various types of core sequences can be obtained from one template DNA. During the process of the present invention, 15 types of core sequences shown by SEQ ID NO: 30 to 44 were obtained.

[0018] It is necessary to make a presumption as to which core sequence is the target sequence among the thus obtained core sequences. In the present invention, the target sequence is presumed by investigating whether the expression/non-expression of each core sequence is genetically linked to the presence/absence of said enzyme activity. In order to investigate the genetic linkage, a petunia which originally has said enzyme (a blue flower cultivar) is backcrossed with a mutant petunia variety which lacks said enzyme (a red flower cultivar) to produce a genetically segregating population (1:1) regarding the presence/absence of said enzyme. Then, the mode of expression of each core sequence in the petals of individual plants in this population is investigated. If the mode of expression of any core sequence agrees with (is genetically linked with) the presence/absence of said enzyme, the core sequence is presumed to be a part of the gene encoding said enzyme.

[0019] In order to investigate whether a core sequence is expressed in the petals, the present invention uses a method called SSP (single specific primer) polymerase chain reaction (PCR). SSP.PCR is a method described in Biochemistry Biophysics Research Communication, vol.167, p.504-506, 1990. By the use of this method, it is possible to amplify a DNA sequence flanking a core sequence and to determine the presence/absence of the corresponding product. First, specific DNA primers are synthesized based on the DNA sequences encoding the core sequences. In the present invention, 15 types of DNAs (K primers 01 to 15) shown by SEQ ID NO: 45 to 59 were synthesized and used as the specific DNA primers. Then, cDNAs are prepared from the petals of each petunia plant in the backcrossed population, digested with appropriate restriction enzymes, and then ligated with appropriate double-strand synthetic DNA (called cassette) which had the corresponding cleaved ends using a ligase to prepare templates. In the present invention, synthetic DNAs shown by SEQ ID NO: 60 and 61 were annealed and used as the cassette. The synthetic DNA shown by SEQ ID NO: 60 was also used as the primer for the cassette. With the template DNA ligated to the cassette, PCR was carried out between the specific primer and the primer for the cassette, whereby the DNA sequence flanking the core sequence is amplified. The presence/absence of its product reflects the expression/non-expression of the core sequence.

[0020] As a result of the search in the petunia population obtained by the backcrossing, it was revealed that the presence/absence of a product (approximately 85 bp) which was amplified by SSP.PCR using the specific primer (K14) shown by SEQ ID NO: 58 was completely linked with the presence/absence of said enzyme activity. As this primer was designed based on the core sequence shown by SEQ ID NO: 43, this sequence is assumed to be the core sequence of said enzyme. On the basis of SEQ ID NO: 43, the primer (J14) shown by SEQ ID NO: 62 was synthesized and SSP.PCR was carried out. As a result, the presence/absence of a product of approximately 280 bp was completely linked

with the presence/absence of the enzyme activity. This result strongly suggests that the core sequence shown by SEQ ID NO: 43 is the target sequence.

[0021] The product of approximately 280 bp thus amplified is assumed to be a part of the cDNA sequence that encodes said enzyme. The full length cDNA sequence shown by SEQ ID NO: 1 can be obtained by preparing petunia flower cDNA library according to the method described in a book by Maniatis et al., and then searching the library using the above-mentioned product as a probe. If the expression of the obtained sequence in a plant which originally does not have said enzyme results in the detection of said enzyme activity in the plant, it will be proved that this sequence is the DNA sequence encoding the polypeptide having said enzyme activity. In the present invention, the DNA shown by SEQ ID NO: 1 was introduced into tobacco and petunia cultivars both of which do not have said enzyme, and expressed. As a result, said enzyme activity was detected in both plants, and thus the DNA was proved to be the DNA encoding the polypeptide having said enzyme activity.

[0022] Cloning of DNAs can be carried out using a material such as a cDNA which is synthesized based on an mRNA extracted from the petals of petunia using a reverse transcriptase.

[0023] DNA cloning and DNA analysis can be carried out according to general techniques described in *Molecular Cloning a Laboratory Manual Second Edition*, J. Sambrook, E. F. Fritsch, T. Maniatis, Cold Spring Harbor Laboratory Press, 1989 (hereinafter referred to as the book by Maniatis et al.), and the like.

[0024] PCR can be carried out according to ordinary techniques described in *PCR Technology*, Edited by H. A. Ehrlich, Stockton Press, 1989, *PCR Protocols*, Edited by M. A. Innis, D. H. Gelfand, J. J. Sninsky, and T. J. White, Academic Press, 1990, and the like.

[0025] Determination of nucleotide sequences can be carried out according to methods using the Taq Dideoxy™ Terminator Cycle Sequencing Kit (ABI Co., Ltd.) and the Model 373A DNA Sequencing System (ABI Co., Ltd.), and the like.

[0026] DNA fragments encoding polypeptides which have analogous sequences and said enzyme activity can be cloned from any of the plants mentioned above as the DNA source by an ordinary method using, as a probe for hybridization, the whole or a part of the DNA sequence shown by SEQ ID NO: 1 which encodes the polypeptide having said enzyme activity and is derived from petunia as above.

[0027] In the present invention, according to the above-mentioned method, a DNA which encodes a polypeptide having the amino acid sequence shown by SEQ ID NO: 63 has been cloned from Russell prairie gentian, and a DNA which encodes a polypeptide having the amino acid sequence shown by SEQ ID NO: 64 has been cloned from campanula.

[0028] New coloration can be introduced into a host plant which does not have said enzyme by introducing a DNA fragment which encodes a polypeptide having said enzyme activity into the host plant, allowing it to express, and thereby hydroxylating the 3'- and 5'-positions of anthocya-

nin pigments. Examples of such host plants include rose (Rosaceae), carnation (Caryophyllaceae), petunia (Solanaceae), tobacco (Solanaceae), chrysanthemum (Compositae), stock (Cruciferae), begonia (Begoniaceae), snapdragon (Scrophulariaceae), camellia (Theaceae), lily (Liliaceae), and orchid (Orchidaceae).

[0029] Further, in plant species which originally have said enzyme, the enzyme activity can be inhibited by introducing said DNA fragment at the antisense or sense orientation and allowing it to express (Nature, vol.333, p.866-869, 1988; The Plant Cell, vol.2, p.279-289, 1990; The Plant Cell, vol.2, p.291-299, 1990). By application of such methods, breeding of a plant species having an unprecedented pigment pattern can be achieved.

[0030] In order to introduce the DNA fragment which encodes the polypeptide having said enzyme activity into plants and allow it to express, it is necessary to introduce an appropriate promoter at the site upstream of the region encoding the polypeptide having said enzyme activity. An example of a promoter that works in plants is 35 S promoter of Cauliflower Mosaic Virus (CaMV) (Cell, vol.21, p.285-294, 1980). An example of a promoter that acts site-specifically is the promoter of petunia chalcone synthase (CHS) gene which works strongly only in the petals (Plant Molecular Biology, vol.15, p.95-109, 1990). The above-mentioned DNA fragment can be expressed in plants by ligating such a promoter. When a DNA which encodes the polypeptide having said enzyme activity is cloned from the genomic DNA, it may have been linked with an inherent promoter, and in such cases, there is no need to further link it with another promoter.

[0031] Further, efficient expression can be expected by introducing a terminator for the termination of transcription at the site downstream of the region encoding the polypeptide having said enzyme activity (EMBO Journal, vol.7, p.791-799, 1988).

[0032] In order to select plant cells or plants in which the DNA has been introduced, it is preferable to introduce an appropriate marker into the DNA. Examples of such markers include the kanamycin resistance gene and the hygromycin resistance gene (Plant Molecular Biology, vol.5, p.299-302, 1985). When a microorganism belonging to the genus *Agrobacterium* is used to introduce the DNA into plant cells or plants, it is necessary to attach the border sequences derived from Ti plasmid at both ends of the sequence to be inserted into plant chromosomes (Nature, vol.313, p.191-196, 1985). Further, it is necessary to link the insert sequence with a sequence that allows stable retention of plasmids in a cell of a microorganism belonging to the genus *Agrobacterium*. An example of an expression vector for plants which meets the above-mentioned requirements is pBI121 (Clontech Co., Ltd.).

[0033] Examples of methods for introducing said DNA fragment inserted in a vector as described above into plants and obtaining genetically stable transformed plants include: 1) a method for dicotyledons in which the DNA is introduced via *Agrobacterium tumefaciens*, the bacterium causing crown gall disease (Methods in Enzymology, vol.118, p.627-640, 1986); 2) a method in which the DNA is pelted in conjunction with microparticles of substances such as gold and tungsten at plant cells at a high speed to be incorporated into cell nuclei and then into chromosomes (the

high-speed microparticle method; *Plant Molecular Biology*, vol.11, p.433-439, 1989; *Bio/Technology*, vol.9, p.1080-1085, 1991); and 3) a method in which the DNA is introduced in conjunction with calcium chloride and polyethylene glycol into protoplasts which have been prepared with cell wall-degrading enzymes (*Nature*, vol.296, p.72-74, 1982; *Nature*, vol.319, p.791-793, 1986). The method 1) can be efficiently carried out by incorporating the insert DNA into a binary vector such as pBI121 (*Nucleic Acids Research*, vol.12, p.8711-8721, 1984). According to the method 2), the DNA can be introduced into plants which cannot be infected with a microorganism belonging to the genus *Agrobacterium* such as monocotyledons. After the introduction of said DNA fragment incorporated into a vector into plant cells according to the methods described above, plant cells in which the introduced DNA is stably retained in the chromosome are selected by utilizing appropriate marker genes such as those for drug resistance. By inducing the differentiation of such plant cells, transformed plants having novel pigment patterns can be obtained.

[0034] In the thus obtained transformed plants, the DNA fragments introduced are retained with genetic stability. In other words, said DNA fragments can be maintained semi-persistently through propagation by vegetative reproduction, or by seeds obtained through self-pollination or cross pollination.

[0035] Further, it is possible to breed new cultivars which have pigment patterns different from those of the first-generation transformants by crossing the transformants with conventional cultivars to combine their genes.

[0036] Thus, a technique is provided which enables the production of unprecedented cultivars having blue or purple flowers by allowing plants having no anthocyanin pigments whose B ring is hydroxylated at both the 3'- and 5'-positions, for example, roses and carnations, to synthesize such pigments.

BRIEF DESCRIPTION OF THE DRAWINGS

[0037] FIG. 1 shows the core sequence which is common to more than 80% of the known amino acid sequences for cytochrome P450.

BEST MODE FOR CARRYING OUT THE INVENTION

EXAMPLE 1

[0038] PCR Amplification and Isolation of the Core Sequences of Cytochrome P450 Genes and Their Sequencing

[0039] (1) Synthesis of Primers

[0040] A part of the gene sequence of cytochrome P450 was amplified and isolated by the polymerase chain reaction (PCR) in the following manner. Cytochrome P450 forms a super family consisting of various molecular species, but the similarity in the amino acid sequence among these molecular species is not so high. However, the sequences of the heme-binding region (core sequence) are relatively common.

[0041] More than 80% of the sequences for cytochrome P450 ever isolated have the core sequence shown in FIG. 1.

DNA sequences which could encode the amino acid sequence of the region indicated by arrows were hypothesized. Then, in order to amplify and isolate the DNA sequences encoding this region by the PCR method, primer DNAs were chemically synthesized using the DNA synthesizer, Cyclone Plus™ (manufactured by Milligen/Bioscience). Thus, 16 types of sense primers shown by SEQ ID NO: 2 to 17, and 12 types of antisense primers shown by SEQ ID NO: 18 to 29 were synthesized.

[0042] The sense primers are synthetic DNA primers each consisting of 18 bases, and each primer has, at the 3' end side, one of the 16 types of 8-base DNA sequences at the 3' end which encode Pro-Phe-Gly or Pro-Phe-Ser, and has, at the 5' end side, a 10-base DNA sequence which includes a recognition site for a restriction enzyme, EcoRI. The antisense primers are synthetic DNA primers each consisting of 18 bases, and each primer has, at the 3' end side, one of the 12 types of sequences which are inversely linked to the 8-base DNA sequence at the 3' end encoding Cys-Xxx-Gly (wherein Xxx represents Ile, Leu, Val, Ala, or Pro), and has, at the 5' end side, a 10-base DNA sequence including a recognition site for a restriction enzyme, BamHI.

[0043] Each primer was used in a 5 μM aqueous solution.

[0044] (2) Extraction of mRNAs from the Petals of *Petunia*

[0045] Extraction of mRNAs from the petals of *petunia* was carried out according to a modification of the method described in *Analytical Biochemistry*, vol.163, p.16-20, 1987. That is, petals were cut off from buds of *petunia* [*Petunia hybrida* cv. Falcon Blue (Sakata Seed Corporation)] which had been grown in a greenhouse. Ten grams (wet weight) of the petals was put into a mortar, frozen by pouring liquid nitrogen, and then ground with a pestle. To the ground petals were added 20 ml of RNA extraction buffer [8 M guanidine hydrochloride, 20 mM Mes buffer (pH 7.0), 20 mM EDTA, 50 mM mercaptoethanol] and then 10 ml of phenol/chloroform/isoamyl alcohol (25:24:1) mixture, and mixed well. The resulting mixture was centrifuged at 10,000×g for 10 minutes, and the upper layer was collected and mixed well with 20 ml of phenol/chloroform/isoamyl alcohol (25:24:1) mixture. The resulting mixture was centrifuged at 10,000×g for 10 minutes, and the upper layer was collected. Then, 14 ml of ethanol and 4 ml of 1 M acetic acid were added to the upper layer, and the mixture was allowed to stand at -70° C. for one hour, followed by centrifugation at 10,000×g for 10 minutes. The precipitate was separated, dissolved in 10 ml of water, and then mixed with 3 ml of 10 M lithium chloride. The resulting mixture was allowed to stand at 4° C. for 2 hours, and centrifuged at 10,000×g for 10 minutes. The precipitate was separated, washed with 10 ml of 70% ethanol, and then dried under vacuum. The dried product was dissolved in 1 ml of elution buffer [10 mM Tris hydrochloride buffer (pH 7.5), 1 mM EDTA, 0.1% sodium dodecyl sulfate (SDS)], and then subjected to purification using 200 μl of oligotex™-dT 30 (Takara Shuzo Co., Ltd.) according to the instructions provided by the manufacturer to give about 3 μg of poly(A)mRNA.

[0046] (3) Synthesis of cDNA from *Petunia* Petal mRNA

[0047] A cDNA was synthesized from oligo dT primer using the above-mentioned mRNA as the template and the

cDNA Synthesis System Plus RPN1256 (Amersham Co., Ltd.) according to the instructions provided by the manufacturer. About 2 μg of double strand cDNA was obtained.

[0048] (4) PCR Amplification of the Consensus Sequence of Cytochrome P450

[0049] The above-mentioned cDNA (1 ng) as a template DNA was dissolved in 25 μl of PCR buffer [10 mM Tris hydrochloride buffer (pH 8.3), 1.5 mM magnesium chloride, 25 mM potassium chloride, 0.05% Tween 20, 100 μM dATP, 100 μM dCTP, 100 μM dGTP, 100 μM dTTP]. The solution was put in a 0.5-ml microcentrifugation tube, and as primers, 1 μl of a sense primer (one type) and 1 μl of an antisense primer (one type) both of which were prepared in the step (1) were added thereto. To the mixture was added 0.5 unit of Taq DNA polymerase (Perkin-Elmer Cetus), and 10 μl of mineral oil was layered over the mixture. The reaction was carried out using the DNA Thermal Cycler (Perkin-Elmer Cetus) with the cycle program set as follows; 30 seconds at 93° C. and 1 minute at 37° C. for 3 cycles, followed by 30 seconds at 93° C. and 1 minute at 55° C. for 37 cycles.

[0050] PCR was carried out under the above conditions for each of all the 192 combinations of 16 sense primers and 12 antisense primers.

[0051] PCR was carried out by reference to PCR Technology, edited by H. A. Ehrlich, Stockton Press, 1989, and PCR Protocols, edited by M. A. Innis, D. H. Gerfand, J. J. Sninsky, and T. J. White, Academic Press, 1990.

[0052] (5) Cloning of PCR Products

[0053] The products of the above reaction were subjected to 10% polyacrylamide gel electrophoresis and stained with ethidium bromide according to the method described in the book by Maniatis et al. As a result, a DNA band of approximately 50 bp was detected for 23 among the 192 combinations of sense primers and antisense primers. Portions containing the DNA band were cut out from the gel, and DNAs were extracted and purified according to the methods described in the book by Maniatis et al. Each of the obtained DNAs was dissolved in 50 μl of H buffer [50 mM Tris hydrochloride buffer (pH 7.5), 10 mM magnesium chloride, 1 mM dithiothreitol, 100 mM sodium chloride]. To the solution were added 10 units of the restriction enzyme BamHI (Takara Shuzo Co., Ltd.) and 10 units of the restriction enzyme EcoRI (Takara Shuzo Co., Ltd.), and the reaction was carried out at 37° C. for 3 hours. After addition of 150 μl of ethanol, the reaction mixture was allowed to stand at -80° C. for 10 minutes, and then centrifuged at 10,000 \times g for 10 minutes. The obtained precipitate was washed with 200 μl of 70% ethanol and dried under vacuum. The obtained DNA was dissolved in 10 μl of TE buffer [10 mM Tris hydrochloride buffer (pH 7.5), 1 mM EDTA].

[0054] The plasmid vector pUC19 (Takara Shuzo Co., Ltd.) (5 μg) was dissolved in 50 μl of H buffer, and 10 units of the restriction enzyme BamHI (Takara Shuzo Co., Ltd.) and 10 units of the restriction enzyme EcoRI (Takara Shuzo Co., Ltd.) were added. The reaction was carried out at 37° C. for 3 hours. After addition of 150 μl of ethanol, the reaction mixture was allowed to stand at -80° C. for 10 minutes, and then centrifuged at 10,000 \times g for 10 minutes. The obtained precipitate was washed with 200 μl of 70% ethanol and dried under vacuum. The obtained vector DNA was dissolved in 100 μl of TE buffer.

[0055] The vector solution thus prepared (1 μl) was mixed with the solution containing the DNA fragment of approximately 50 bp (10 μl) prepared above, and subjected to ligation at 16° C. for 30 minutes using the DNA Ligation Kit (Takara Shuzo Co., Ltd.) according to the instructions provided by the manufacturer. The volume of the reaction mixture was 60 μl . Highly competent cells of *E. coli* JM109 (Toyobo Co., Ltd.) were transformed with 2 μl of the reaction mixture according to the instructions provided by the manufacturer. According to the method described in the book by Maniatis et al., the cells were cultured at 37° C. for 20 hours on X-gal ampicillin LB agar medium [1% Bacto Tryptone (Difco Laboratories), 0.5% Bacto Yeast Extract (Difco Laboratories), 1% sodium chloride, 40 $\mu\text{g}/\text{ml}$ X-gal, 40 $\mu\text{g}/\text{ml}$ isopropyl-1-thio- β -D-galactopyranoside (IPTG), 100 $\mu\text{g}/\text{ml}$ ampicillin, 1.5% Bacto Agar (Difco Laboratories)]. One of the formed white colonies was isolated and cultured, and plasmid DNA was extracted from the culture and purified.

[0056] (6) Determination of DNA Sequences of PCR Products

[0057] The nucleotide sequence of the insert fragment in each of the 108 clones prepared as described above were determined using the Taq Dideoxy™ Terminator Cycle Sequencing Kit (ABI) and the Model 373A DNA Sequencing System (ABI) according to the instructions provided by the manufacturer. As a result, 15 types of core sequences shown by SEQ ID NO: 30 to 44 were determined for the cytochrome P450 genes.

EXAMPLE 2

[0058] Production of Petunia Backcrossed Population

[0059] (1) Analysis of Pigments in the Petals

[0060] Pigments in the petals were analyzed after converting anthocyanins into anthocyanidins according to the method described in Phytochemical Methods, Second Edition, edited by J. B. Harbone, p.64, Chapman and Hall, 1989. That is, 0.1 to 0.5 g of the petals was cut off and 1 ml of 2N hydrochloric acid was added. The mixture was heated at 95° C. for 40 minutes, and then brought to room temperature. After addition of 300 μl of ethyl acetate followed by thorough mixing, the mixture was allowed to stand still, and the upper ethyl acetate layer was discarded. The residue was heated at 80° C. for 3 minutes to evaporate ethyl acetate, and then brought to room temperature. After addition of 100 μl of isoamyl alcohol followed by thorough mixing, the mixture was allowed to stand still, and the upper isoamyl alcohol layer was collected. Aliquots of 1 to 5 μl of the obtained solution were spotted on a cellulose thin layer plate (Merck & Co., Inc.) and chromatographed using Solvent 1 (con-c. hydrochloric acid:acetic acid:water=3:30:10) or Solvent 2 (n-butanol:acetic acid:water=4:1:5) to identify anthocyanidins based on the Rf values and coloration of the pigment spots. Separately, analysis was also carried out using the Hitachi Ion Chromato System (Model L6200 pump and Model L4200 detector), YMC-Pack ODS-A Reversed Phase Column (YMC), and mobile phase consisting of water, acetic acid and methanol (71:10:19) [New High Performance Liquid Chromatography, Application II, p.528, Hirokawa Shoten, 1983]. Anthocyanidins were identified by using commercially available cyanidin, delphinidin, peonidin, and maruvidin (all produced by Extrasynthese) as standards.

[0061] (2) Production of Petunia Backcrossed Population

[0062] Crossing of petunia was carried out according to the method described in Petunia, edited by K. C. Sink, p.180-202, Springer Verlag, 1984. A blue flower petunia cultivar, Purple Joy (NPI Seeds) was crossed with a red flower petunia cultivar, Falcon Red (Sakata Seed Corporation) to obtain hybrids. The hybrids were backcrossed with Falcon Red, and anthocyanidins in the petals of the obtained hybrids were analyzed. A hybrid plant which had delphinidin as the anthocyanidin component was selected and then backcrossed with Falcon Red. After such backcrossing was repeated four times in total, anthocyanidins in the petals of 18 plants of the obtained hybrid population were analyzed. Among them ten hybrids had delphinidin (delphinidin-type) as anthocyanidin and eight hybrids had cyanidin (cyanidin-type). The color of petals of the former type was grayish purple, and that of the latter was red.

[0063] (3) Detection of Flavonoid-3',5'-hydroxylase Activity

[0064] Detection of flavonoid-3',5'-hydroxylase activity was carried out according to a modification of the method described in Z. Naturforsch, vol.37c, p.19-23, 1982. That is, 5 g (wet weight) of petals of buds was disrupted using mortar and pestle at 0° C., with 2.5 g of quartz sand (Sigma), 2.5 g of Dow X 1x2 (The Dow Chemical), and 10 ml of buffer for enzyme extraction [0.1 M potassium phosphate buffer (pH 7.5), 20% glycerol, 10 mg/ml sodium ascorbate]. After centrifugation at 12,000xg for 20 minutes, the obtained supernatant (10 ml) was mixed with 0.4 ml of 1 M magnesium chloride. The mixture was allowed to stand at 0° C. for 10 minutes, and centrifuged at 17,000xg for 20 minutes to obtain precipitate. The precipitate was suspended in a small quantity of the buffer for enzyme extraction to make a final volume of 500 μ l, and the suspension was used as the microsome fraction.

[0065] An aliquot of 100 μ l of the microsome fraction was mixed with 400 μ l of a reaction mixture [0.1 M potassium phosphate buffer (pH 7.5), 20% glycerol, 10 mg/ml sodium ascorbate, 0.25 mM NADPH (Sigma), 0.25 mM dihydroquercetin (Sigma)], and allowed to react at 25° C. for 30 minutes. After addition of 250 μ l of ethyl acetate, the mixture was allowed to stand still, and the upper layer (ethyl acetate layer) was collected, followed by evaporation of ethyl acetate. The residue was dissolved in 10 μ l of ethyl acetate, and an aliquot of 5 μ l of the solution was spotted on a cellulose thin layer plate (Merck & Co., Inc.) and chromatographed using Solvent 3 (chloroform:acetic acid:water=10:9:1). Flavonoids detected under the UV light were identified based on the Rf values. As a result, it was shown that dihydroquercetin had been converted into dihydromyricetin by the action of flavonoid-3',5'-hydroxylase.

[0066] Among the plants of the above-mentioned population obtained by backcrossing, said enzyme activity was detected in the delphinidin-type plants. On the other hand, it was not detected in the cyanidin-type plants. Further, said enzyme activity was detected in Falcon Blue (Sakata Seed Corporation) and Purple Joy (NPI Seeds), which were blue flower petunia cultivars, but was not detected in Falcon Red (Sakata Seed Corporation) and Falcon Salmon (Sakata Seed Corporation), which were red flower petunia cultivars.

EXAMPLE 3

[0067] SSP.PCR Using the Core Sequence of Cytochrome P450**[0068]** (1) Synthesis of K Primers

[0069] On the basis of 15 types of the core sequences for cytochrome P450 shown by SEQ ID NO: 30 to 44 which were obtained in Example 1 (6), 15 types of PCR primers shown by SEQ ID NO: 45 to 59 were chemically synthesized using the DNA Synthesizer Cyclone Plus (Milligen/Biosearch). Each primer was used in a 5 μ M aqueous solution. The primers were named K01 to K15 primers, respectively, and collectively referred to as K primers. K primers are synthetic DNA primers each having a 17-base sequence which starts from the codon for glycine located at the C-terminus of the amino acid sequence of the core and extends toward the N-terminus, and correspond to the sequences from the 32nd nucleotide to the 16th nucleotide in the core DNA sequences shown by SEQ ID NO: 30 to 44.

[0070] (2) Synthesis of a Cassette and a Primer for the Cassette

[0071] Oligonucleotides indicated by SEQ ID NO: 60 and 61 were chemically synthesized using the DNA Synthesizer Cyclone Plus (Milligen/Biosearch), and a 20 μ M aqueous solution of each oligonucleotide was prepared. After 100 μ l each of the solutions were mixed, the mixture was heated at 95° C. for 10 minutes, and then kept at 50° C. for one hour to obtain a double strand DNA, which is called a cassette. One end of the cassette forms a cohesive end of CG-protruding type, and therefore, can be efficiently linked with a restriction end digested with restriction enzymes, such as HinPI, MaeII, MspI and TthHB8I.

[0072] Separately, a 5 μ M aqueous solution of the oligonucleotide shown by SEQ ID NO: 60 was prepared and used as the primer for the cassette.

[0073] (3) Synthesis of Petunia Petal cDNA

[0074] Four plants of the delphinidin-type and two plants of the cyanidin-type were selected from the backcrossed population produced in Example 2, and cDNAs were synthesized using mRNAs extracted from the petals of each plant according to the methods described in Example 1 (2) and (3). Similarly, cDNAs were synthesized from petals of Falcon Blue, Falcon Red, Falcon Salmon, and Purple Joy.

[0075] (4) TthHB8I Digestion of cDNA and Linkage to Cassette

[0076] An aliquot of 0.1 μ g of each of the ten types of cDNAs obtained in (3) above was dissolved in 50 μ l of H buffer, and one unit of the restriction enzyme TthHB8I (Takara Shuzo Co., Ltd.) was added. The reaction was carried out at 65° C. for one hour. Then, the reaction mixture was mixed with 5 μ l of phenol/chloroform (1:1) mixture, followed by addition of 150 ml of ethanol. The resulting mixture was allowed to stand at -80° C. for 10 minutes, and centrifuged at 10,000xg for 10 minutes. The obtained precipitate was washed with 200 μ l of 70% ethanol, and dried under vacuum. The obtained DNA was dissolved in 9 μ l of TE buffer.

[0077] After adding 1 μ l of the cassette to each DNA solution, ligation reaction was carried out at 16° C. for 30 minutes using the DNA Ligation Kit (Takara Shuzo Co.,

Ltd.) according to the instructions provided by the manufacturer. The volume of the reaction mixture was 60 μ l.

[0078] (5) PCR Between K Primers and the Primer for Cassette

[0079] By the use of each of the above reaction mixtures as a template, the sequence to a near restriction site can be amplified by PCR between a K primer (01 to 15) and the primer for the cassette.

[0080] An aliquot of 1 μ l of each of the above-mentioned reaction mixtures, which was used as a template, was added to 1 μ l of a K primer and 1 μ l of the primer for the cassette, and mixed with 25 μ l of PCR buffer. The mixture was transferred into a 0.5-ml microcentrifugation tube, 0.5 unit of Taq DNA polymerase (Perkin-Elmer Cetus) was added thereto, and 10 μ l of mineral oil was layered over the mixture. The reaction was carried out using the DNA Thermal Cycler (Perkin-Elmer Cetus) for 40 cycles with the cycle profile consisting of 30 seconds at 93° C. and 1 minute at 55° C. According to the methods described in the book by Maniatis et al., the PCR products were subjected to 10% polyacrylamide gel electrophoresis, and DNA bands were stained with ethidium bromide and examined under UV light.

[0081] As a result, in the SSP.PCR using K14 primer, a DNA band of about 85 bp was obtained when one of the six types of cDNAs obtained from Falcon Blue, Purple Joy, and four delphinidin-type backcrossed plants was used as the template. On the other hand, the band was not detected when one of the four types of cDNAs obtained from Falcon Red, Falcon Salmon, and two cyanidin-type backcrossed plants was used as the template. That is, it was demonstrated that the presence/absence of the SSP.PCR products of about 85 bp was genetically linked to the presence/absence of said enzyme activity. When the other primers were used, no such product was detected. As the K14 primer was designed based on the core sequence shown by SEQ ID NO: 43, it was suggested that the sequence shown by SEQ ID NO: 43 was a part of the DNA sequence encoding the polypeptide which had said enzyme activity.

[0082] (6) Synthesis of J14 Primer

[0083] On the basis of the core sequence of cytochrome P450 shown by SEQ ID NO: 43, according to which K14 primer was synthesized, a primer shown by SEQ ID NO: 62 was chemically synthesized using the DNA Synthesizer Cyclone Plus (Milligen/Bioscience). The primer was named J14 primer, and used in a 5 μ M aqueous solution.

[0084] (7) HinPI Digestion of cDNA and Linkage to Cassette

[0085] An aliquot of 0.1 μ g of each of the ten types of cDNAs obtained in (3) above was dissolved in 50 μ l of M buffer [10 mM Tris hydrochloride buffer (pH 7.5), 10 mM magnesium chloride, 1 mM dithiothreitol, 50 mM sodium chloride], and one unit of the restriction enzyme HinPI (New England Biolabs) was added. The reaction was carried out at 37° C. for one hour. Then, the reaction mixture was mixed with 5 μ l of phenol/chloroform (1:1) mixture, followed by addition of 150 μ l of ethanol. The resulting mixture was allowed to stand at -80° C. for 10 minutes, and centrifuged at 10,000 \times g for 10 minutes. The obtained precipitate was washed with 200 μ l of 70% ethanol, and dried under

vacuum. The obtained DNA was dissolved in 9 μ l of TE buffer. After adding 1 μ l of the cassette to each DNA solution, ligation reaction was carried out at 16° C. for 30 minutes using the DNA Ligation Kit (Takara Shuzo Co., Ltd.) according to the instructions provided by the manufacturer. The volume of the reaction mixture was 60 μ l.

[0086] (8) PCR Between J14 Primer and the Primer for Cassette

[0087] An aliquot of 1 μ l of each of the above-mentioned reaction mixtures, which was used as a template, was added to 1 μ l of J14 primer and 1 μ l of the primer for the cassette, and mixed with 25 μ l of PCR buffer. The mixture was transferred into a 0.5-ml microcentrifugation tube, 0.5 unit of Taq DNA polymerase (Perkin-Elmer Cetus) was added thereto, and 10 μ l of mineral oil was layered over the mixture. The reaction was carried out using the DNA Thermal Cycler (Perkin-Elmer Cetus) for 40 cycles with the cycle profile consisting of 30 seconds at 93° C. and 1 minute at 55° C. According to the methods described in the book by Maniatis et al., the PCR products were subjected to 10% polyacrylamide gel electrophoresis, and DNA bands were stained with ethidium bromide and examined under UV light.

[0088] As a result, in the SSP.PCR using J14 primer, a DNA band of about 280 bp was obtained when one of the six types of cDNAs obtained from Falcon Blue, Purple Joy, and four delphinidin-type backcrossed plants was used as the template. On the other hand, the band was not detected when one of the four types of cDNAs obtained from Falcon Red, Falcon Salmon, and two cyanidin-type backcrossed plants was used as the template. That is, it was demonstrated that the presence/absence of the SSP.PCR products of about 280 bp was genetically linked to the presence/absence of said enzyme activity. It was strongly suggested that the core sequence shown by SEQ ID NO: 43 was a part of the DNA sequence encoding the polypeptide which had said enzyme activity.

EXAMPLE 4

[0089] Construction and Sequencing of Plasmid pEAK14

[0090] A library is constructed by incorporating petunia petal cDNAs into an appropriate vector. The library is searched using the SSP.PCR product of about 280 bp obtained in Example 3 as a probe, and the sequence of a clone which hybridizes with the probe is determined.

[0091] (1) Construction of Petunia Petal cDNA Library

[0092] One microgram of cDNA prepared from the petals of petunia (Falcon Blue) in Example 1 (3) was cloned using the cDNA Cloning System λ gt10.RPN1257 (Amersham Co., Ltd.) according to the instructions provided by the manufacturer. The final product was subjected to the packaging reaction using the XDNA in vitro packaging kit Giga Pack Gold (Stratagene Co., Ltd.) according to the instructions provided by the manufacturer. Cells of *E. coli* NM 514 (Amersham Co., Ltd.) were infected with the appropriately diluted packaging products according to the instructions provided by the manufacturer, and spread on LB agar medium [1% Bacto Tryptone (Difco Laboratories), 0.5% Bacto Yeast Extract (Difco Laboratories), 1% sodium chloride, 1.5% Bacto Agar (Difco Laboratories)] in plastic plates

of 15 cm in diameter (Iwaki Glass Co., Ltd.) to obtain about 10,000 plaques per plate. A total of five plates were prepared.

[0093] (2) Radiolabeling of DNA Probe

[0094] A portion containing the PCR product of about 280 bp obtained in Example 3 was cut out from the polyacrylamide gel, and the PCR product was extracted and purified according to the method described in the book by Maniatis et al. Approximately 50 ng of the purified DNA was labeled with [α - 32 P]dCTP (Amersham Co., Ltd.) using the Multiprime™ DNA Labeling System (Amersham Co., Ltd.) according to the instructions provided by the manufacturer.

[0095] (3) Screening by Plaque Hybridization

[0096] The plaques on the five plates obtained in Example 4 (1) were transferred onto nylon filters (MSI Co., Ltd.), alkali-denatured, and fixed by heating at 90° C. for 3 hours, according to the methods described in the book by Maniatis et al. The labeled DNA probe prepared in Example 4 (2) was added to the filters and hybridization was carried out according to the method described in the book by Maniatis et al. At the final step, the filters were washed with 0.1×SSC (15 mM sodium chloride, 1.5 mM sodium citrate, pH 7.0) at 60° C., and were analyzed by autoradiography to search for positive clones. As a result, 11 positive clones were obtained. One of the clones was selected, and according to the methods described in the book by Maniatis et al., phages were multiplied and DNA was extracted from them.

[0097] (4) Subcloning into Plasmid Vectors

[0098] About 5 μ g of the above-mentioned phage DNA was dissolved in 20 μ l of H buffer, and 10 units of the restriction enzyme BamHI (Takara Shuzo Co., Ltd.) was added. The reaction was carried out at 30° C. for 2 hours. After separation of the reaction products by electrophoresis through a 0.8% GTG agarose gel (Takara Shuzo Co., Ltd.), a portion containing the inserted DNA fragment of about 1.9 kb was cut out. The DNA fragment was extracted and purified using the SUPREC™-01 (Takara Shuzo Co., Ltd.) according to the instructions provided by the manufacturer.

[0099] The obtained DNA fragment was dissolved in 10 μ l of TE buffer and 0.2 μ g of pUC18 BamHI BAP (Pharmacia Co., Ltd.) was added. Ligation reaction was carried out at 16° C. for 30 minutes using the DNA Ligation Kit (Takara Shuzo Co., Ltd.) according to the instructions provided by the manufacturer. The volume of the reaction mixture was 60 μ l. Highly competent cells of *E. coli* JM109 (Toyobo Co., Ltd.) were transformed with 2 μ l of the reaction mixture according to the instructions provided by the manufacturer. The cells were cultured at 37° C. for 20 hours on X-gal ampicillin LB agar medium according to the method described in the book by Maniatis et al. One of the formed white colonies was isolated and cultured, and plasmid DNA was extracted from the culture and purified. The obtained plasmid was named pEAK14.

[0100] (5) Determination of DNA Sequence

[0101] The nucleotide sequence of about 1.9 kb which was contained in the plasmid pEAK14 and derived from petunia cDNA was determined by the Model 373A DNA Sequencing System (ABI Co., Ltd.) using the Deletion Kit for Kilosequence (Takara Shuzo Co., Ltd.) and the Taq Dideoxy™ Terminator Cycle Sequencing Kit (ABI Co., Ltd.) according

to the instructions provided by the manufacturers. The sequence was analyzed using a sequence analysis software, DNASIS™ (Hitachi Software Engineering Co., Ltd.).

[0102] As a result, the DNA sequence of 1824 bp shown by SEQ ID NO: 1 was obtained. This sequence contained an open reading frame starting at the 116th nucleotide and ending at the 1633rd nucleotide, and coding for a polypeptide consisting of 506 amino acid residues. The amino acid sequence of the polypeptide showed approximately 33% homology to that of cytochrome P450 of avocado which had been reported (Proc. Natl. Acad. Sci. USA, vol.87, p.3904-3908, 1990).

[0103] This open reading frame was named AK14 sequence.

EXAMPLE 5

[0104] Introduction of the AK14 Sequence into Plant Expression Vectors

[0105] (1) Deletion of ATG Sequence in 5' Non-coding Region

[0106] An aliquot of 2 μ g of the plasmid pEAK14 obtained in Example 4 (4) was dissolved in 20 μ l of H buffer, and 10 units of the restriction enzyme BamHI (Takara Shuzo Co., Ltd.) was added. The reaction was carried out at 30° C. for 2 hours. The reaction products were separated by electrophoresis through a 0.8% GTG agarose gel (Takara Shuzo Co., Ltd.), and a portion containing the inserted DNA fragment of about 1.9 kb was cut out. Then, the DNA fragment was extracted and purified using the SUPREC™-01 (Takara Shuzo Co., Ltd.) according to the instructions provided by the manufacturer.

[0107] The obtained DNA fragment was dissolved in 50 μ l of BAL31 buffer [20 mM Tris hydrochloride buffer (pH 8.0), 600 mM sodium chloride, 12 mM calcium chloride, 12 mM magnesium chloride, 1 mM EDTA], and one unit of BAL31 nuclease S (Takara Shuzo Co., Ltd.) was added. The reaction was carried out at 30° C. for one minute, followed by addition of 5 μ l of phenol:chloroform (1:1) mixture to terminate the reaction. After addition of 150 μ l of ethanol, the reaction mixture was allowed to stand at -80° C. for 10 minutes, and then centrifuged at 10,000×g for 10 minutes. The obtained precipitate was washed with 200 μ l of 70% ethanol, and dried under vacuum. The obtained DNA was dissolved in 50 μ l of the Klenow buffer [50 mM Tris hydrochloride buffer (pH 7.5), 10 mM magnesium chloride, 1 mM dithiothreitol, 100 μ M dATP, 100 μ M dCTP, 100 μ M dGTP, 100 μ M dTTP], and one unit of Klenow fragment (Takara Shuzo Co., Ltd.) was added. The reaction was carried out at 30° C. for 30 minutes. After addition of 150 μ l of ethanol, the reaction mixture was allowed to stand at -80° C. for 10 minutes, and then centrifuged at 10,000×g for 10 minutes. The obtained precipitate was washed with 200 μ l of 70% ethanol, and dried under vacuum. The obtained DNA was dissolved in 10 μ l of TE buffer.

[0108] (2) Subcloning into Plasmid Vectors

[0109] An aliquot of 1 μ g of pUC19 (Pharmacia Co., Ltd.) was dissolved in 50 μ l of Sma buffer [10 mM Tris hydrochloride buffer (pH 7.5), 10 mM magnesium chloride, 1 mM dithiothreitol, 20 mM potassium chloride], and 10 units of the restriction enzyme SmaI (Takara Shuzo Co., Ltd.) was

added. The reaction was carried out at 30° C. for 2 hours. After addition of 150 μ l of ethanol, the reaction mixture was allowed to stand at -80° C. for 10 minutes, and then centrifuged at 10,000 \times g for 10 minutes. The obtained precipitate was washed with 200 μ l of 70% ethanol, and dried under vacuum. The obtained DNA was dissolved in 50 μ l of CIP buffer [50 mM Tris hydrochloride buffer (pH 9.0), 1 mM magnesium chloride, 0.1 mM zinc chloride, 1 mM spermidine], and 0.1 unit of calf intestine alkaline phosphatase (Boehringer Mannheim GmbH) was added. The reaction was carried out at 37° C. for 30 minutes, and then at 56° C. for 30 minutes, followed by addition of 5 μ l of phenol:chloroform (1:1) mixture to terminate the reaction. After addition of 150 μ l of ethanol, the reaction mixture was allowed to stand at -80° C. for 10 minutes, and then centrifuged at 10,000 \times g for 10 minutes. The obtained precipitate was washed with 200 μ l of 70% ethanol, and dried under vacuum. The obtained DNA was dissolved in 10 μ l of TE buffer [10 mM Tris hydrochloride buffer (pH 7.5), 1 mM EDTA] to obtain a vector DNA solution.

[0110] An aliquot of 1 μ l of the above-mentioned vector DNA solution and 2 μ l of the DNA solution obtained in Example 5 (1) were mixed and subjected to ligation at 16° C. for 30 minutes using the DNA Ligation Kit (Takara Shuzo Co., Ltd.) according to the instructions provided by the manufacturer. The volume of the reaction mixture was 18 μ l. Highly competent cells of *E. coli* JM109 (Toyobo Co., Ltd.) were transformed with 2 μ l of the reaction mixture according to the instructions provided by the manufacturer. The cells were cultured at 37° C. for 20 hours on X-gal ampicillin LB agar medium according to the method described in the book by Maniatis et al. One of the formed white colonies was isolated and cultured, and plasmid DNA was extracted from the culture and purified. The obtained plasmid was named pEAK14S.

[0111] The nucleotide sequence of the region bound to the SmaI site derived from pUC19 vector in pEAK14S was analyzed. As a result, it was shown that the sequence of pEAK14S lacked the 1st to the 91st nucleotides of the sequence shown by SEQ ID NO: 1. It was also revealed that the direction of the insertion was such that the BamHI site of pUC19 vector was linked to the amino terminus of the AK14 sequence.

[0112] (3) Subcloning into a Plant Expression Vector, pBI121

[0113] An aliquot of 1 μ g of pEAK14S was dissolved in 50 μ l of M buffer, and 10 units of the restriction enzyme SacI (Takara Shuzo Co., Ltd.) and 10 units of the restriction enzyme XbaI (Takara Shuzo Co., Ltd.) were added. The reaction was carried out at 37° C. for 2 hours. The reaction products were separated by electrophoresis through a 0.8% GTG agarose gel (Takara Shuzo Co., Ltd.), and a portion containing the inserted DNA fragment of about 1.9 kb was cut out. Then, the DNA fragment was extracted and purified using the SUPREC™-01 (Takara Shuzo Co., Ltd.) according to the instructions provided by the manufacturer. The obtained DNA fragment was dissolved in 10 μ l of TE buffer.

[0114] Similarly, an aliquot of 1 μ g of a plant expression vector, pBI121 (GUS Gene Fusion System: Clonetech Co., Ltd.) was dissolved in 50 μ l of M buffer, and 10 units of the restriction enzyme SacI (Takara Shuzo Co., Ltd.) and 10 units of the restriction enzyme XbaI (Takara Shuzo Co.,

Ltd.) were added. The reaction was carried out at 37° C. for 2 hours. The reaction products were separated by electrophoresis through a 0.8% GTG agarose gel (Takara Shuzo Co., Ltd.), and a portion containing the vector DNA fragment of about 11 kb was cut out. Then, the vector DNA fragment was extracted and purified using the SUPREC™-01 (Takara Shuzo Co., Ltd.) according to the instructions provided by the manufacturer. The obtained vector DNA fragment was dissolved in 10 μ l of TE buffer.

[0115] An aliquot of 1 μ l of the above-mentioned TE buffer containing the AK14 DNA fragment of about 1.9 kb and 1 μ l of the above-mentioned TE buffer containing the vector DNA fragment of about 11 kb were mixed, and subjected to ligation at 16° C. for 30 minutes using the DNA Ligation Kit (Takara Shuzo Co., Ltd.) according to the instructions provided by the manufacturer. The volume of the reaction mixture was 12 μ l. Highly competent cells of *E. coli* JM109 (Toyobo Co., Ltd.) were transformed with 2 μ l of the reaction mixture according to the instructions provided by the manufacturer. The cells were cultured at 37° C. for 20 hours on kanamycin LB agar medium [1% Bacto Tryptone (Difco Laboratories), 0.5% Bacto Yeast Extract (Difco Laboratories), 1% sodium chloride, 50 μ g/ml kanamycin, 1.5% Bacto Agar (Difco Laboratories)] according to the method described in the book by Maniatis et al. One of the formed colonies was isolated and cultured, and plasmid DNA was extracted from the culture and purified. The obtained plasmid was named pBAK14.

[0116] (4) Introduction of pBAK14 into *Agrobacterium tumefaciens* LBA4404

[0117] The plasmid pBAK14 was introduced into *Agrobacterium tumefaciens* LBA4404 by triparental mating using the GUS Gene Fusion System (Clonetech Co., Ltd.) according to the instructions provided by the manufacturer. *E. coli* JM109 strain which carries pBAK14 and *E. coli* HB101 strain which carries pRK2013 (Clonetech Co., Ltd.) were cultured, respectively, in 1 ml of kanamycin LB liquid medium [1% Bacto Tryptone (Difco Laboratories), 0.5% Bacto Yeast Extract (Difco Laboratories), 1% sodium chloride, 50 μ g/ml kanamycin] with shaking at 37° C. for 12 hours. Separately, *Agrobacterium tumefaciens* LBA4404 which carries pAL4404 (Clonetech Co., Ltd.) was cultured in 1 ml of streptomycin LB liquid medium [1% Bacto Tryptone (Difco Laboratories), 0.5% Bacto Yeast Extract (Difco Laboratories), 1% sodium chloride, 300 μ g/ml streptomycin] with shaking at 28° C. for 36 hours. Three types of cultured cells were individually collected by centrifugation at 5,000 \times g for 10 minutes, washed with 1 ml of water, and suspended in small amount of water. The suspensions were mixed together, and the whole of the combined suspension was spread on LB agar medium and incubated at 28° C. for 20 hours. The obtained cells were applied on LB agar medium containing 50 μ g/ml kanamycin and 300 μ g/ml streptomycin, and incubated at 28° C. over 2 nights. One of the formed colonies was isolated to obtain *Agrobacterium tumefaciens* LBA4404 carrying both pBAK14 and pAL4404.

EXAMPLE 6

[0118] Introduction of the AK14 Sequence into Tobacco and Its Expression

[0119] (1) Introduction into Tobacco Using a Microorganism of the Genus *Agrobacterium*

[0120] *Agrobacterium tumefaciens* LBA 4404 strain carrying pBAK14 and pAL4404 which was obtained in Example 5 was cultured in 10 ml of LB liquid medium [1% Bacto Tryptone (Difco Laboratories), 0.5% Bacto Yeast Extract (Difco Laboratories), 1% sodium chloride] containing 50 $\mu\text{g/ml}$ kanamycin and 300 $\mu\text{g/ml}$ streptomycin, with shaking at 28° C. for 40 hours. The cultured cells were collected by centrifugation at 5,000 \times g for 10 minutes, washed with 10 ml of water, and then suspended in an equal amount of water.

[0121] Leaves of tobacco (*Nicotiana tabacum* cv. petit Havana SR-1) aseptically subcultured at 25° C. were cut into one centimeter squares, soaked in the above-mentioned cell suspension, and wiped with sterilized filter paper. The leaves were placed on MS medium containing 1 $\mu\text{g/ml}$ 6-benzyladenine, 0.3 $\mu\text{g/ml}$ 1-naphthaleneacetic acid, 3% sucrose, and 0.2% Gelrite (Physiol. Plant., vol.15, p.473-497, 1962) (hereinafter referred to as the solid PD4 medium) with the abaxial side up, and cultured at 25° C. for 2 days under continuous light illumination at 2,500 lux. Then, the leaves were transplanted to the solid PD4 medium containing 500 $\mu\text{g/ml}$ Claforan (for injection, Hoechst Japan Co., Ltd.) and 200 $\mu\text{g/ml}$ kanamycin for culturing, and transplanted to the same medium every 2 weeks afterward. About one month after the start of culturing, adventitious buds were induced. The buds were cut off and subcultured on MS medium containing 500 $\mu\text{g/ml}$ Claforan and 50 $\mu\text{g/ml}$ kanamycin to induce rooting. Plants which took roots were transferred into pots, after checked for their aseptic condition, and cultivated at 25° C. in an artificial weather system. Transgenic plants were thus obtained.

[0122] (2) Detection of Enzyme Activity in Leaves of Tobacco which had been Transformed (hereinafter Referred to as the Transgenic Tobacco)

[0123] Microsome fraction was prepared from 20 g of the transgenic tobacco leaves obtained as above according to the method described in Example 2 (3), and flavonoid-3',5'-hydroxylase activity in the fraction was determined. As a control, microsome fraction prepared from non-transgenic tobacco leaves was used. As a result, said enzyme activity, which catalyzes the conversion of dihydroquercetin to dihydromyricetin, was detected only in the microsome fraction of the transgenic tobacco.

[0124] (3) Change in Pigments in Petals of the Transgenic Tobacco

[0125] Anthocyanidins were prepared from petals of the transgenic and non-transgenic tobacco plants, respectively, according to the method described in Example 2 (1), and analyzed. As a result, only cyanidin was detected in the non-transgenic tobacco, whereas cyanidin and delphinidin were detected in almost the same amounts in the transgenic tobacco.

[0126] The flower colors were compared with The Japan Color Standard For Horticultural Plants (Japan Color Research Institute). The color of flowers of the transgenic

tobacco corresponded to Color No. 8904 or 8905, and that of the non-transgenic tobacco corresponded to Color No. 9503 or 9504. That is, flowers of the transgenic tobacco showed more bluish color.

EXAMPLE 7

[0127] Introduction of the AK14 Sequence into a Petunia Cultivar with Pink Glowlers and its Expression

[0128] (1) Introduction into Petunia Using a Microorganism of the Genus *Agrobacterium*

[0129] Kanamycin-resistant transgenic plants were obtained by infecting leaves of aseptically subcultured petunia (*Petunia hybrida* cv. Falcon Pinkvein: Sakata Seed Corporation) with *Agrobacterium tumefaciens* LBA4404 strain which carries pBAK14 and pAL4404 according to a method similar to that used in Example 6.

[0130] (2) Change in Pigments in Petals of the Transgenic Petunia

[0131] Anthocyanidins were prepared from petals of the above-mentioned transgenic petunia according to the method described in Example 2 (1), and compared with those prepared from the control, non-transgenic petunia (Falcon Pinkvein). As a result, little malvidin or delphinidin was detected in the non-transgenic petunia. On the other hand, the transgenic petunia had both of them as major components. The major component in the control plants was peonidin.

[0132] The flower colors at the center area of petals were compared with The Japan Color Standard For Horticultural Plants (Japan Color Research Institute). The color of flowers of the transgenic petunia corresponded to Color No. 9206 or 9207, and that of the non-transgenic petunia (Falcon Pinkvein) corresponded to Color No. 9204 or 9205. That is, flowers of the transgenic petunia showed more bluish color.

EXAMPLE 8

[0133] Introduction of the AK14 Sequence into Rose and its Expression

[0134] (1) Introduction into Rose Using a Microorganism of the Genus *Agrobacterium*

[0135] Leaves of aseptically subcultured rose (*Rosa hybrida* cv. deep red) were infected with *Agrobacterium tumefaciens* LBA4404 strain carrying pBAK14 and pAL4404 according to a method similar to that used in Example 6 (1). The leaves were placed on MS medium containing 0.01 $\mu\text{g/ml}$ 6-benzyladenine, 10 $\mu\text{g/ml}$ 2,4-dichlorophenoxyacetic acid, 3% sucrose, and 0.2% Gelrite (hereinafter referred to as the solid BE medium), and cultured at 25° C. for 2 days under continuous light illumination at 2,500 lux. Then, the leaves were transplanted to the solid BE medium containing 500 $\mu\text{g/ml}$ Claforan, and after 7 days, transplanted to the solid BE medium containing 500 $\mu\text{g/ml}$ Claforan and 200 $\mu\text{g/ml}$ kanamycin. Thereafter, the leaves were transplanted to the same medium every 2 weeks. After about 2 months, approximately 20 g of kanamycin-resistant callus was obtained.

[0136] (2) Expression of Enzyme Activity in the Rose Callus

[0137] Microsome fraction was prepared from the callus obtained in Example 8 (1) according to the method described in Example 2 (3), and flavonoid-3',5'-hydroxylase activity in the fraction was determined. As a control, microsome fraction prepared from untransformed callus of rose was used. As a result, said enzyme activity, which catalyzes the conversion of dihydroquercetin to dihydromyricetin, was detected only in the microsome fraction of the transformed callus.

EXAMPLE 9

[0138] Introduction of the AK14 Sequence into Carnation and its Expression

[0139] (1) Introduction of pBAK14 into *Agrobacterium rhizogenes* NIAES1724 Strain

[0140] According to a method similar to that described in Example 5 (4), pBAK14 was introduced into *Agrobacterium rhizogenes* NIAES1724 strain (obtained from National Institute of Agrobiological Resources, the Japanese Ministry of Agriculture, Forestry and Fisheries). In this example, JM103 was used as the *E. coli* strain, and 25 $\mu\text{g/ml}$ nalidixic acid (Sigma Co., Ltd.) was used instead of streptomycin.

[0141] (2) Introduction of the AK14 Sequence into Carnation Using a Microorganism of the Genus *Agrobacterium*

[0142] Petals cut off from buds of carnation (*Dianthus caryophyllus* cv. Nora) were infected with *Agrobacterium rhizogenes* NIAES1724 carrying pBAK14 according to a method similar to that described in Example 6 (1). The infected petals were placed on solid MS medium containing 0.3 $\mu\text{g/ml}$ 6-benzyladenine, 0.3 $\mu\text{g/ml}$ naphthaleneacetic acid, 3% sucrose, and 0.2% Gelrite, and cultured at 25° C. for 3 days under continuous light illumination at 2,500 lux. Then, the petals were transplanted to the same medium containing 250 $\mu\text{g/ml}$ Claforan, and after 7 days, transplanted to the same medium containing 250 $\mu\text{g/ml}$ Claforan and 300 $\mu\text{g/ml}$ kanamycin. Thereafter, the petals were transplanted to the same medium every 2 weeks. After about 4 months, approximately 10 g of kanamycin-resistant hairy roots were obtained.

[0143] (3) Expression of Enzyme Activity in Hairy Roots of Carnation

[0144] Microsome fraction was prepared from the hairy roots obtained in Example 8 (1) according to the method described in Example 2 (3), and flavonoid-3',5'-hydroxylase activity in the fraction was determined. As a control, microsome fraction prepared from hairy roots infected with *Agrobacterium rhizogenes* NIAES1724 strain which did not carry pBAK14 was used. As a result, said enzyme activity, which catalyzes the conversion of dihydroquercetin to dihydromyricetin, was detected only in the microsome fraction of the transformed hairy roots.

EXAMPLE 10

[0145] Detection of AK14 Homologous Sequences in Genomic DNAs of Heterogeneous Plants

[0146] (1) Preparation of Plant Genomic DNA

[0147] Ten to twenty grams of green leaves of each of the following plants was freeze-dried, and their genomic DNAs were extracted according to the method described in DNA Cloning A Practical Approach, vol.2, p.103, 1985, IRL Press: petunia (*Petunia hybrida* cv. Purple Joy: NPI Seeds), nicotiana (*Nicotiana affinis* cv. F1 Domino: Daiichi Seed Co., Ltd.), Japanese gentian (*Gentiana triflora* cv. Japonica), sweet pea (*Lathyrus odoratus* cv. Royal Deep Blue: Daiichi Seed Co., Ltd.), pansy (*Viola tricolor*; blue cultivar), primrose (*Primula polyantha*, purple cultivar), Russell prairie gentian (*Eustoma russellianum* cv. Royal Light Purple: Takii Seed Co., Ltd.), campanula (*Campanula medium*, light purple cultivar), delphinium (*Delphinium hybridum*, pale blue cultivar), and hyacinth (*Hyacinthus orientalis*, purple cultivar).

[0148] (2) Preparation of Genomic DNA Blots

[0149] An aliquot of 5 μg of each of the genomic DNAs obtained in Example 10 (1) was dissolved in 20 μl of H buffer, and 10 units of the restriction enzyme EcoRV (Takara Shuzo Co., Ltd.) was added. The reaction was carried out at 37° C. for 2 hours. According to the method described in the book by Maniatis et al, the digested DNA was subjected to 0.8% agarose gel electrophoresis, alkali-denatured, and neutralized. Then, the DNA was transferred onto nylon filters (MSI Co., Ltd.), and fixed by heating at 90° C. for 3 hours for fixation to prepare genomic DNA blots.

[0150] (3) Radiolabeling of AK14 Sequence Probe

[0151] An aliquot of 1 μg of pEAK14 obtained in Example 4 was dissolved in 20 μl of H buffer, and 10 units of the restriction enzyme BamHI (Takara Shuzo Co., Ltd.) was added. The reaction was carried out at 37° C. for 2 hours. The reaction products were separated by electrophoresis through a 0.8% GTG agarose gel (Takara Shuzo Co., Ltd.), and a portion containing the inserted DNA fragment of about 1.9 kb was cut out. The inserted DNA fragment was extracted and purified using the SUPREC™-01 (Takara Shuzo Co., Ltd.) according to the instructions provided by the manufacturer. An aliquot of 50 ng of the DNA fragment containing the AK14 sequence was labeled with [α -³²P] dCTP (Amersham Co., Ltd.) using the Multiprime™ DNA Labeling System (Amersham Co., Ltd.) according to the instructions provided by the manufacturer.

[0152] (4) Hybridization

[0153] The genomic DNA blots obtained in Example 10 (2) were hybridized with the labeled probe of (3) according to the method described in the book by Maniatis et al. At the final step, the filters were washed twice with 2 \times SSC (0.3 M sodium chloride, 0.03 M sodium citrate, pH 7.0) at 50° C. for 30 minutes. The obtained filters were examined by autoradiography using X-ray films (New RX: Fuji Photo Film Co., Ltd.). As a result, the DNAs prepared from petunia (Purple Joy), nicotiana, Japanese gentian, Russell prairie gentian, and campanula showed a clear band. The DNAs prepared from sweet pea and primrose showed a band hybridized with the probe though unclear. That is, the result

showed that homologous sequences which hybridize with the AK14 sequence existed in the genomic DNAs of these plants.

EXAMPLE 11

[0154] Detection of AK14 Homologous Sequences in Petal cDNAs of Heterogeneous Plants

[0155] (1) Preparation of Petal cDNA

[0156] About 10 g of petals was collected from buds of each of the following plants; petunia (*Petunia hybrida* cv. Purple Joy: NPI Seeds Co., Ltd.), nicotiana (*Nicotiana affinis* cv. F1 Domino: Daiichi Seed Co., Ltd.), Japanese gentian (*Gentiana triflora* cv. Japonica), Russell prairie gentian (*Eustoma russellianum* cv. Royal Light Purple: Takii Seed Co., Ltd.), and campanula (*Campanula medium*, light purple cultivar). mRNAs were extracted from the petals according to the method described in Example 1 (2). By using the obtained mRNAs as templates, double strand cDNAs were synthesized using the cDNA Synthesis System Plus RPN1256 (Amersham Co., Ltd.) according to the instructions provided by the manufacturer.

[0157] (2) Preparation of cDNA Blots

[0158] According to the method described in the book by Maniatis et al, about 0.1 μ g of each of the above-mentioned cDNAs was subjected to 0.8% agarose gel electrophoresis, alkali-denatured, and neutralized. Then, the cDNA was transferred onto nylon filters (MSI Co., Ltd.), and fixed by heating at 90° C. for 3 hours to prepare cDNA blots.

[0159] (3) Hybridization

[0160] A radiolabeled AK14 sequence probe was prepared according to a method similar to that used in Example 10 (3), and hybridized with each of the above-mentioned cDNA blots according to a method similar to that used in Example 10 (4). At the final step, the filters were washed twice with 2 \times SSC at 50° C. for 30 minutes, and then examined by autoradiography. As a result, each plant showed a clear band at the location corresponding to about 2 kb. That is, it was demonstrated that analogous sequences which hybridize with the AK14 sequence existed in the petal cDNAs of these plants.

EXAMPLE 12

[0161] Cloning of the AK14 Homologous Sequence from Russell Prairie Gentian and Campanula

[0162] (1) Construction of Petal cDNA Library

[0163] About 20 g of petals was collected from buds of Russell prairie gentian (*Eustoma russellianum* cv. Royal Light Purple: Takii Seed Co., Ltd.) and campanula (*Campanula medium*, light purple cultivar), and mRNAs were extracted from them, respectively, according to the method described in Example 1 (2). By using the obtained mRNAs as templates, double strand cDNAs were synthesized and cloned into λ gt122 vectors using the SuperscriptTM Lambda System (BRL Life Technologies Co., Ltd.) according to the instructions provided by the manufacturer.

[0164] Each final product was subjected to the packaging reaction using the XDNA in vitro packaging kit Giga Pack Gold (Stratagene Co., Ltd.) according to the instructions provided by the manufacturer. Cells of *E. coli* Y1090 (τ^-)

(BRL Life Technologies Co., Ltd.) were infected with the appropriately diluted packaging products according to the instructions provided by the manufacturer, and spread on LB agar medium [1% Bacto Tryptone (Difco Laboratories), 0.5% Bacto Yeast Extract (Difco Laboratories), 1% sodium chloride, 1.5% Bacto Agar (Difco Laboratories)] in plastic plates of 15 cm in diameter (Iwaki Glass Co., Ltd.) to obtain about 10,000 plaques per plate. Five plates were prepared for Russell prairie gentian and campanula, respectively, to obtain cDNA libraries.

[0165] (2) Screening by Plaque Hybridization

[0166] The plaques on the five plates obtained in Example 4 (1) were transferred onto nylon filters (MSI Co., Ltd.), alkali-denatured, and fixed by heating at 90° C. for 3 hours according to the methods described in the book by Maniatis et al. The radiolabeled probe DNA prepared by a method similar to that used in Example 11 (3) was added to the filters and hybridization was carried out according to the method described in the book by Maniatis et al. At the final step, the filters were washed with 2 \times SSC (0.3 M sodium chloride, 0.03 M sodium citrate, pH 7.0) at 50° C., and were examined by autoradiography to search for positive clones. As a result, 12 and 7 positive clones were obtained from the library of Russell prairie gentian and that of campanula, respectively. One clone was selected from each library, and according to the method described in the book by Maniatis et al, phages were multiplied and DNAs were extracted from them.

[0167] About 5 μ g of each phage DNA was dissolved in 20 μ l of H buffer, and 10 units of the restriction enzyme NotI (Takara Shuzo Co., Ltd.) and 10 units of the restriction enzyme Sall (Takara Shuzo Co., Ltd.) were added. The reaction was carried out at 37° C. for 2 hours. The reaction products were separated by electrophoresis through a 0.8% GTG agarose gel (Takara Shuzo Co., Ltd.), and a portion containing the inserted DNA fragment of about 2 kb was cut out from each gel. The DNA fragments were extracted and purified using the SUPRECTM-01 (Takara Shuzo Co., Ltd.) according to the instructions provided by the manufacturer, and dissolved in 10 μ l of TE buffer, respectively.

[0168] (3) Subcloning into Plasmid Vectors

[0169] About 1 μ g of DNA of a plasmid vector, pBlue-scriptIIKS+ (Stratagene Co., Ltd.) was dissolved in 20 μ l of H buffer, and 10 units of the restriction enzyme NotI (Takara Shuzo Co., Ltd.) and 10 units of the restriction enzyme Sall (Takara Shuzo Co., Ltd.) were added. The reaction was carried out at 37° C. for 2 hours. The reaction products were separated by electrophoresis through a 0.8% GTG agarose gel (Takara Shuzo Co., Ltd.), and a portion containing the vector DNA fragment of about 3 kb was cut out. The DNA fragment was extracted and purified using the SUPRECTM-01 (Takara Shuzo Co., Ltd.) according to the instructions provided by the manufacturer, and dissolved in 10 μ l of TE buffer.

[0170] To 4 μ l each of the two types of inserted DNA fragments obtained in Example 12 (2) was added 1 μ l of the above-mentioned vector DNA fragment, respectively, and ligation was carried out at 16° C. for 30 minutes using the DNA Ligation Kit (Takara Shuzo Co., Ltd.) according to the instructions provided by the manufacturer. The volume of each reaction mixture was 30 μ l. Highly competent cells of *E. coli* JM109 (Toyobo Co., Ltd.) were transformed with 2

μ l each of the reaction mixtures, respectively, according to the instructions provided by the manufacturer. The transformed cells were cultured at 37° C. for 20 hours on X-gal ampicillin LB agar medium according to the method described in the book by Maniatis et al. From each culture, one of the white colonies formed was isolated and cultured, and plasmid DNA was extracted from the culture and purified. The plasmid derived from the library of Russell prairie gentian was named pETg1, and that from the library of campanula was named pEKa1.

[0171] (4) Determination of DNA Sequence

[0172] The nucleotide sequences of the DNA fragments which were derived from the petal cDNAs and contained in the plasmids pETg1 and pEKa1 were determined by the Model 373A DNA Sequencing System (ABI Co., Ltd.) using the Deletion Kit for Kilosequence (Takara Shuzo Co., Ltd.) and the Taq Dideoxy™ Terminator Cycle Sequencing Kit (ABI Co., Ltd.) according to the instructions provided by manufacturers. The sequences were analyzed using a sequence analysis software, DNASIS™ (Hitachi Software Engineering Co., Ltd.).

[0173] As a result, the DNA sequence of 2174 bp shown by SEQ ID NO: 63 was obtained from Russell prairie gentian. This sequence contained an open reading frame starting at the 92nd nucleotide and ending at the 1621st nucleotide, and coding for a polypeptide consisting of 510 amino acid residues. The amino acid sequence of the polypeptide showed 74% homology to that of AK14. This open reading frame was named Tg1 sequence.

[0174] The DNA sequence of 1927 bp shown by SEQ ID NO: 64 was obtained from campanula. This sequence contained an open reading frame starting at the 180th nucleotide and ending at the 1748th nucleotide, and coding for a polypeptide consisting of 523 amino acid residues. The amino acid sequence of the polypeptide showed 66% homology to that of AK14. This open reading frame was named Ka1 sequence.

EXAMPLE 13

[0175] Introduction of Tg1 and Ka1 into Plant Expression Vectors

[0176] (1) Subcloning into Plant Expression Vector pBI121

[0177] An aliquot of 1 μ g of pETg1 was dissolved in 50 μ l of H buffer, and 10 units of the restriction enzyme Sall (Takara Shuzo Co., Ltd.) was added. The reaction was carried out at 37° C. for 2 hours. After addition of 150 μ l of ethanol, the reaction mixture was allowed to stand at -80° C. for 10 minutes, and then centrifuged at 10,000 \times g for 10 minutes. The obtained precipitate was washed with 200 μ l of 70% ethanol, and dried under vacuum. The obtained DNA was dissolved in 50 μ l of Klenow buffer, and one unit of Klenow fragment (Takara Shuzo Co., Ltd.) was added. The reaction was carried out at 30° C. for 30 minutes. After addition of 150 μ l of ethanol, the reaction mixture was allowed to stand at -80° C. for 10 minutes, and then centrifuged at 10,000 \times g for 10 minutes. The obtained precipitate was washed with 200 μ l of 70% ethanol, and dried under vacuum. The obtained DNA was dissolved in 50 μ l of M buffer, and 10 units of the restriction enzyme SacI (Takara Shuzo Co., Ltd.) was added. The reaction was carried out at

37° C. for 2 hours. The reaction products were separated by electrophoresis through a 0.8% GTG agarose gel (Takara Shuzo Co., Ltd.), and a portion containing the inserted DNA fragment of about 2.2 kb was cut out. The DNA fragment was extracted and purified using the SUPREC™-01 (Takara Shuzo Co., Ltd.) according to the instructions provided by the manufacturer, and dissolved in 10 μ l of TE buffer.

[0178] Separately, 1 μ g of pEKa1 was dissolved in 50 μ l of H buffer, and 10 units of the restriction enzyme Sall (Takara Shuzo Co., Ltd.) was added. The reaction was carried out at 37° C. for 2 hours. After addition of 150 μ l of ethanol, the reaction mixture was allowed to stand at -80° C. for 10 minutes, and then centrifuged at 10,000 \times g for 10 minutes. The obtained precipitate was washed with 200 μ l of 70% ethanol, and dried under vacuum. The obtained DNA was dissolved in 50 μ l of Klenow buffer, and one unit of Klenow fragment (Takara Shuzo Co., Ltd.) was added. The reaction was carried out at 30° C. for 30 minutes. After addition of 150 μ l of ethanol, the reaction mixture was allowed to stand at -80° C. for 10 minutes, and then centrifuged at 10,000 \times g for 10 minutes. The obtained precipitate was washed with 200 μ l of 70% ethanol, and dried under vacuum. The obtained DNA was dissolved in 50 μ l of M buffer, and 0.5 unit of the restriction enzyme SacI (Takara Shuzo Co., Ltd.) was added. The reaction was carried out at 37° C. for one hour. The reaction products were separated by electrophoresis through a 0.8% GTG agarose gel (Takara Shuzo Co., Ltd.), and a portion containing the inserted DNA fragment of about 1.9 kb was cut out. The DNA fragment was extracted and purified using the SUPREC™-01 (Takara Shuzo Co., Ltd.) according to the instructions provided by the manufacturer, and dissolved in 10 μ l of TE buffer.

[0179] An aliquot of 1 μ g of the plant expression vector pBI121 (GUS Gene Fusion System: Clontech Co., Ltd.) was dissolved in 50 μ l of Sma buffer [10 mM Tris hydrochloride buffer (pH 7.5), 10 mM magnesium chloride, 1 mM dithiothreitol, 20 mM potassium chloride], and 10 units of the restriction enzyme SmaI (Takara Shuzo Co., Ltd.) was added. The reaction was carried out at 30° C. for 2 hours. After addition of 150 μ l of ethanol, the reaction mixture was allowed to stand at -80° C. for 10 minutes, and then centrifuged at 10,000 \times g for 10 minutes. The obtained precipitate was washed with 200 μ l of 70% ethanol, and dried under vacuum. The obtained DNA was dissolved in 50 μ l of M buffer, and 10 units of the restriction enzyme SacI (Takara Shuzo Co., Ltd.) was added. The reaction was carried out at 37° C. for 2 hours. The reaction products were separated by electrophoresis through a 0.8% GTG agarose gel (Takara Shuzo Co., Ltd.), and a portion containing the vector DNA fragment of about 11 kb was cut out. The vector DNA fragment was extracted and purified using the SUPREC™-01 (Takara Shuzo Co., Ltd.) according to the instructions provided by the manufacturer, and dissolved in 10 μ l of TE buffer.

[0180] An aliquot of 1 μ l of the TE buffer containing the vector DNA fragment and 1 μ l of the TE buffer containing the DNA insert fragment of pETg1 were mixed, and ligation was carried out at 16° C. for 30 minutes using the DNA Ligation Kit (Takara Shuzo Co., Ltd.) according to the instructions provided by the manufacturer. The volume of

the reaction mixture was 12 μ l. Highly competent cells of *E. coli* JM109 (Toyobo Co., Ltd.) were transformed with 2 μ l of the reaction mixture according to the instructions provided by the manufacturer. The transformed cells were cultured at 37° C. for 20 hours on kanamycin LB agar medium [1% Bacto Tryptone (Difco Laboratories), 0.5% Bacto Yeast Extract (Difco Laboratories), 1% sodium chloride, 50 μ g/ml kanamycin, 1.5% Bacto Agar (Difco Laboratories)] according to the method described in the book by Maniatis et al. One of the formed colonies was isolated and cultured, and plasmid DNA was extracted and purified. The obtained plasmid was named pBTg1. pBTg1 is a plasmid composed of the plant expression vector pBI121, and inserted therein, Tg1, which is the AK14 homologous cDNA sequence derived from Russell prairie gentian.

[0181] An aliquot of 1 μ l of the TE buffer containing the vector DNA fragment and 1 μ l of the TE buffer containing the inserted DNA fragment of pEKa1 were mixed, and ligation was carried out at 16° C. for 30 minutes using the DNA Ligation Kit (Takara Shuzo Co., Ltd.) according to the instructions provided by the manufacturer. The volume of the reaction mixture was 12 μ l. Highly competent cells of *E. coli* JM109 (Toyobo Co., Ltd.) were transformed with 2 μ l of the reaction mixture according to the instructions provided by the manufacturer. The transformed cells were cultured at 37° C. for 20 hours on kanamycin LB agar medium [1% Bacto Tryptone (Difco Laboratories), 0.5% Bacto Yeast Extract (Difco Laboratories), 1% sodium chloride, 50 μ g/ml kanamycin, 1.5% Bacto Agar (Difco Laboratories)] according to the method described in the book by Maniatis et al. One of the formed colonies was isolated and cultured, and plasmid DNA was extracted and purified. The obtained plasmid was named pBKa1. pBKa1 is a plasmid composed of the plant expression vector pBI121, and inserted therein, Ka1, which is the AK14 homologous cDNA sequence derived from campanula.

[0182] (2) Introduction of pBTg1 and pBKa1 into *Agrobacterium tumefaciens* LBA4404 Strain

[0183] The plasmids pBTg1 and pBKa1 were respectively introduced into *Agrobacterium tumefaciens* LBA4404 strain using the triparental mating technique described in Example 5 (4).

EXAMPLE 14

[0184] Introduction of Tg1 and Ka1 into Tobacco and Their Expression

[0185] (1) Introduction into Tobacco Using a Microorganism of the Genus *Agrobacterium*

[0186] Leaves of tobacco (*Nicotiana tabacum* cv. petit Havana SR-1) were infected with each of the two types of *Agrobacterium* strains prepared in Example 13 (2) according to a method similar to that described in Example 6 (1) to obtain kanamycin-resistant transgenic tobacco.

[0187] (2) Detection of Enzyme Activity in Leaves of Transgenic Tobacco

[0188] Microsome fractions were prepared from 20 g each of the leaves of two types of transgenic tobacco obtained as above according to the method described in Example 2 (3), and flavonoid-3',5'-hydroxylase activity in the fractions was determined. As a result, said enzyme activity, which catalyzes the conversion of dihydroquercetin to dihydromyricetin, was detected in the microsome fractions of both transgenic tobacco. On the other hand, said enzyme activity was not detected in the microsome fraction prepared from leaves of the non-transgenic tobacco.

[0189] (3) Change in Pigments in Petals of the Transgenic Tobacco

[0190] Anthocyanidins were prepared from petals of the transgenic and non-transgenic tobacco plants, respectively, according to the method described in Example 2 (1), and analyzed. As a result, only cyanidin was detected in the non-transgenic tobacco, whereas cyanidin and delphinidin were detected in almost the same amounts in both the transgenic tobacco plants.

[0191] The flower colors were compared with The Japan Color Standard For Horticultural Plants (Japan Color Research Institute). The color of flowers of the transgenic tobacco corresponded to Color No. 8904 or 8905, and that of the non-transgenic tobacco corresponded to Color No. 9503 or 9504. That is, flowers of the transgenic tobacco showed more bluish color.

[0192] Industrial Applicability

[0193] According to the present invention, a plant having a pigment pattern which flowers or fruits of the plant do not originally have can be provided.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(iii) NUMBER OF SEQUENCES: 67

(2) INFORMATION FOR SEQ ID NO: 1 :

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1824 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

-continued

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Petunia hybrida*
 (B) STRAIN: Falcon Blue

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 116 to 1633
 (C) IDENTIFICATION METHOD: by experiment

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1

GCTACTTCGT TATATATATG TAAAAATTGTG ACTTTGAAAA TCATTTAAAT TATCATAAGG60

TTCATTTTAT CTGATCAAAA ATATTTACTT CGGCCATATA CGTTTTCTT TAGTC ATG118

Met

1

ATG CTA CTT ACT GAG CTT GGT GCA GCA ACT TCA ATC TTT CTA ATA GCA 166
 Met Leu Leu Thr Glu Leu Gly Ala Ala Thr Ser Ile Phe Leu Ile Ala
 5 10 15

CAC ATA ATC ATT TCA ACT CTT ATT TCA AAA ACT ACC GGC CGG CAT CTA 214
 His Ile Ile Ile Ser Thr Leu Ile Ser Lys Thr Thr Gly Arg His Leu
 20 25 30

CCG CCG GGG CCA AGA GGG TGG CCG GTG ATC GGA GCA CTT CCA CTT TTA 262
 Pro Pro Gly Pro Arg Gly Trp Pro Val Ile Gly Ala Leu Pro Leu Leu
 35 40 45

GGA GCC ATG CCA CAT GTT TCC TTA GCT AAA ATG GCA AAA AAA TAT GGA 310
 Gly Ala Met Pro His Val Ser Leu Ala Lys Met Ala Lys Lys Tyr Gly
 50 55 60 65

GCA ATC ATG TAT CTC AAA GTT GGA ACA TGT GGC ATG GCA GTT GCT TCT 358
 Ala Ile Met Tyr Leu Lys Val Gly Thr Cys Gly Met Ala Val Ala Ser
 70 75 80

ACC CCT GAT GCT GCT AAA GCA TTC TTG AAA ACA CTT GAT ATC AAC TTC 406
 Thr Pro Asp Ala Ala Lys Ala Phe Leu Lys Thr Leu Asp Ile Asn Phe
 85 90 95

TCC AAT CGT CCA CCT AAT GCA GGT GCC ACT CAC TTA GCT TAT AAT GCT 454
 Ser Asn Arg Pro Pro Asn Ala Gly Ala Thr His Leu Ala Tyr Asn Ala
 100 105 110

CAA GAC ATG GTT TTT GCA CAT TAT GGA CCA CGA TGG AAG TTG CTA AGG 502
 Gln Asp Met Val Phe Ala His Tyr Gly Pro Arg Trp Lys Leu Leu Arg
 115 120 125

AAA TTA AGC AAC TTG CAT ATG CTA GGG GGA AAA GCC TTA GAG AAT TGG 550
 Lys Leu Ser Asn Leu His Met Leu Gly Gly Lys Ala Leu Glu Asn Trp
 130 135 140 145

GCA AAT GTT CGT GCC AAT GAG CTA GGG CAC ATG CTA AAA TCA ATG TCC 598
 Ala Asn Val Arg Ala Asn Glu Leu Gly His Met Leu Lys Ser Met Ser
 150 155 160

GAT ATG AGT CGA GAG GGC CAG AGG GTT GTG GTG GCG GAG ATG TTG ACA 646
 Asp Met Ser Arg Glu Gly Gln Arg Val Val Val Ala Glu Met Leu Thr
 165 170 175

TTT GCC ATG GCC AAT ATG ATC GGA CAA GTG ATG CTA AGC AAA AGA GTA 694
 Phe Ala Met Ala Asn Met Ile Gly Gln Val Met Leu Ser Lys Arg Val
 180 185 190

TTT GTA GAT AAA GGT GTT GAG GTA AAT GAA TTT AAG GAC ATG GTT GTA 742
 Phe Val Asp Lys Gly Val Glu Val Asn Glu Phe Lys Asp Met Val Val
 195 200 205

GAG TTA ATG ACA ATA GCA GGG TAT TTC AAC ATT GGT GAT TTT ATT CCT 790
 Glu Leu Met Thr Ile Ala Gly Tyr Phe Asn Ile Gly Asp Phe Ile Pro
 210 215 220 225

TGT TTA GCT TGG ATG GAT TTA CAA GGG ATA GAA AAA CGA ATG AAA CGT 838

-continued

Cys Leu Ala Trp Met Asp Leu Gln Gly Ile Glu Lys Arg Met Lys Arg
 230 235 240

TTA CAT AAG AAG TTT GAT GCT TTA TTG ACA AAG ATG TTT GAT GAA CAC 886
 Leu His Lys Lys Phe Asp Ala Leu Leu Thr Lys Met Phe Asp Glu His
 245 250 255

AAA GCA ACT ACC TAT GAA CGT AAG GGG AAA CCA GAT TTT CTT GAT GTT 934
 Lys Ala Thr Thr Tyr Glu Arg Lys Gly Lys Pro Asp Phe Leu Asp Val
 260 265 270

GTT ATG GAA AAT GGG GAC AAT TCT GAA GGA GAA AGA CTC AGT ACA ACC 982
 Val Met Glu Asn Gly Asp Asn Ser Glu Gly Glu Arg Leu Ser Thr Thr
 275 280 285

AAC ATC AAA GCA CTT TTG CTG AAT TTG TTC ACA GCT GGT ACG GAC ACT1030
 Asn Ile Lys Ala Leu Leu Leu Asn Leu Phe Thr Ala Gly Thr Asp Thr
 290 295 300 305

TCT TCT AGT GCA ATA GAA TGG GCA CTT GCA GAA ATG ATG AAG AAC CCT1078
 Ser Ser Ser Ala Ile Glu Trp Ala Leu Ala Glu Met Met Lys Asn Pro
 310 315 320

GCC ATT TTG AAA AAA GCA CAA GCA GAA ATG GAT CAA GTC ATT GGA AGA1126
 Ala Ile Leu Lys Lys Ala Gln Ala Glu Met Asp Gln Val Ile Gly Arg
 325 330 335

AAT AGG CGT TTA CTC GAA TCC GAT ATC CCA AAT CTC CCT TAC CTC CGA1174
 Asn Arg Arg Leu Leu Glu Ser Asp Ile Pro Asn Leu Pro Tyr Leu Arg
 340 345 350

GCA ATT TGC AAA GAA ACA TTT CGA AAA CAC CCT TCT ACA CCA TTA AAT1222
 Ala Ile Cys Lys Glu Thr Phe Arg Lys His Pro Ser Thr Pro Leu Asn
 355 360 365

CTT CCT AGG ATC TCG AAC GAA CCA TGC ATA GTC GAT GGT TAT TAC ATA1270
 Leu Pro Arg Ile Ser Asn Glu Pro Cys Ile Val Asp Gly Tyr Tyr Ile
 370 375 380 385

CCA AAA AAC ACT AGG CTT AGT GTT AAC ATA TGG GCA ATT GGA AGA GAT1318
 Pro Lys Asn Thr Arg Leu Ser Val Asn Ile Trp Ala Ile Gly Arg Asp
 390 395 400

CCC CAA GTT TGG GAA AAT CCA CTA GAG TTT AAT CCC GAA AGA TTC TTG1366
 Pro Gln Val Trp Glu Asn Pro Leu Glu Phe Asn Pro Glu Arg Phe Leu
 405 410 415

AGT GGA AGA AAC TCC AAG ATT GAT CCT CGA GGG AAC GAT TTT GAA TTG1414
 Ser Gly Arg Asn Ser Lys Ile Asp Pro Arg Gly Asn Asp Phe Glu Leu
 420 425 430

ATA CCA TTT GGT GCT GGA CGA AGA ATT TGT GCA GGA ACA AGA ATG GGA1462
 Ile Pro Phe Gly Ala Gly Arg Arg Ile Cys Ala Gly Thr Arg Met Gly
 435 440 445

ATT GTA ATG GTG GAA TAT ATA TTA GGA ACT TTG GTT CAT TCA TTT GAT1510
 Ile Val Met Val Glu Tyr Ile Leu Gly Thr Leu Val His Ser Phe Asp
 450 455 460 465

TGG AAA TTA CCA AGT GAA GTT ATT GAG TTG AAT ATG GAA GAA GCT TTT1558
 Trp Lys Leu Pro Ser Glu Val Ile Glu Leu Asn Met Glu Glu Ala Phe
 470 475 480

GGC TTA GCT TTG CAG AAA GCT GTC CCT CTT GAA GCT ATG GTT ACT CCA1606
 Gly Leu Ala Leu Gln Lys Ala Val Pro Leu Glu Ala Met Val Thr Pro
 485 490 495

AGG TTA CAA TTG GAT GTT TAT GTA CCA TAGCTATAGA TGTGTATTGT 1653
 Arg Leu Gln Leu Asp Val Tyr Val Pro
 500 505

GCTATAATTG CGCATGTTGT TGGTTGTAGC ATGAGATATT AAAAGGAGTA CATGAAGC1713

ATTGCATGAG TTTAAGTGT AGCTCCTTAA TATTTTAGGT ATTTTCAAT TAATAAGT1773

TTGTGTTGTTG GGTAAAAAAA AAAAAAATAA AAAAAAATAA AAAAAAATAA A 1824

-continued

(2) INFORMATION FOR SEQ ID NO: 2 :

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2

TCGAATTCTN CCATTGG

18

(2) INFORMATION FOR SEQ ID NO: 3 :

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3

TCGAATTCTN CCATTGG

18

(2) INFORMATION FOR SEQ ID NO: 4 :

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4

TCGAATTCTN CCCTTGG

18

(2) INFORMATION FOR SEQ ID NO: 5 :

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5

TCGAATTCTN CCCTTGG

18

(2) INFORMATION FOR SEQ ID NO: 6 :

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Synthetic DNA

-continued

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6
TCGAATTCTN CCGTTCGG 18
- (2) INFORMATION FOR SEQ ID NO: 7 :
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7
TCGAATTCTN CCGTTTGG 18
- (2) INFORMATION FOR SEQ ID NO: 8 :
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8
TCGAATTCTN CCTTTCGG 18
- (2) INFORMATION FOR SEQ ID NO: 9 :
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9
TCGAATTCTN CCTTTTGG 18
- (2) INFORMATION FOR SEQ ID NO: 10 :
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10
TCGAATTCTN CCATTCTC 18
- (2) INFORMATION FOR SEQ ID NO: 11 :
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single

-continued

- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11
- TCGAATTC TN CCATTTTC 18
- (2) INFORMATION FOR SEQ ID NO: 12 :
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Genomic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12
- TCGAATTC TN CCCTTCTC 18
- (2) INFORMATION FOR SEQ ID NO: 13 :
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13
- TCGAATTC TN CCCTTTTC 18
- (2) INFORMATION FOR SEQ ID NO: 14 :
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14
- TCGAATTC TN CCGTTCTC 18
- (2) INFORMATION FOR SEQ ID NO: 15 :
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15
- TCGAATTC TN CCGTTTTC 18
- (2) INFORMATION FOR SEQ ID NO: 16 :
- (i) SEQUENCE CHARACTERISTICS:

-continued

- (A) LENGTH: 18 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other nucleic acid
- (A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16
- TCGAATTC TN CCTTTCTC 18
- (2) INFORMATION FOR SEQ ID NO: 17 :
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 18 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: Other nucleic acid
 - (A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17
- TCGAATTC TN CCTTTTTC 18
- (2) INFORMATION FOR SEQ ID NO: 18 :
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 18 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: Other nucleic acid
 - (A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18
- GCGGATCCCN CCNAAACA 18
- (2) INFORMATION FOR SEQ ID NO: 19 :
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 18 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: Other nucleic acid
 - (A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19
- GCGGATCCCN CCNAAAGCA 18
- (2) INFORMATION FOR SEQ ID NO: 20 :
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 18 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: Other nucleic acid
 - (A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20
- GCGGATCCCN CCNACACA 18

-continued

(2) INFORMATION FOR SEQ ID NO: 21 :

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid

- (A) DESCRIPTION: Synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21

GCGGATCCCN CCNACGCA

18

(2) INFORMATION FOR SEQ ID NO: 22 :

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid

- (A) DESCRIPTION: Synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22

GCGGATCCCN CCNAGACA

18

(2) INFORMATION FOR SEQ ID NO: 23 :

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid

- (A) DESCRIPTION: Synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23

GCGGATCCCN CCNAGGCA

18

(2) INFORMATION FOR SEQ ID NO: 24 :

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid

- (A) DESCRIPTION: Synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24

GCGGATCCCN CCNATACA

18

(2) INFORMATION FOR SEQ ID NO: 25 :

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid

- (A) DESCRIPTION: Synthetic DNA

-continued

-
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25
- GCGGATCCCN CCNATGCA 18
- (2) INFORMATION FOR SEQ ID NO: 26 :
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 18 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other nucleic acid
- (A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26
- GCGGATCCTN CCNGGACA 18
- (2) INFORMATION FOR SEQ ID NO: 27 :
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 18 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other nucleic acid
- (A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27
- GCGGATCCTN CCNGGGCA 18
- (2) INFORMATION FOR SEQ ID NO: 28 :
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 18 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other nucleic acid
- (A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 28
- GCGGATCCCN CCNGCACA 18
- (2) INFORMATION FOR SEQ ID NO: 29 :
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 18 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other nucleic acid
- (A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 29
- GCGGATCCCN CCNGCGCA 18
- (2) INFORMATION FOR SEQ ID NO: 30 :
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 32 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear

-continued

(ii) MOLECULE TYPE: cDNA to mRNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Petunia hybrida
- (B) STRAIN: Falcon Blue
- (F) TISSUE TYPE: flower limbs in the bud

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 30

CCN TTT GGT AGT GGA AGG AGG ATT TGC CCN GG 32
Pro Phe Gly Ser Gly Arg Arg Ile Cys Pro Gly
1 5 10

(2) INFORMATION FOR SEQ ID NO: 31 :

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Petunia hybrida
- (B) STRAIN: Falcon Blue
- (F) TISSUE TYPE: flower limbs in the bud

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 31

CCN TTT GGT GCT GGA AGA CGT ATA TGT CCN GG 32
Pro Phe Gly Ala Gly Arg Arg Ile Cys Pro Gly
1 5 10

(2) INFORMATION FOR SEQ ID NO: 32 :

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Petunia hybrida
- (B) STRAIN: Falcon Blue
- (F) TISSUE TYPE: flower limbs in the bud

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 32

CCN TTT GGT GCT GGT CGA AGA ATA TGC CCN GG 32
Pro Phe Gly Ala Gly Arg Arg Ile Cys Pro Gly
1 5 10

(2) INFORMATION FOR SEQ ID NO: 33 :

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Petunia hybrida
- (B) STRAIN: Falcon Blue
- (F) TISSUE TYPE: flower limbs in the bud

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 33

-continued

-
- CCN TTT GGG ACT GGT CGA CGA ATT TGT CCN GG 32
 Pro Phe Gly Thr Gly Arg Arg Ile Cys Pro Gly
 1 5 10
- (2) INFORMATION FOR SEQ ID NO: 34 :
- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 32 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA to mRNA
- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Petunia hybrida
 (B) STRAIN: Falcon Blue
 (F) TISSUE TYPE: flower limbs in the bud
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 34
- CCN TTT GGC TCG GGA AGA CGA TCT TGT CCN GG 32
 Pro Phe Gly Ser Gly Arg Arg Ser Cys Pro Gly
 1 5 10
- (2) INFORMATION FOR SEQ ID NO: 35 :
- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 32 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA to mRNA
- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Petunia hybrida
 (B) STRAIN: Falcon Blue
 (F) TISSUE TYPE: flower limbs in the bud
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 35
- CCN TTT GGT GCT GGT AGA AGA GTG TGT CCN GG 32
 Pro Phe Gly Ala Gly Arg Arg Val Cys Pro Gly
 1 5 10
- (2) INFORMATION FOR SEQ ID NO: 36 :
- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 32 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA to mRNA
- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Petunia hybrida
 (B) STRAIN: Falcon Blue
 (F) TISSUE TYPE: flower limbs in the bud
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 36
- CCN TTT GGA GTA GGC CTA AGA ATG TGC CCN GG 32
 Pro Phe Gly Val Gly Leu Arg Met Cys Pro Gly
 1 5 10
- (2) INFORMATION FOR SEQ ID NO: 37 :
- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 32 base pairs
 (B) TYPE: nucleic acid

-continued

- (C) STRANDEDNESS: double
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA to mRNA
- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Petunia hybrida
(B) STRAIN: Falcon Blue
(F) TISSUE TYPE: flower limbs in the bud
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 37
- CCN TTT GGT GGA GGA CCA CGG CGA TGT CCN GG 32
Pro Phe Gly Gly Gly Pro Arg Arg Cys Pro Gly
1 5 10
- (2) INFORMATION FOR SEQ ID NO: 38 :
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 32 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA to mRNA
- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Petunia hybrida
(B) STRAIN: Falcon Blue
(F) TISSUE TYPE: flower limbs in the bud
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 38
- CCN TTT GGT GTT GGT AGG AGG AGT TGC CCN GG 32
Pro Phe Gly Val Gly Arg Arg Ser Cys Pro Gly
1 5 10
- (2) INFORMATION FOR SEQ ID NO: 39 :
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 32 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA to mRNA
- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Petunia hybrida
(B) STRAIN: Falcon Blue
(F) TISSUE TYPE: flower limbs in the bud
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 39
- CCN TTC GGA GTC GGC CCC AAA ATG TGC CCN GG 32
Pro Phe Gly Val Gly Pro Lys Met Cys Pro Gly
1 5 10
- (2) INFORMATION FOR SEQ ID NO: 40 :
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 32 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA to mRNA
- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Petunia hybrida
(B) STRAIN: Falcon Blue
(F) TISSUE TYPE: flower limbs in the bud

-continued

-
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 40
- CCN TTC GGT GGA GGA CCA AGA AAA TGC GTN GG 32
 Pro Phe Gly Gly Gly Pro Arg Lys Cys Val Gly
 1 5 10
- (2) INFORMATION FOR SEQ ID NO: 41 :
- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 32 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA to mRNA
- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Petunia hybrida
 (B) STRAIN: Falcon Blue
 (F) TISSUE TYPE: flower limbs in the bud
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 41
- CCN TTC GGC TTT GGT CCT CGA AAA TGC GTN GG 32
 Pro Phe Gly Phe Gly Pro Arg Lys Cys Val Gly
 1 5 10
- (2) INFORMATION FOR SEQ ID NO: 42 :
- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 32 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA to mRNA
- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Petunia hybrida
 (B) STRAIN: Falcon Blue
 (F) TISSUE TYPE: flower limbs in the bud
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 42
- CCN TTT GGC AGT GGT TTC TGT TCA TGT CCN GG 32
 Pro Phe Gly Ser Gly Phe Cys Ser Cys Pro Gly
 1 5 10
- (2) INFORMATION FOR SEQ ID NO: 43 :
- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 32 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA to mRNA
- (iii) HYPOTHETICAL: <Unknown>
- (iv) ANTI-SENSE: <Unknown>
- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Petunia hybrida
 (B) STRAIN: Falcon Blue
 (F) TISSUE TYPE: flower limbs in the bud
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 43
- CCN TTT GGT GCT GGA CGA AGA ATT TGT GCN GG 32
 Pro Phe Gly Ala Gly Arg Arg Ile Cys Ala Gly
 1 5 10

-continued

(2) INFORMATION FOR SEQ ID NO: 44 :

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Petunia hybrida
- (B) STRAIN: Falcon Blue
- (F) TISSUE TYPE: flower limbs in the bud

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 44

```
CCN TTT GGT GGT GGA AGA AGG ATA TGT CCN GG      32
Pro Phe Gly Gly Gly Arg Arg Ile Cys Pro Gly
1           5           10
```

(2) INFORMATION FOR SEQ ID NO: 45 :

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid

- (A) DESCRIPTION: Synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 45

```
CCNGGGCAAA TCCTCCT      17
```

(2) INFORMATION FOR SEQ ID NO: 46 :

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid

- (A) DESCRIPTION: Synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 46

```
CCNGGACATA TACGTCT      17
```

(2) INFORMATION FOR SEQ ID NO: 47 :

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid

- (A) DESCRIPTION: Synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 47

```
CCNGGCATA TTCTTCG      17
```

(2) INFORMATION FOR SEQ ID NO: 48 :

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 base pairs
- (B) TYPE: nucleic acid

-continued

- (C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 48
CCNGGACAAA TTCGTCG 17
- (2) INFORMATION FOR SEQ ID NO: 49 :
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 49
CCNGGACAAG ATCGTCT 17
- (2) INFORMATION FOR SEQ ID NO: 50 :
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 50
CCNGGACACA CTCTTCT 17
- (2) INFORMATION FOR SEQ ID NO: 51 :
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 51
CCNGGCACA TTCTTAG 17
- (2) INFORMATION FOR SEQ ID NO: 52 :
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 52
CCNGGACATC GCCGTGG 17
- (2) INFORMATION FOR SEQ ID NO: 53 :

-continued

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 53
- CCNGGGCAAC TCCTCCT 17
- (2) INFORMATION FOR SEQ ID NO: 54 :
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 54
- CCNGGGCACA TTTTGGG 17
- (2) INFORMATION FOR SEQ ID NO: 55 :
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 55
- CCNACGCATT TTCTTGG 17
- (2) INFORMATION FOR SEQ ID NO: 56 :
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 56
- CCNACACATT TTCGAGG 17
- (2) INFORMATION FOR SEQ ID NO: 57 :
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 57

-continued

- CCNGGACATG AACAGAA 17
- (2) INFORMATION FOR SEQ ID NO: 58 :
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 17 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: Other nucleic acid
 - (A) DESCRIPTION: Synthetic DNA
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 58
- CCNGCACAAA TTCTTCG 17
- (2) INFORMATION FOR SEQ ID NO: 59 :
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 17 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: Other nucleic acid
 - (A) DESCRIPTION: Synthetic DNA
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 59
- CCNGGACATA TCCTTCT 17
- (2) INFORMATION FOR SEQ ID NO: 60 :
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: Other nucleic acid
 - (A) DESCRIPTION: Synthetic DNA
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 60
- TGATCCGGAA TTCGTCCAT CAAG 24
- (2) INFORMATION FOR SEQ ID NO: 61 :
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 26 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: Other nucleic acid
 - (A) DESCRIPTION: Synthetic DNA
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 61
- CGCTTGATGG CACGAATCC GGATCA 26
- (2) INFORMATION FOR SEQ ID NO: 62 :
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 15 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: Other nucleic acid

-continued

(A) DESCRIPTION: Synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 62

CCNTTTGGTG CTGGA

15

(2) INFORMATION FOR SEQ ID NO: 63 :

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2174 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Eustoma russellianum

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 92 to 1621
- (C) IDENTIFICATION METHOD: by experiment

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 63

GAAACTATC CATTCTTACC AAGATAAGCA CATTTCCTCGT TTCTTTCTAA GAAGAGCATT60

AGGCCAATTC TTAAAGCCCG TACTTAACGA T ATG GCT GTT GGA AAT GGC GTT 112

Met Ala Val Gly Asn Gly Val
1 5

TTA CTT CAC ATT GCT GCA TCA TTG ATG CTG TTC TTT CAT GTG CAA AAA 160
Leu Leu His Ile Ala Ala Ser Leu Met Leu Phe Phe His Val Gln Lys
10 15 20

CTT GTG CAA TAT CTA TGG ATG AAT TCC AGG CGC CAC CGG CTT CCA CCT 208
Leu Val Gln Tyr Leu Trp Met Asn Ser Arg Arg His Arg Leu Pro Pro
25 30 35

GGC CCG ATA GGG TGG CCG GTT CTC GGT GCC CTT CGG CTT TTA GGC ACC 256
Gly Pro Ile Gly Trp Pro Val Leu Gly Ala Leu Arg Leu Leu Gly Thr
40 45 50 55

ATG CCT CAT GTT GCA CTA GCT AAC ATG GCC AAA AAA TAT GGT CCT GTT 304
Met Pro His Val Ala Leu Ala Asn Met Ala Lys Lys Tyr Gly Pro Val
60 65 70

ATG TAC TTA AAG GTA GGC AGC TGT GGT CTG GCC GTG GCA TCG ACT CCT 352
Met Tyr Leu Lys Val Gly Ser Cys Gly Leu Ala Val Ala Ser Thr Pro
75 80 85

GAG GCT GCT AAG GCA TTC CTC AAA ACA CTT GAC ATG AAC TTC TCG AAT 400
Glu Ala Ala Lys Ala Phe Leu Lys Thr Leu Asp Met Asn Phe Ser Asn
90 95 100

CGG CCG CCT AAT GCA GGG GCT ACC CAT TTG GCC TAT AAT GCT CAG GAC 448
Arg Pro Pro Asn Ala Gly Ala Thr His Leu Ala Tyr Asn Ala Gln Asp
105 110 115

ATG GTG TTT GCA GAC TAT GGT CCC AGA TGG AAG CTG CTA CGT AAA CTC 496
Met Val Phe Ala Asp Tyr Gly Pro Arg Trp Lys Leu Leu Arg Lys Leu
120 125 130 135

AGC AAC ATA CAC ATT CTT GGT GGC AAG GCC CTG CAG GGC TGG GAA GAA 544
Ser Asn Ile His Ile Leu Gly Gly Lys Ala Leu Gln Gly Trp Glu Glu
140 145 150

GTT CGA AAG AAA GAG CTT GGG TAT ATG CTC TAT GCA ATG GCT GAA TCA 592
Val Arg Lys Lys Glu Leu Gly Tyr Met Leu Tyr Ala Met Ala Glu Ser
155 160 165

GGG CGA CAT GGC CAG CCA GTG GTG GTG TCA GAG ATG CTA ACA TAT GCC 640
Gly Arg His Gly Gln Pro Val Val Val Ser Glu Met Leu Thr Tyr Ala

-continued

| 170 | 175 | 180 |
|---|-----|-----|
| ATG GCA AAC ATG TTA GGC CAA GTG ATG CTC AGC AAG CGA GTT TTC GGG 688 Met Ala Asn Met Leu Gly Gln Val Met Leu Ser Lys Arg Val Phe Gly 185 190 195 | | |
| TCT CAA GGA TCA GAA TCC AAT GAG TTC AAA GAT ATG GTG GTT GAG TTG 736 Ser Gln Gly Ser Glu Ser Asn Glu Phe Lys Asp Met Val Val Glu Leu 200 205 210 215 | | |
| ATG ACT GTT GCT GGC TAT TTC AAC ATA GGT GAT TTT ATC CCC TCG ATT 784 Met Thr Val Ala Gly Tyr Phe Asn Ile Gly Asp Phe Ile Pro Ser Ile 220 225 230 | | |
| GCA TGG ATG GAT TTG CAG GGG ATT CAG GGC GGA ATG AAA CGG TTG CAT 832 Ala Trp Met Asp Leu Gln Gly Ile Gln Gly Gly Met Lys Arg Leu His 235 240 245 | | |
| AAG AAG TTT GAT GCT TTG TTG ACT CGG TTG CTG GAA GAG CAC ACT GCA 880 Lys Lys Phe Asp Ala Leu Leu Thr Arg Leu Leu Glu Glu His Thr Ala 250 255 260 | | |
| TCG GCT CAT GAG CGT AAA GGC AGC CCT GAT TTC CTT GAT TTT GTC GTT 928 Ser Ala His Glu Arg Lys Gly Ser Pro Asp Phe Leu Asp Phe Val Val 265 270 275 | | |
| GCA AAT GGC GAC AAT TCT GAA GGC GAA AGG CTT CAG ACA GTC AAT ATC 976 Ala Asn Gly Asp Asn Ser Glu Gly Glu Arg Leu Gln Thr Val Asn Ile 280 285 290 295 | | |
| AAG GCT CTT TTA TTG AAC ATG TTT ACC GCT GGT ACG GAT ACA TCT TCA1024 Lys Ala Leu Leu Leu Asn Met Phe Thr Ala Gly Thr Asp Thr Ser Ser 300 305 310 | | |
| AGC GTC ATA GAG TGG GCG CTG GCC GAG TTG CTA AAG AAT CCA ATC ATC1072 Ser Val Ile Glu Trp Ala Leu Ala Glu Leu Leu Lys Asn Pro Ile Ile 315 320 325 | | |
| CTA AGA CGA GCC CAA GAA GAA ATG GAC GGT GTG ATC GGC CGA GAC CGG1120 Leu Arg Arg Ala Gln Glu Glu Met Asp Gly Val Ile Gly Arg Asp Arg 330 335 340 | | |
| CGG TTT CTT GAG GCA GAC ATA TCA AAG TTG CCA TAT CTC CAA GCC ATC1168 Arg Phe Leu Glu Ala Asp Ile Ser Lys Leu Pro Tyr Leu Gln Ala Ile 345 350 355 | | |
| TGC AAA GAA GCT TTC AGA AAG CAT CCT TCC ACG CCT TTA AAT CTC CCA1216 Cys Lys Glu Ala Phe Arg Lys His Pro Ser Thr Pro Leu Asn Leu Pro 360 365 370 375 | | |
| CGA ATC GCG TCG CAA GCA TGT GAA GTA AAT GGA CAC TAC ATA CCA AAG1264 Arg Ile Ala Ser Gln Ala Cys Glu Val Asn Gly His Tyr Ile Pro Lys 380 385 390 | | |
| GGC ACT AGG CTC AGC GTT AAC ATA TGG GCT ATT GGA AGA GAT CCA TCT1312 Gly Thr Arg Leu Ser Val Asn Ile Trp Ala Ile Gly Arg Asp Pro Ser 395 400 405 | | |
| GTG TGG GAA AAT CCA AAT GAA TTT AAC CCT GAT AGG TTT TTG GAA CGA1360 Val Trp Glu Asn Pro Asn Glu Phe Asn Pro Asp Arg Phe Leu Glu Arg 410 415 420 | | |
| AAG AAT GCC AAG ATC GAT CCA CGA GGA AAT GAT TTT GAG CTG ATC CCA1408 Lys Asn Ala Lys Ile Asp Pro Arg Gly Asn Asp Phe Glu Leu Ile Pro 425 430 435 | | |
| TTT GGA GCT GGA AGA AGA ATT TGC GCT GGA ACA AGA TTG GGA ATA CTT1456 Phe Gly Ala Gly Arg Arg Ile Cys Ala Gly Thr Arg Leu Gly Ile Leu 440 445 450 455 | | |
| CTA GTG GAG TAT ATT TTG GGA ACT TTG GTG CAT TCT TTT GTT TGG GAA1504 Leu Val Glu Tyr Ile Leu Gly Thr Leu Val His Ser Phe Val Trp Glu 460 465 470 | | |
| TTG CCA TCC TCT GTG ATT GAA CTT AAC ATG GAT GAG TCT TTT GGG CTT1552 Leu Pro Ser Ser Val Ile Glu Leu Asn Met Asp Glu Ser Phe Gly Leu | | |

-continued

| | | | |
|--|-----|-----|--|
| 475 | 480 | 485 | |
| GCT CTG CAG AAG GCA GTG CCT CTT GCT GCT ATG GTC ACT CCA CGG CTG1600 | | | |
| Ala Leu Gln Lys Ala Val Pro Leu Ala Ala Met Val Thr Pro Arg Leu | | | |
| 490 | 495 | 500 | |
| CCT CTC CAT ATT TAC TCT CCT TGAGATCTGT GTTCTATGGG TCATTGAGAA 1651 | | | |
| Pro Leu His Ile Tyr Ser Pro | | | |
| 505 | 510 | | |
| ACAACCCGCTG TGTGTTTCTA ACACATGAAT ATGGTTGTGT ACATCTGGCT TATTTATA1711 | | | |
| TCCCTATAGA CGAGAAGCCT CGAAGGCAAT GGGTAATGT TGTGTTGTGC GTGAGACA1771 | | | |
| TCTTCTATGT TTCTAAGCAG ATGAGATCTA AGTAGATGAC ATATGCTGTC TTCTACTA1831 | | | |
| TTGAAATTAG ATATGCCCCA GAATAAACGC ATCAAACCTCG TAATTCGATA CAAAAAAT1891 | | | |
| TTGTTGTGGT TTTGAATAAA CACTTATAGA TAATTTGAGA TTTAGAATCG GGTATTTT1951 | | | |
| TATATTTTCC ACGTTCATAG GAGTTCGTCC ATGTTTCTGA TTTACAAATA TGATTTT2011 | | | |
| TGGACATTC TAATAATATC AATTTGTATT CCTGTTTAA GTTTTTTAAT TTCTCAAG2071 | | | |
| TTAGTCCTAA TTAGCAAAGG ACCAGAAAAA CTGCTAGTT ATGAATCGGG GATAGAAC2131 | | | |
| GCAGGAGATG CTGTTACAA TTTGATTAA AAAAAAAAAA AAA 2174 | | | |

(2) INFORMATION FOR SEQ ID NO: 64 :

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1927 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Campanula medium

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 180 to 1748
- (C) IDENTIFICATION METHOD: by experiment

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 64

ACCAAATGAG CTTTGTAATT TGAGATTAAT CATAATTGCA TGCTCAACTA ACATTCTGTA60

TTCATATATC CATATGTATT TTGACCTATA GATATTACAT TACACCTTGA GGCCTTTAT120

TATAGAGAGT GTATCTACTT CCCTTAATAT CACCTTTTCA TTCACAAGT GAAGCCACC179

ATG TCT ATA GAC ATA TCC ACC CTC TTC TAT GAA CTT GTT GCA GCA ATT 227
 Met Ser Ile Asp Ile Ser Thr Leu Phe Tyr Glu Leu Val Ala Ala Ile
 1 5 10 15

TCA CTC TAC TTA GCT ACC TAC TCT TTC ATT CGT TTC CTC TTC AAA CCC 275
 Ser Leu Tyr Leu Ala Thr Tyr Ser Phe Ile Arg Phe Leu Phe Lys Pro
 20 25 30

TCT CAC CAC CAC CAC CTC CCT CCC GGC CCA ACC GGA TGG CCG ATC ATC 323
 Ser His His His His Leu Pro Pro Gly Pro Thr Gly Trp Pro Ile Ile
 35 40 45

GGA GCC CTT CCA CTC TTA GGC ACC ATG CCA CAT GTT TCC TTA GCC GAC 371
 Gly Ala Leu Pro Leu Leu Gly Thr Met Pro His Val Ser Leu Ala Asp
 50 55 60

ATG GCC GTT AAA TAC GGT CCT ATA ATG TAC CTA AAA CTT GGT TCA AAG 419
 Met Ala Val Lys Tyr Gly Pro Ile Met Tyr Leu Lys Leu Gly Ser Lys
 65 70 75 80

GGC ACC GTC GTG GCC TCA AAT CCA AAA GCC GCC CGA GCC TTC TTG AAA 467

-continued

| | | | | | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|--|--|--|
| Gly | Thr | Val | Val | Ala | Ser | Asn | Pro | Lys | Ala | Ala | Arg | Ala | Phe | Leu | Lys | | | | |
| | | | | 85 | | | | 90 | | | | | | 95 | | | | | |
| ACC | CAT | GAT | GCC | AAT | TTT | TCT | AAC | CGT | CCG | ATT | GAT | GGG | GGC | CCT | ACC | 515 | | | |
| Thr | His | Asp | Ala | Asn | Phe | Ser | Asn | Arg | Pro | Ile | Asp | Gly | Gly | Pro | Thr | | | | |
| | | | 100 | | | | | 105 | | | | | | 110 | | | | | |
| TAC | CTC | GCG | TAT | AAT | GCA | CAA | GAC | ATG | GTT | TTT | GCA | GAA | TAT | GGC | CCA | 563 | | | |
| Tyr | Leu | Ala | Tyr | Asn | Ala | Gln | Asp | Met | Val | Phe | Ala | Glu | Tyr | Gly | Pro | | | | |
| | | 115 | | | | | 120 | | | | | 125 | | | | | | | |
| AAA | TGG | AAG | CTT | TTG | CGA | AAG | CTA | TGT | AGC | TTG | CAC | ATG | TTA | GGC | CCG | 611 | | | |
| Lys | Trp | Lys | Leu | Leu | Arg | Lys | Leu | Cys | Ser | Leu | His | Met | Leu | Gly | Pro | | | | |
| | | 130 | | | | 135 | | | | | | 140 | | | | | | | |
| AAG | GCA | CTC | GAG | GAT | TGG | GCT | CAT | GTC | AAA | GTT | TCA | GAG | GTC | GGT | CAT | 659 | | | |
| Lys | Ala | Leu | Glu | Asp | Trp | Ala | His | Val | Lys | Val | Ser | Glu | Val | Gly | His | | | | |
| | | 145 | | | 150 | | | | | 155 | | | | | 160 | | | | |
| ATG | CTC | AAA | GAA | ATG | TAC | GAG | CAA | TCG | AGT | AAG | TCA | GTG | CCA | GTG | CCA | 707 | | | |
| Met | Leu | Lys | Glu | Met | Tyr | Glu | Gln | Ser | Ser | Lys | Ser | Val | Pro | Val | Pro | | | | |
| | | | | 165 | | | | | | 170 | | | | | 175 | | | | |
| GTG | GTG | GTG | CCA | GAG | ATG | TTA | ACT | TAT | GCC | ATG | GCT | AAT | ATG | ATT | GGA | 755 | | | |
| Val | Val | Val | Pro | Glu | Met | Leu | Thr | Tyr | Ala | Met | Ala | Asn | Met | Ile | Gly | | | | |
| | | | 180 | | | | | 185 | | | | | 190 | | | | | | |
| CGA | ATC | ATA | CTC | AGC | CGA | CGC | CCT | TTT | GTT | ATC | ACG | AGC | AAA | TTA | GAC | 803 | | | |
| Arg | Ile | Ile | Leu | Ser | Arg | Arg | Pro | Phe | Val | Ile | Thr | Ser | Lys | Leu | Asp | | | | |
| | | 195 | | | | | 200 | | | | | | 205 | | | | | | |
| TCG | TCT | GCT | TCT | GCT | TCT | GCT | TCT | GTT | AGT | GAA | TTC | CAA | TAT | ATG | GTT | 851 | | | |
| Ser | Ser | Ala | Ser | Ala | Ser | Ala | Ser | Val | Ser | Glu | Phe | Gln | Tyr | Met | Val | | | | |
| | | 210 | | | | 215 | | | | | 220 | | | | | | | | |
| ATG | GAG | CTC | ATG | AGG | ATG | GCA | GGG | TTG | TTC | AAT | ATT | GGT | GAT | TTC | ATA | 899 | | | |
| Met | Glu | Leu | Met | Arg | Met | Ala | Gly | Leu | Phe | Asn | Ile | Gly | Asp | Phe | Ile | | | | |
| | | 225 | | | 230 | | | | | 235 | | | | | 240 | | | | |
| CCA | TAT | ATT | GCA | TGG | ATG | GAT | TTG | CAA | GGC | ATT | CAA | CGT | GAT | ATG | AAG | 947 | | | |
| Pro | Tyr | Ile | Ala | Trp | Met | Asp | Leu | Gln | Gly | Ile | Gln | Arg | Asp | Met | Lys | | | | |
| | | | | 245 | | | | | 250 | | | | | | 255 | | | | |
| GTT | ATA | CAG | AAA | AAG | TTT | GAT | GTC | TTG | TTG | AAC | AAA | ATG | ATC | AAG | GAA | 995 | | | |
| Val | Ile | Gln | Lys | Lys | Phe | Asp | Val | Leu | Leu | Asn | Lys | Met | Ile | Lys | Glu | | | | |
| | | | 260 | | | | | 265 | | | | | 270 | | | | | | |
| CAT | ACA | GAA | TCC | GCT | CAT | GAT | CGC | AAA | GAT | AAT | CCT | GAT | TTT | CTT | GAT | 1043 | | | |
| His | Thr | Glu | Ser | Ala | His | Asp | Arg | Lys | Asp | Asn | Pro | Asp | Phe | Leu | Asp | | | | |
| | | 275 | | | | | 280 | | | | | | 285 | | | | | | |
| ATT | CTT | ATG | GCG | GCT | ACC | CAA | GAA | AAC | ACG | GAG | GGA | ATT | CAG | CTT | AAT | 1091 | | | |
| Ile | Leu | Met | Ala | Ala | Thr | Gln | Glu | Asn | Thr | Glu | Gly | Ile | Gln | Leu | Asn | | | | |
| | | 290 | | | | 295 | | | | | | 300 | | | | | | | |
| CTT | GTA | AAT | GTT | AAG | GCA | CTT | CTT | TTG | GAT | TTA | TTC | ACG | GCG | GGC | ACG | 1139 | | | |
| Leu | Val | Asn | Val | Lys | Ala | Leu | Leu | Leu | Asp | Leu | Phe | Thr | Ala | Gly | Thr | | | | |
| | | 305 | | | 310 | | | | | | 315 | | | | 320 | | | | |
| GAT | ACA | TCA | TCA | AGT | GTG | ATC | GAA | TGG | GCA | CTA | GCC | GAA | ATG | TTG | AAC | 1187 | | | |
| Asp | Thr | Ser | Ser | Ser | Val | Ile | Glu | Trp | Ala | Leu | Ala | Glu | Met | Leu | Asn | | | | |
| | | | | 325 | | | | | 330 | | | | | | 335 | | | | |
| CAT | CGA | CAG | ATC | CTA | AAC | CGG | GCC | CAC | GAA | GAA | ATG | GAC | CAA | GTC | ATT | 1235 | | | |
| His | Arg | Gln | Ile | Leu | Asn | Arg | Ala | His | Glu | Glu | Met | Asp | Gln | Val | Ile | | | | |
| | | | 340 | | | | | | 345 | | | | | 350 | | | | | |
| GGC | AGA | AAC | AGA | AGA | CTA | GAA | CAA | TCT | GAC | ATA | CCA | AAC | TTG | CCA | TAT | 1283 | | | |
| Gly | Arg | Asn | Arg | Arg | Leu | Glu | Gln | Ser | Asp | Ile | Pro | Asn | Leu | Pro | Tyr | | | | |
| | | 355 | | | | | 360 | | | | | | 365 | | | | | | |
| TTC | CAA | GCC | ATA | TGC | AAA | GAA | ACA | TTC | CGA | AAA | CAC | CCT | TCC | ACG | CCC | 1331 | | | |
| Phe | Gln | Ala | Ile | Cys | Lys | Glu | Thr | Phe | Arg | Lys | His | Pro | Ser | Thr | Pro | | | | |
| | | 370 | | | | 375 | | | | | 380 | | | | | | | | |
| TTA | AAC | CTC | CCA | AGA | ATC | TCA | ACA | GAA | GCA | TGT | GAA | GTG | GAC | GGA | TTT | 1379 | | | |

-continued

```

Leu Asn Leu Pro Arg Ile Ser Thr Glu Ala Cys Glu Val Asp Gly Phe
385                390                395                400

CAC ATA CCA AAA AAC ACT AGA CTA ATA GTG AAC ATA TGG GCA ATA GGG1427
His Ile Pro Lys Asn Thr Arg Leu Ile Val Asn Ile Trp Ala Ile Gly
                405                410                415

AGG GAC CCT AAA GTG TGG GAA AAT CCA TTA GAT TTT ACT CCG GAA CGT1475
Arg Asp Pro Lys Val Trp Glu Asn Pro Leu Asp Phe Thr Pro Glu Arg
                420                425                430

TTC TTG AGT GAA AAA CAC GCG AAA ATT GAT CCG CGA GGT AAT CAT TTT1523
Phe Leu Ser Glu Lys His Ala Lys Ile Asp Pro Arg Gly Asn His Phe
                435                440                445

GAG TTA ATC CCA TTT GGG GCT GGA CGA AGG ATA TGT GCA GGG GCT AGA1571
Glu Leu Ile Pro Phe Gly Ala Gly Arg Arg Ile Cys Ala Gly Ala Arg
                450                455                460

ATG GGA GCG GCC TCG GTC GAG TAC ATA TTA GGT ACA TTG GTG CAC TCA1619
Met Gly Ala Ala Ser Val Glu Tyr Ile Leu Gly Thr Leu Val His Ser
                465                470                475                480

TTT GAT TGG AAA TTG CCT GAT GGA GTT GTG GAA GTT AAT ATG GAA GAG1667
Phe Asp Trp Lys Leu Pro Asp Gly Val Val Glu Val Asn Met Glu Glu
                485                490                495

AGC TTT GGG ATC GCA TTG CAA AAA AAA GTG CCT CTT TCT GCT ATT GTT1715
Ser Phe Gly Ile Ala Leu Gln Lys Lys Val Pro Leu Ser Ala Ile Val
                500                505                510

ACT CCA AGA TTG CCT CCA AGT TCT TAC ACT GTC TAGGCAAATG CTTATATA1768
Thr Pro Arg Leu Pro Pro Ser Ser Tyr Thr Val
                515                520

TGAATAATTG ATTGAGTTGT TTAGTTGTAT GAAAGATTTG AGAAAATAAA TTATTAGG1828

TTGCACCATT ATGTTGAGAT GGTGTGTGTT AGTGTTAAGG AAGTCGATTG TAGTAATA1888

AATTTTATTT TTTTCGAAAA AAAAAAAAAA AAAAAAAAAA                1927
    
```

(2) INFORMATION FOR SEQ ID NO: 65 :

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 506 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: *Petunia hybrida*
 - (B) STRAIN: Falcon Blue
- (ix) FEATURE:
 - (A) NAME/KEY: CDS
 - (B) LOCATION: 116 to 1633
 - (C) IDENTIFICATION METHOD: by experiment
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 65

Met

1

```

Met Leu Leu Thr Glu Leu Gly Ala Ala Thr Ser Ile Phe Leu Ile Ala
    5                10                15

His Ile Ile Ile Ser Thr Leu Ile Ser Lys Thr Thr Gly Arg His Leu
    20                25                30

Pro Pro Gly Pro Arg Gly Trp Pro Val Ile Gly Ala Leu Pro Leu Leu
    35                40                45
    
```

-continued

Gly Ala Met Pro His Val Ser Leu Ala Lys Met Ala Lys Lys Tyr Gly
 50 55 60 65
 Ala Ile Met Tyr Leu Lys Val Gly Thr Cys Gly Met Ala Val Ala Ser
 70 75 80
 Thr Pro Asp Ala Ala Lys Ala Phe Leu Lys Thr Leu Asp Ile Asn Phe
 85 90 95
 Ser Asn Arg Pro Pro Asn Ala Gly Ala Thr His Leu Ala Tyr Asn Ala
 100 105 110
 Gln Asp Met Val Phe Ala His Tyr Gly Pro Arg Trp Lys Leu Leu Arg
 115 120 125
 Lys Leu Ser Asn Leu His Met Leu Gly Gly Lys Ala Leu Glu Asn Trp
 130 135 140 145
 Ala Asn Val Arg Ala Asn Glu Leu Gly His Met Leu Lys Ser Met Ser
 150 155 160
 Asp Met Ser Arg Glu Gly Gln Arg Val Val Val Ala Glu Met Leu Thr
 165 170 175
 Phe Ala Met Ala Asn Met Ile Gly Gln Val Met Leu Ser Lys Arg Val
 180 185 190
 Phe Val Asp Lys Gly Val Glu Val Asn Glu Phe Lys Asp Met Val Val
 195 200 205
 Glu Leu Met Thr Ile Ala Gly Tyr Phe Asn Ile Gly Asp Phe Ile Pro
 210 215 220 225
 Cys Leu Ala Trp Met Asp Leu Gln Gly Ile Glu Lys Arg Met Lys Arg
 230 235 240
 Leu His Lys Lys Phe Asp Ala Leu Leu Thr Lys Met Phe Asp Glu His
 245 250 255
 Lys Ala Thr Thr Tyr Glu Arg Lys Gly Lys Pro Asp Phe Leu Asp Val
 260 265 270
 Val Met Glu Asn Gly Asp Asn Ser Glu Gly Glu Arg Leu Ser Thr Thr
 275 280 285
 Asn Ile Lys Ala Leu Leu Leu Asn Leu Phe Thr Ala Gly Thr Asp Thr
 290 295 300 305
 Ser Ser Ser Ala Ile Glu Trp Ala Leu Ala Glu Met Met Lys Asn Pro
 310 315 320
 Ala Ile Leu Lys Lys Ala Gln Ala Glu Met Asp Gln Val Ile Gly Arg
 325 330 335
 Asn Arg Arg Leu Leu Glu Ser Asp Ile Pro Asn Leu Pro Tyr Leu Arg
 340 345 350
 Ala Ile Cys Lys Glu Thr Phe Arg Lys His Pro Ser Thr Pro Leu Asn
 355 360 365
 Leu Pro Arg Ile Ser Asn Glu Pro Cys Ile Val Asp Gly Tyr Tyr Ile
 370 375 380 385
 Pro Lys Asn Thr Arg Leu Ser Val Asn Ile Trp Ala Ile Gly Arg Asp
 390 395 400
 Pro Gln Val Trp Glu Asn Pro Leu Glu Phe Asn Pro Glu Arg Phe Leu
 405 410 415
 Ser Gly Arg Asn Ser Lys Ile Asp Pro Arg Gly Asn Asp Phe Glu Leu
 420 425 430
 Ile Pro Phe Gly Ala Gly Arg Arg Ile Cys Ala Gly Thr Arg Met Gly
 435 440 445
 Ile Val Met Val Glu Tyr Ile Leu Gly Thr Leu Val His Ser Phe Asp

-continued

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 450 | | 455 | | 460 | | 465 | | | | | | | | | |
| Trp | Lys | Leu | Pro | Ser | Glu | Val | Ile | Glu | Leu | Asn | Met | Glu | Glu | Ala | Phe |
| | | | | 470 | | | | | 475 | | | | | 480 | |
| Gly | Leu | Ala | Leu | Gln | Lys | Ala | Val | Pro | Leu | Glu | Ala | Met | Val | Thr | Pro |
| | | | 485 | | | | | 490 | | | | | 495 | | |
| Arg | Leu | Gln | Leu | Asp | Val | Tyr | Val | Pro | | | | | | | |
| | | 500 | | | | | 505 | | | | | | | | |

(2) INFORMATION FOR SEQ ID NO: 66 :

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 510 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Eustoma russellianum
- (ix) FEATURE:
 - (A) NAME/KEY: CDS
 - (B) LOCATION: 92 to 1621
 - (C) IDENTIFICATION METHOD:by experiment
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 66

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | | | | | | | | Met | Ala | Val | Gly | Asn | Gly | Val | |
| | | | | | | | | 1 | | | | 5 | | | |
| Leu | Leu | His | Ile | Ala | Ala | Ser | Leu | Met | Leu | Phe | Phe | His | Val | Gln | Lys |
| | | 10 | | | | | 15 | | | | | 20 | | | |
| Leu | Val | Gln | Tyr | Leu | Trp | Met | Asn | Ser | Arg | Arg | His | Arg | Leu | Pro | Pro |
| | | 25 | | | | 30 | | | | | 35 | | | | |
| Gly | Pro | Ile | Gly | Trp | Pro | Val | Leu | Gly | Ala | Leu | Arg | Leu | Leu | Gly | Thr |
| | | | | | 45 | | | | | 50 | | | | 55 | |
| Met | Pro | His | Val | Ala | Leu | Ala | Asn | Met | Ala | Lys | Lys | Tyr | Gly | Pro | Val |
| | | | | 60 | | | | | 65 | | | | | 70 | |
| Met | Tyr | Leu | Lys | Val | Gly | Ser | Cys | Gly | Leu | Ala | Val | Ala | Ser | Thr | Pro |
| | | | 75 | | | | | 80 | | | | | 85 | | |
| Glu | Ala | Ala | Lys | Ala | Phe | Leu | Lys | Thr | Leu | Asp | Met | Asn | Phe | Ser | Asn |
| | | 90 | | | | | 95 | | | | | 100 | | | |
| Arg | Pro | Pro | Asn | Ala | Gly | Ala | Thr | His | Leu | Ala | Tyr | Asn | Ala | Gln | Asp |
| | | 105 | | | | 110 | | | | | | 115 | | | |
| Met | Val | Phe | Ala | Asp | Tyr | Gly | Pro | Arg | Trp | Lys | Leu | Leu | Arg | Lys | Leu |
| | | | | | 125 | | | | | 130 | | | | 135 | |
| Ser | Asn | Ile | His | Ile | Leu | Gly | Gly | Lys | Ala | Leu | Gln | Gly | Trp | Glu | Glu |
| | | | | 140 | | | | | 145 | | | | | 150 | |
| Val | Arg | Lys | Lys | Glu | Leu | Gly | Tyr | Met | Leu | Tyr | Ala | Met | Ala | Glu | Ser |
| | | | 155 | | | | | 160 | | | | | 165 | | |
| Gly | Arg | His | Gly | Gln | Pro | Val | Val | Val | Ser | Glu | Met | Leu | Thr | Tyr | Ala |
| | | | 170 | | | | 175 | | | | | | 180 | | |
| Met | Ala | Asn | Met | Leu | Gly | Gln | Val | Met | Leu | Ser | Lys | Arg | Val | Phe | Gly |
| | | 185 | | | | 190 | | | | | 195 | | | | |
| Ser | Gln | Gly | Ser | Glu | Ser | Asn | Glu | Phe | Lys | Asp | Met | Val | Val | Glu | Leu |
| | | 200 | | | 205 | | | | | 210 | | | | 215 | |
| Met | Thr | Val | Ala | Gly | Tyr | Phe | Asn | Ile | Gly | Asp | Phe | Ile | Pro | Ser | Ile |
| | | | | 220 | | | | | 225 | | | | | 230 | |

-continued

Ala Trp Met Asp Leu Gln Gly Ile Gln Gly Gly Met Lys Arg Leu His
235 240 245

Lys Lys Phe Asp Ala Leu Leu Thr Arg Leu Leu Glu Glu His Thr Ala
250 255 260

Ser Ala His Glu Arg Lys Gly Ser Pro Asp Phe Leu Asp Phe Val Val
265 270 275

Ala Asn Gly Asp Asn Ser Glu Gly Glu Arg Leu Gln Thr Val Asn Ile
280 285 290 295

Lys Ala Leu Leu Leu Asn Met Phe Thr Ala Gly Thr Asp Thr Ser Ser
300 305 310

Ser Val Ile Glu Trp Ala Leu Ala Glu Leu Leu Lys Asn Pro Ile Ile
315 320 325

Leu Arg Arg Ala Gln Glu Glu Met Asp Gly Val Ile Gly Arg Asp Arg
330 335 340

Arg Phe Leu Glu Ala Asp Ile Ser Lys Leu Pro Tyr Leu Gln Ala Ile
345 350 355

Cys Lys Glu Ala Phe Arg Lys His Pro Ser Thr Pro Leu Asn Leu Pro
360 365 370 375

Arg Ile Ala Ser Gln Ala Cys Glu Val Asn Gly His Tyr Ile Pro Lys
380 385 390

Gly Thr Arg Leu Ser Val Asn Ile Trp Ala Ile Gly Arg Asp Pro Ser
395 400 405

Val Trp Glu Asn Pro Asn Glu Phe Asn Pro Asp Arg Phe Leu Glu Arg
410 415 420

Lys Asn Ala Lys Ile Asp Pro Arg Gly Asn Asp Phe Glu Leu Ile Pro
425 430 435

Phe Gly Ala Gly Arg Arg Ile Cys Ala Gly Thr Arg Leu Gly Ile Leu
440 445 450 455

Leu Val Glu Tyr Ile Leu Gly Thr Leu Val His Ser Phe Val Trp Glu
460 465 470

Leu Pro Ser Ser Val Ile Glu Leu Asn Met Asp Glu Ser Phe Gly Leu
475 480 485

Ala Leu Gln Lys Ala Val Pro Leu Ala Ala Met Val Thr Pro Arg Leu
490 495 500

Pro Leu His Ile Tyr Ser Pro
505 510

(2) INFORMATION FOR SEQ ID NO: 67 :

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 523 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Campanula medium

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 180 to 1748
- (C) IDENTIFICATION METHOD: by experiment

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 67

-continued

Met Ser Ile Asp Ile Ser Thr Leu Phe Tyr Glu Leu Val Ala Ala Ile
1 5 10 15

Ser Leu Tyr Leu Ala Thr Tyr Ser Phe Ile Arg Phe Leu Phe Lys Pro
20 25 30

Ser His His His His Leu Pro Pro Gly Pro Thr Gly Trp Pro Ile Ile
35 40 45

Gly Ala Leu Pro Leu Leu Gly Thr Met Pro His Val Ser Leu Ala Asp
50 55 60

Met Ala Val Lys Tyr Gly Pro Ile Met Tyr Leu Lys Leu Gly Ser Lys
65 70 75 80

Gly Thr Val Val Ala Ser Asn Pro Lys Ala Ala Arg Ala Phe Leu Lys
85 90 95

Thr His Asp Ala Asn Phe Ser Asn Arg Pro Ile Asp Gly Gly Pro Thr
100 105 110

Tyr Leu Ala Tyr Asn Ala Gln Asp Met Val Phe Ala Glu Tyr Gly Pro
115 120 125

Lys Trp Lys Leu Leu Arg Lys Leu Cys Ser Leu His Met Leu Gly Pro
130 135 140

Lys Ala Leu Glu Asp Trp Ala His Val Lys Val Ser Glu Val Gly His
145 150 155 160

Met Leu Lys Glu Met Tyr Glu Gln Ser Ser Lys Ser Val Pro Val Pro
165 170 175

Val Val Val Pro Glu Met Leu Thr Tyr Ala Met Ala Asn Met Ile Gly
180 185 190

Arg Ile Ile Leu Ser Arg Arg Pro Phe Val Ile Thr Ser Lys Leu Asp
195 200 205

Ser Ser Ala Ser Ala Ser Ala Ser Val Ser Glu Phe Gln Tyr Met Val
210 215 220

Met Glu Leu Met Arg Met Ala Gly Leu Phe Asn Ile Gly Asp Phe Ile
225 230 235 240

Pro Tyr Ile Ala Trp Met Asp Leu Gln Gly Ile Gln Arg Asp Met Lys
245 250 255

Val Ile Gln Lys Lys Phe Asp Val Leu Leu Asn Lys Met Ile Lys Glu
260 265 270

His Thr Glu Ser Ala His Asp Arg Lys Asp Asn Pro Asp Phe Leu Asp
275 280 285

Ile Leu Met Ala Ala Thr Gln Glu Asn Thr Glu Gly Ile Gln Leu Asn
290 295 300

Leu Val Asn Val Lys Ala Leu Leu Leu Asp Leu Phe Thr Ala Gly Thr
305 310 315 320

Asp Thr Ser Ser Ser Val Ile Glu Trp Ala Leu Ala Glu Met Leu Asn
325 330 335

His Arg Gln Ile Leu Asn Arg Ala His Glu Glu Met Asp Gln Val Ile
340 345 350

Gly Arg Asn Arg Arg Leu Glu Gln Ser Asp Ile Pro Asn Leu Pro Tyr
355 360 365

Phe Gln Ala Ile Cys Lys Glu Thr Phe Arg Lys His Pro Ser Thr Pro
370 375 380

Leu Asn Leu Pro Arg Ile Ser Thr Glu Ala Cys Glu Val Asp Gly Phe
385 390 395 400

His Ile Pro Lys Asn Thr Arg Leu Ile Val Asn Ile Trp Ala Ile Gly

-continued

| | | | |
|---|-----|-----|-----|
| | 405 | 410 | 415 |
| Arg Asp Pro Lys Val Trp Glu Asn Pro Leu Asp Phe Thr Pro Glu Arg | 420 | 425 | 430 |
| Phe Leu Ser Glu Lys His Ala Lys Ile Asp Pro Arg Gly Asn His Phe | 435 | 440 | 445 |
| Glu Leu Ile Pro Phe Gly Ala Gly Arg Arg Ile Cys Ala Gly Ala Arg | 450 | 455 | 460 |
| Met Gly Ala Ala Ser Val Glu Tyr Ile Leu Gly Thr Leu Val His Ser | 465 | 470 | 475 |
| Phe Asp Trp Lys Leu Pro Asp Gly Val Val Glu Val Asn Met Glu Glu | 485 | 490 | 495 |
| Ser Phe Gly Ile Ala Leu Gln Lys Lys Val Pro Leu Ser Ala Ile Val | 500 | 505 | 510 |
| Thr Pro Arg Leu Pro Pro Ser Ser Tyr Thr Val | 515 | 520 | |

1) A DNA encoding a polypeptide which has flavonoid-3',5'-hydroxylase activity and which is represented by the amino acid sequence shown by SEQ ID NO: 1, 63 or 64, or a DNA which hybridizes with said DNA.

2) The DNA according to claim (1), wherein a part of the nucleotide sequence of said DNA is deleted or replaced by another nucleotide sequence.

3) A recombinant DNA composed of a vector DNA and the DNA of claim (1) or claim (2) which is inserted in the vector DNA.

4) A plant or plant cell which carries the recombinant DNA according to claim (3).

5) The plant or plant cell according to claim (4), wherein said plant belongs to the genus *Rosa*, *Nicotiana*, *Petunia*, or *Dianthus*.

6) A DNA which hybridizes with a DNA represented by the nucleotide sequence shown by SEQ ID NO: 1, 63 or 64 in 2xSSC (0.3 M sodium chloride, 0.03 M sodium citrate, pH 7.0) at 50° C.

7) A method for producing a plant, which comprises: introducing a recombinant DNA composed of a vector DNA

fragment and a DNA fragment which encodes a polypeptide having flavonoid-3',5'-hydroxylase activity into a plant; breeding a plant which can express a pigment based on the genetic information of the DNA encoding said polypeptide; and harvesting said plant thus obtained.

8) The method for producing a plant according to claim (7), wherein said DNA fragment which encodes the polypeptide having flavonoid-3',5'-hydroxylase activity is the DNA of claim (1) or claim (2).

9) A DNA which has the nucleotide sequence shown by any of SEQ ID NO: 2 to 29.

10) A DNA which has a sequence comprising a sequence identical with the eight-nucleotide sequence from the 3'-terminus in the sequence of the DNA of claim (9).

11) A method for amplifying and isolating a gene fragment which encodes the amino acid sequence of the heme-binding region of cytochrome P450 enzyme, by polymerase chain reaction (PCR) using the DNA of claim (9) or claim (10) as primers.

* * * * *