



US005457280A

United States Patent [19]

[11] **Patent Number:** **5,457,280**

Eby

[45] **Date of Patent:** **Oct. 10, 1995**

- [54] **SOYBEAN CULTIVAR 0585540**
- [75] Inventor: **William H. Eby**, Adel, Iowa
- [73] Assignee: **Stine Seed Farm, Inc.**, Adel, Iowa
- [21] Appl. No.: **229,245**
- [22] Filed: **Apr. 18, 1994**
- [51] Int. Cl.⁶ **A01H 4/00**; A01H 5/00;
A01H 5/10; C12N 5/04
- [52] U.S. Cl. **800/200**; 800/255; 800/DIG. 26;
435/240.1; 435/240.4; 435/240.49; 435/240.5;
47/58; 47/DIG. 1
- [58] **Field of Search** 435/172.1, 172.3,
435/240.1, 240.4, 240.49, 240.5; 800/200,
250, DIG. 9, DIG. 23, DIG. 26, 255; 47/58.03

- [56] **References Cited**
- U.S. PATENT DOCUMENTS
- 3,903,645 9/1975 Bradner 47/58

OTHER PUBLICATIONS

Luedders. 1977. *Crop Science*. 17:971-972.
 Kilen. 1980. *Crop Science*. 20:261-262.
 Nelson et al. 1984. *Crop Science*. 24:549-553.

Barwale et al. 1987. *Plant Cell Reports*. 6:365-368.
 Barwale. 1986. *Planta*. 167:473-481.
 Larkin. 1987. *Iowa State Journal of Research*.
 61(4):393-434.
 Palmer et al. 1987. In *Soybeans: Improvement, Production,
 and Uses*. 2nd Ed. Wilcox, ed. Ch. 5:135-207.
 Fehr. 1987. In *Soybeans: Improvement, Production, and
 Uses*. 2nd Ed. Wilcox, ed. Ch. 7:249-293.

Primary Examiner—Gary Benzion
Assistant Examiner—Erich E. Veitenheimer
Attorney, Agent, or Firm—Henderson & Sturm

- [57] **ABSTRACT**
- A novel soybean cultivar, designated 0585540, is disclosed. The invention relates to the seeds of soybean cultivar 0585540, to the plants of soybean 0585540 and to methods for producing a soybean plant produced by crossing the cultivar 0585540 with itself or another soybean variety. The invention further relates to hybrid soybean seeds and plants produced by crossing the cultivar 0585540 with another soybean cultivar.

8 Claims, No Drawings

SOYBEAN CULTIVAR 0585540

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinctive soybean cultivar, designated 0585540. There are numerous steps in the development of any novel, desirable plant germplasm. Plant breeding begins with the analysis and definition of problems and weaknesses of the current germplasm, the establishment of program goals, and the definition of specific breeding objectives. The next step is selection of germplasm that possess the traits to meet the program goals. The goal is to combine in a single variety an improved combination of desirable traits from the parental germplasm. These important traits may include higher seed yield, resistance to diseases and insects, better stems and roots, tolerance to drought and heat, and better agronomic quality.

Choice of breeding or selection methods depends on the mode of plant reproduction, the heritability of the trait(s) being improved, and the type of cultivar used commercially (e.g., F_1 hybrid cultivar, pureline cultivar, etc.). For highly heritable traits, a choice of superior individual plants evaluated at a single location will be effective, whereas for traits with low heritability, selection should be based on mean values obtained from replicated evaluations of families of related plants. Popular selection methods commonly include pedigree selection, modified pedigree selection, mass selection, and recurrent selection.

The complexity of inheritance influences choice of the breeding method. Backcross breeding is used to transfer one or a few favorable genes for a highly heritable trait into a desirable cultivar. This approach has been used extensively for breeding disease-resistant cultivars. Various recurrent selection techniques are used to improve quantitatively inherited traits controlled by numerous genes. The use of recurrent selection in self-pollinating crops depends on the ease of pollination, the frequency of successful hybrids from each pollination, and the number of hybrid offspring from each successful cross.

Each breeding program should include a periodic, objective evaluation of the efficiency of the breeding procedure. Evaluation criteria vary depending on the goal and objectives, but should include gain from selection per year based on comparisons to an appropriate standard, overall value of the advanced breeding lines, and number of successful cultivars produced per unit of input (e.g., per year, per dollar expended, etc.).

Promising advanced breeding lines are thoroughly tested and compared to appropriate standards in environments representative of the commercial target area(s) for three or more years. The best lines are candidates for new commercial cultivars; those still deficient in a few traits may be used as parents to produce new populations for further selection.

These processes, which lead to the final step of marketing and distribution, usually take from eight to 12 years from the time the first cross is made. Therefore, development of new cultivars is a time-consuming process that requires precise forward planning, efficient use of resources, and a minimum of changes in direction.

A most difficult task is the identification of individuals that are genetically superior, because for most traits the true genotypic value is masked by other confounding plant traits or environmental factors. One method of identifying a superior plant is to observe its performance relative to other

experimental plants and to a widely grown standard cultivar. If a single observation is inconclusive, replicated observations provide a better estimate of its genetic worth.

The goal of plant breeding is to develop new, unique and superior soybean cultivars. The breeder initially selects and crosses two or more parental lines, followed by repeated selfing and selection, producing many new genetic combinations. The breeder can theoretically generate billions of different genetic combinations via crossing, selfing and mutations. The breeder has no direct control at the cellular level. Therefore, two breeders will never develop the same line, or even very similar lines, having the same soybean traits.

Each year, the plant breeder selects the germplasm to advance to the next generation. This germplasm is grown under unique and different geographical, climatic and soil conditions, and further selections are then made, during and at the end of the growing season. The cultivars which are developed are unpredictable. This unpredictability is because the breeder's selection occurs in unique environments, with no control at the DNA level (using conventional breeding procedures), and with millions of different possible genetic combinations being generated. A breeder of ordinary skill in the art cannot predict the final resulting lines he develops, except possibly in a very gross and general fashion. The same breeder cannot produce the same cultivar twice by using the exact same original parents and the same selection techniques. This unpredictability results in the expenditure of large research monies to develop a superior new soybean cultivar.

Pedigree breeding and recurrent selection breeding methods are used to develop cultivars from breeding populations. Breeding programs combine desirable traits from two or more cultivars or various broad-based sources into breeding pools from which cultivars are developed by selfing and selection of desired phenotypes. The new cultivars are evaluated to determine which have commercial potential.

Pedigree breeding is used commonly for the improvement of self-pollinating crops. Two parents which possess favorable, complementary traits are crossed to produce an F_1 . An F_2 population is produced by selfing one or several F_1 's or by intercrossing two F_1 's (sib mating). Selection of the best individuals may begin in the F_2 population; then, beginning in the F_3 , the best individuals in the best families are selected. Replicated testing of families can begin in the F_4 generation to improve the effectiveness of selection for traits with low heritability. At an advanced stage of inbreeding (i.e., F_6 and F_7), the best lines or mixtures of phenotypically similar lines are tested for potential release as new cultivars.

Mass and recurrent selections can be used to improve populations of either self- or cross-pollinating crops. A genetically variable population of heterozygous individuals is either identified or created by intercrossing several different parents. The best plants are selected based on individual superiority, outstanding progeny, or excellent combining ability. The selected plants are intercrossed to produce a new population in which further cycles of selection are continued.

Backcross breeding has been used to transfer genes for a simply inherited, highly heritable trait into a desirable homozygous cultivar or inbred line which is the recurrent parent. The source of the trait to be transferred is called the donor parent. The resulting plant is expected to have the attributes of the recurrent parent (e.g., cultivar) and the desirable trait transferred from the donor parent. After the initial cross, individuals possessing the phenotype of the

donor parent are selected and repeatedly crossed (back-crossed) to the recurrent parent. The resulting plant is expected to have the attributes of the recurrent parent (e.g., cultivar) and the desirable trait transferred from the donor parent.

The single-seed descent procedure in the strict sense refers to planting a segregating population, harvesting a sample of one seed per plant, and using the one-seed sample to plant the next generation. When the population has been advanced from the F_2 to the desired level of inbreeding, the plants from which lines are derived will each trace to different F_2 individuals. The number of plants in a population declines each generation due to failure of some seeds to germinate or some plants to produce at least one seed. As a result, not all of the F_2 plants originally sampled in the population will be represented by a progeny when generation advance is completed.

In a multiple-seed procedure, soybean breeders commonly harvest one or more pods from each plant in a population and thresh them together to form a bulk. Part of the bulk is used to plant the next generation and part is put in reserve. The procedure has been referred to as modified single-seed descent or the pod-bulk technique.

The multiple-seed procedure has been used to save labor at harvest. It is considerably faster to thresh pods with a machine than to remove one seed from each by hand for the single-seed procedure. The multiple-seed procedure also makes it possible to plant the same number of seeds of a population each generation of inbreeding. Enough seeds are harvested to make up for those plants that did not germinate or produce seed.

Descriptions of other breeding methods that are commonly used for different traits and crops can be found in one of several reference books (e.g., Allard, 1960; Simmonds, 1979; Sneep et al., 1979; Fehr, 1987).

Proper testing should detect any major faults and establish the level of superiority or improvement over current cultivars. In addition to showing superior performance, there must be a demand for a new cultivar that is compatible with industry standards or which creates a new market. The introduction of a new cultivar will incur additional costs to the seed producer, the grower, processor and consumer; for special advertising and marketing, altered seed and commercial production practices, and new product utilization. The testing preceding release of a new cultivar should take into consideration research and development costs as well as technical superiority of the final cultivar. For seed-propagated cultivars, it must be feasible to produce seed easily and economically.

Soybean, *Glycine max* (L), is an important and valuable field crop. Thus, a continuing goal of plant breeders is to develop stable, high yielding soybean cultivars that are agronomically sound. The reasons for this goal are obviously to maximize the amount of grain produced on the land used and to supply food for both animals and humans. To accomplish this goal, the soybean breeder must select and develop soybean plants that have the traits that result in superior cultivars.

SUMMARY OF THE INVENTION

According to the invention, there is provided a novel soybean cultivar, designated 0585540. This invention thus relates to the seeds of soybean cultivar 0585540, to the plants of soybean 0585540 and to methods for producing a soybean plant produced by crossing the soybean 0585540 with itself or another soybean line.

DEFINITIONS

In the description and tables which follow, a number of terms are used. In order to provide a clear and consistent understanding of the specification and claims, including the scope to be given such terms, the following definitions are

provided:

Maturity. Plants are considered mature when 95 % of the pods have reached their mature color.

Seed Yield (Bushels/Acre). The yield in bushels/acre is the actual yield of the grain at harvest.

Lodging Resistance. Lodging is rated on a scale of 1 to 9. A score of 9 indicates erect plants. A score of 5 indicates plants are leaning at a 45° angle in relation to the ground and a score of 1 indicates plants are laying on the ground.

Phytophthora Tolerance. Tolerance to Phytophthora root rot is rated on a scale of 1 to 9, with a score of 9 being the best or highest tolerance ranging down to a score of 1 which indicates the plants have no tolerance to phytophthora.

Emergence. This score indicates the ability of the seed to emerge when planted 3" deep in sand and with a controlled temperature of 25° C. The number of plants that emerge each day are counted. Based on this data, each genotype is given a 1 to 9 score based on its rate of emergence and percent of emergence. A score of 9 indicates an excellent rate and percent of emergence, an intermediate score of 5 indicates average ratings and a 1 score indicates a very poor rate and percent of emergence.

Iron-Deficiency Chlorosis. Plants are scored 1 to 9 based on visual observations. A score of 1 indicates the plants are dead or dying caused by iron-deficiency chlorosis, a score of 5 means plants have intermediate health with some leaf yellowing and a score of 9 means no stunting of the plants or yellowing of the leaves.

Brown Stem Rot. This is a visual disease score from 1 to 9 comparing all genotypes in a given test. The score is based on leaf symptoms of yellowing and necrosis caused by brown stem rot. A score of 9 indicates no symptoms. Visual scores range down to a score of 1 which indicates severe symptoms of leaf yellowing and necrosis.

Shattering. The amount of pod dehiscence prior to harvest. Pod dehiscence involves seeds falling from the pods to the soil. This is a visual score from 1 to 9 comparing all genotypes within a given test. A score of 9 means pods have not opened and no seeds have fallen out. A score of 5 indicates approximately 50% of the pods have opened, with seeds falling to the ground and a score of 1 indicates 100% of the pods are opened.

Plant Height. Plant height is taken from the top of soil to top of the plant and is measured in inches.

DETAILED DESCRIPTION OF THE INVENTION

Soybean cultivar 0585540 has superior characteristics and was developed from the cross Sherman x 63639. F_3 seed was advanced by the single seed decent method of breeding. In the fall of 1988, F_3 derived single plants were selected and harvested individually. These lines were tested in experimental plots in 1989 at Adel, Iowa. The highest yielding lines were retained for additional testing in 1990. A second year of testing was conducted during 1990 at four midwest locations where the lines were adapted. In the fall of 1990, 20 F_3 derived single plants of 05855 were selected and harvested individually.

During 1991 these 20 plant rows were evaluated for maturity and other visual characters. Also during 1991, 05855 was evaluated at ten midwest locations. 05855 was advanced to the 1992 Elite Trial where it was evaluated at eleven midwest locations. Six plant row selections were advanced to the 1992 Plant Row Yield Trial.

Based on the 1991-1992 Elite data on 05855 and plant row Yield Trial data, selection 0585540 was advanced to the 1993 Elite Trial which included ten midwest locations as

TABLE 2

COMPARISON OF 0585540 VERSUS STINE 3260					
0585540					
	OIL %	PROTEIN %	LODGING	HEIGHT	SEED SIZE GR/100 SEEDS
0585540	18.6	35.5	6.5	102	16.9
Stine 3260	19.0	35.9	6.2	99	15.8

10

DEPOSIT INFORMATION

Seeds of 0585540 have been placed on deposit with the American Type Culture Collection (ATCC), Rockville, Md. 20852, under Deposit Accession Number 75732 on Apr. 7, 1994.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity and understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the invention, as limited only by the scope of the appended claims.

What is claimed is:

1. A soybean seed designated 0585540 having ATCC accession No. 75732.
2. A soybean plant produced by growing the seed of claim 1.
3. Pollen of the plant of claim 2.
4. An ovule of the plant of claim 2.
5. A soybean plant having all of the physiological and morphological characteristics of the soybean plant designated 0585540, wherein seed of 0585540 has been deposited as ATCC accession No. 75732.

6. A tissue culture comprising regenerable cells of the plant of claim 2.

7. A soybean plant with all of the physiological and morphological characteristics of the soybean plant designated 0585540, seed of which has been deposited as ATCC accession No. 75732, said soybean plant produced by the tissue culture process using soybean plant 0585540 as the starting material of such a process.

8. A first generation (F₁) hybrid soybean plant produced by growing said hybrid soybean seed, said seed produced by the process of:

- (a) planting in pollinating proximity seeds of soybean cultivar 0585540, seed of which has been deposited as ATCC accession No. 75732, and another soybean genotype;
- (b) cultivating soybean plants resulting from said seeds until said plants bear flowers;
- (c) emasculating the male flowers of the plants of either soybean genotype;
- (d) allowing cross pollination to occur between said soybean genotypes; and,
- (e) harvesting seeds produced on said emasculated plants of the soybean line.

35

* * * * *

40

45

50

55

60

65