

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization

International Bureau

(43) International Publication Date
19 April 2018 (19.04.2018)



(10) International Publication Number
WO 2018/069108 A1

(51) International Patent Classification:

A01N 43/90 (2006.01) *A01N 43/713* (2006.01)
A01N 43/42 (2006.01) *A01N 43/72* (2006.01)
A01N 43/653 (2006.01) *A01P 3/00* (2006.01)

(21) International Application Number:

PCT/EP2017/075189

(22) International Filing Date:

04 October 2017 (04.10.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

16192985.6 10 October 2016 (10.10.2016) EP

(71) Applicant: **BASF SE** [DE/DE]; Carl-Bosch-Strasse 38,
67056 Ludwigshafen am Rhein (DE).

(72) Inventor: **GEWEHR, Markus**; Friedrich-Ebert-Strasse
18, 69502 Hemsbach (DE).

(74) Agent: **BASF IP ASSOCIATION**; BASF SE, G-FLP -
C006, 67056 Ludwigshafen (DE).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: PESTICIDAL MIXTURES

(57) Abstract: The present invention relates to pesticidal mixtures comprising tetramic acids as compound I and a compound II selected from the group consisting of mefentrifluconazole, picarbutrazox, 2-[2-fluoro-6-[(8-fluoro-2-methyl-3-quinolyl)oxy]phenyl]propan-2-ol, 2-[2-[(7,8-difluoro-2-methyl-3-quinolyl)oxy]-6-fluoro-phenyl]propan-2-ol and 9-fluoro-2,2-dimethyl-5-(3-quinolyl)-3H 1,4-benzoxazepine.




WO 2018/069108 A1

Pesticidal mixtures

Description

5 The present invention relates to pesticidal mixtures comprising

1) a compound of formula I



(I), as compound I;

10

wherein R¹ is hydrogen, methyl, methoxy or ethoxy; and
R² and R⁴ are methyl, R³ is methyl or chlorine

and

15 2) one fungicidal compound II, wherein compound II is selected from the group consisting of mefen-trifluconazole, picarbutrazox, 2-[2-fluoro-6-[(8-fluoro-2-methyl-3-quinolyl)oxy]phenyl]propan-2-ol, 2-[2-[(7,8-difluoro-2-methyl-3-quinolyl)oxy]-6-fluoro-phenyl]propan-2-ol, 9-fluoro-2,2-dimethyl-5-(3-quinolyl)-3H 1,4-benzoxazepine.

20 The above-referred mixtures are herein below also referred as "inventive mixtures".

Moreover, the invention relates to a method for controlling pests, this refers to includes animal pests and harmful fungi, using the inventive mixtures and to the use of compound I and compound II for preparing such mixtures, and also to compositions comprising such mixtures.

25

Moreover, the invention relates to a method for controlling harmful fungi, using the inventive mixtures of compound I and II for preparing such mixtures, and also to compositions comprising such mixtures.

30 Additionally, the present invention also comprises a method for protection of plant propagation material (preferably seed) from harmful pests, such as fungi or animal pests (insects, arachnids or nematodes) comprising contacting the plant propagation materials (preferably seeds) with an inventive mixture in pesticidally effective amounts

35 The term "plant propagation material" is to be understood to denote all the generative parts of the plant such as seeds and vegetative plant material such as cuttings and tubers (e. g. pota-

toes), which can be used for the multiplication of the plant. This includes seeds, roots, fruits, tubers, bulbs, rhizomes, shoots, sprouts and other parts of plants, including seedlings and young plants, which are to be transplanted after germination or after emergence from soil. These young plants may also be protected before transplantation by a total or partial treatment
5 by immersion or pouring. In a particular preferred embodiment, the term propagation material denotes seeds.

Additionally, the present invention also comprises a method for protection of plant propagation material (preferably seed) from harmful fungi comprising contacting the plant propagation mate-
10 rials (preferably seeds) with the inventive mixture in pesticidally effective amounts.

Moreover, the invention relates to a method for controlling harmful fungi using the inventive mix-
tures and to the use of the compounds present in the inventive mixtures for preparing such mix-
tures, and also to compositions comprising such mixtures.

The present invention further relates to plant-protecting active ingredient mixtures having syner-
15 gistically enhanced action of improving the health of plants and to a method of applying such inventive mixtures to the plants.

Compounds I and analogues as well as their pesticidal action and methods for producing them
20 are known e.g. from WO 2014/191271 and WO 2009/049851. Compounds II as well as their pesticidal action and methods for producing them are generally known from, e.g. from the e-Pesticide Manual V5.2 (ISBN 978 1 901396 85 0) (2008-2011), WO2012143127, WO 14/060502, WO 14/060521, WO 2015036379 AI, WO13127704, WO 2009115557,
25 WO2012045798, WO 2013047441 und WO2011081174, WO 2013047441 and WO2011081174.

Mixtures of compound I with selected fungicides and insecticides are known from
30 WO16001120: WO 2013079564 , WO 2013107795 , WO 2013107796 , WO 2013107793 and WO 2013107794. However, the combinations set forth above are not described therein.

One typical problem arising in the field of pest control lies in the need to reduce the dosage
rates of the active ingredient in order to reduce or avoid unfavorable environmental or toxicolog-
ical effects whilst still allowing effective pest control.

35 In regard to the instant invention the term pests embrace harmful fungi and animal pests.

Another problem encountered concerns the need to have available pest control agents which
are effective against a broad spectrum of harmful fungi and harmful animal pests.

40 There also exists the need for pest control agents that combine knock-down activity with prolonged control, that is, fast action with long lasting action.

Another difficulty in relation to the use of pesticides is that the repeated and exclusive application of an individual pesticidal compound leads in many cases to a rapid selection of pests, that means animal pests, and harmful fungi, which have developed natural or adapted resistance against the active compound in question. Therefore there is a need for pest control agents that help prevent or overcome resistance.

Another problem underlying the present invention is the desire for compositions that improve plants, a process which is commonly and hereinafter referred to as "plant health".

The term plant health comprises various sorts of improvements of plants that are not connected to the control of pests. For example, advantageous properties that may be mentioned are improved crop characteristics including: emergence, crop yields, protein content, oil content, starch content, more developed root system (improved root growth), improved stress tolerance (e.g. against drought, heat, salt, UV, water, cold), reduced ethylene (reduced production and/or inhibition of reception), tillering increase, increase in plant height, bigger leaf blade, less dead basal leaves, stronger tillers, greener leaf color, pigment content, photosynthetic activity, less input needed (such as fertilizers or water), less seeds needed, more productive tillers, earlier flowering, early grain maturity, less plant verse (lodging), increased shoot growth, enhanced plant vigor, increased plant stand and early and better germination; or any other advantages familiar to a person skilled in the art.

It was therefore an object of the present invention to provide pesticidal mixtures which solve the problems of reducing the dosage rate and / or enhancing the spectrum of activity and / or combining knock-down activity with prolonged control and / or to resistance management and/or promoting the health of plants.

We have found that this object is in part or in whole achieved by the mixtures comprising the active compounds defined in the outset.

Especially, it has been found that the mixtures as defined in the outset show markedly enhanced action against pests compared to the control rates that are possible with the individual compounds and/or is suitable for improving the health of plants when applied to plants, parts of plants, seeds, or at their locus of growth.

It has been found that the action of the inventive mixtures comprising compound I and compound II goes far beyond the fungicidal and/or insecticidal and/or plant health improving action of the active compounds present in the mixture alone (synergistic action).

Moreover, we have found that simultaneous, that is joint or separate, application of the compound I and the compound II or successive application of the compound I and compound II allows enhanced control of harmful fungi, compared to the control rates that are possible with the individual compounds (synergistic mixtures).

Moreover, we have found that simultaneous, that is joint or separate, application of the compound I and the compound II or successive application of the compound I and the compound II allows enhanced control of harmful animal pests, compared to the control rates that are possible with the individual compounds (synergistic mixtures).

5


Moreover, we have found that simultaneous, that is joint or separate, application of the compound I and the compound II or successive application of the compound I and the compound II provides enhanced plant health effects compared to the plant health effects that are possible with the individual compounds.

10

The ratio by weight of compound I and compound II in binary mixtures is from 20000:1 to 1:20000, from 500:1 to 1:500, preferably from 100:1 to 1:100 more preferably from 50:1 to 1:50, most preferably from 20:1 to 1:20, including also ratios from 10:1 to 1:10, 1:5 to 5:1, or 1:1.

15 As set forth above, the present specifically relates to the inventive mixtures as defined above, namely pesticidal mixtures comprising

1) a compound of formula I



20

(I), as compound I;

wherein R¹ is hydrogen, methyl, methoxy or ethoxy; and
R² and R⁴ are methyl, R³ is methyl or chlorine

and

2) one fungicidal compound II, wherein compound II is selected from the group consisting of
25 mefentrifluconazole, picarbutrazox, 2-[2-fluoro-6-[(8-fluoro-2-methyl-3-quinolyl)oxy]phenyl]propan-2-ol, 2-[2-[(7,8-difluoro-2-methyl-3-quinolyl)oxy]-6-fluorophenyl]propan-2-ol, 9-fluoro-2,2-dimethyl-5-(3-quinolyl)-3H 1,4-benzoxazepine.

30 Preferably, compound I comprises the compounds listed in table 1 below, wherein in the tables 3 to 16 compounds of formula I are abbreviated as indicated in table 1:

Table 1

Compound I	Abbreviation
[3-(4-chloro-2,6-dimethylphenyl)-8-methoxy-1-methyl-2-oxo-1,8-diazaspiro[4.5]dec-3-en-4-yl] carbonic acid ethyl ester	I-1
[3-(4-chloro-2,6-dimethylphenyl)-8-methoxy-2-oxo-1,8-diazaspiro[4.5]dec-3-en-4-yl] carbonic acid ethyl ester	I-2

Compound I	Abbreviation
[3-(4-chloro-2,6-dimethyl-phenyl)-1,8-dimethoxy-2-oxo-1,8-diazaspiro[4.5]dec-3-en-4-yl] carbonic acid ethyl ester	I-3
[3-(4-chloro-2,6-dimethyl-phenyl)-1-ethoxy-8-methoxy-2-oxo-1,8-diazaspiro[4.5]dec-3-en-4-yl] carbonic acid ethyl ester	I-4
[1-ethoxy-8-methoxy-2-oxo-3-(2,4,6-trimethylphenyl)-1,8-diazaspiro[4.5]dec-3-en-4-yl] carbonic acid ethyl ester	I-5
[1,8-dimethoxy-2-oxo-3-(2,4,6-trimethylphenyl)-1,8-diazaspiro[4.5]dec-3-en-4-yl] carbonic acid ethyl ester	I-6
[8-methoxy-1-methyl-2-oxo-3-(2,4,6-trimethylphenyl)-1,8-diazaspiro[4.5]dec-3-en-4-yl] carbonic acid ethyl ester	I-7
[8-methoxy-2-oxo-3-(2,4,6-trimethylphenyl)-1,8-diazaspiro[4.5]dec-3-en-4-yl] carbonic acid ethyl ester	I-8

In the tables 3 to 16 compounds II are abbreviated as indicated in table 2:

Table 2

Compound II	Abbreviation
mefentrifluconazole	II-1
picarbutrazox	II-2
2-[2-fluoro-6-[(8-fluoro-2-methyl-3-quinolyl)oxy]phenyl]propan-2-ol	II-3
9-fluoro-2,2-dimethyl-5-(3-quinolyl)-3H 1,4-benzoxazepine	II-4
2-[2-[(7,8-difluoro-2-methyl-3-quinolyl)oxy]-6-fluoro-phenyl]propan-2-ol	II-5

- 5 The present invention furthermore relates to mixtures of a compound of formula I and mefentrifluconazole, wherein the mixtures as set forth in table 3 are herein preferred:

Table 3

No	Compound I	Compound II
M5-1	I-1	II-1
M5-2	I-2	II-1
M5-3	I-3	II-1
M5-4	I-4	II-1
M5-5	I-5	II-1

No	Compound I	Compound II
M5-1	I-6	II-1
M5-2	I-7	II-1
M5-3	I-8	II-1

- 10 The present invention furthermore relates to mixtures of a compound of formula I and picarbutrazox, wherein the mixtures as set forth in table 4 are herein preferred:

Table 4

No	Compound I	Compound II
M6-1	I-1	II-2
M6-2	I-2	II-2
M6-3	I-3	II-2
M6-4	I-4	II-2

No	Compound I	Compound II
M6-1	I-6	II-2
M6-2	I-7	II-2
M6-3	I-8	II-2

No	Compound I	Compound II
M6-5	I-5	II-2

No	Compound I	Compound II

The present invention furthermore relates to mixtures of a compound of formula I and 2-[2-fluoro-6-[(8-fluoro-2-methyl-3-quinolyl)oxy]phenyl]propan-2-ol, wherein the mixtures as set forth in table 5 are herein preferred:

5

Table 5

No	Compound I	Compound II
M14-1	I-1	II-3
M14-2	I-2	II-3
M14-3	I-3	II-3
M14-4	I-4	II-3
M14-5	I-5	II-3

No	Compound I	Compound II
M14-1	I-6	II-3
M14-2	I-7	II-3
M14-3	I-8	II-3

The present invention furthermore relates to mixtures of a compound of formula I and 9-fluoro-2,2-dimethyl-5-(3-quinolyl)-3H 1,4-benzoxazepine, wherein the mixtures as set forth in table 6 are herein preferred:

10

Table 6

No	Compound I	Compound II
M15-1	I-1	II-4
M15-2	I-2	II-4
M15-3	I-3	II-4
M15-4	I-4	II-4
M15-5	I-5	II-4

No	Compound I	Compound II
M15-1	I-6	II-4
M15-2	I-7	II-4
M15-3	I-8	II-4

The present invention furthermore relates to mixtures of a compound of formula I and 2-[2-[(7,8-difluoro-2-methyl-3-quinolyl)oxy]-6-fluoro-phenyl]propan-2-ol, wherein the mixtures as set forth in table 7 are herein preferred:

15

Table 7

No	Compound I	Compound II
M16-1	I-1	II-5
M16-2	I-2	II-5
M16-3	I-3	II-5
M16-4	I-4	II-5

No	Compound I	Compound II
M16-5	I-5	II-5
M16-1	I-6	II-5
M16-2	I-7	II-5
M16-3	I-8	II-5

20 The inventive mixtures can further contain one or more insecticides, fungicides, herbicides.

The inventive mixtures can be converted into customary types of agrochemical compositions, e. g. solutions, emulsions, suspensions, dusts, powders, pastes, granules, pressings, capsules, and mixtures thereof. Examples for composition types are suspensions (e.g. SC, OD, FS), emulsifiable concentrates (e.g. EC), emulsions (e.g. EW, EO, ES, ME), capsules (e.g. CS, ZC), pastes, pastilles, wettable powders or dusts (e.g. WP, SP, WS, DP, DS), pressings (e.g.

25

BR, TB, DT), granules (e.g. WG, SG, GR, FG, GG, MG), insecticidal articles (e.g. LN), as well as gel formulations for the treatment of plant propagation materials such as seeds (e.g. GF). These and further compositions types are defined in the "Catalogue of pesticide formulation types and international coding system", Technical Monograph No. 2, 6th Ed. May 2008, CropLife International.

The compositions are prepared in a known manner, such as described by Mollet and Grubemann, Formulation technology, Wiley VCH, Weinheim, 2001; or Knowles, New developments in crop protection product formulation, Agrow Reports DS243, T&F Informa, London, 2005.

Suitable auxiliaries are solvents, liquid carriers, solid carriers or fillers, surfactants, dispersants, emulsifiers, wetters, adjuvants, solubilizers, penetration enhancers, protective colloids, adhesion agents, thickeners, humectants, repellents, attractants, feeding stimulants, compatibilizers, bactericides, anti-freezing agents, anti-foaming agents, colorants, tackifiers and binders.

Suitable solvents and liquid carriers are water and organic solvents, such as mineral oil fractions of medium to high boiling point, e.g. kerosene, diesel oil; oils of vegetable or animal origin; aliphatic, cyclic and aromatic hydrocarbons, e. g. toluene, paraffin, tetrahydronaphthalene, alkylated naphthalenes; alcohols, e.g. ethanol, propanol, butanol, benzylalcohol, cyclohexanol; glycols; DMSO; ketones, e.g. cyclohexanone; esters, e.g. lactates, carbonates, fatty acid esters, gamma-butyrolactone; fatty acids; phosphonates; amines; amides, e.g. N-methylpyrrolidone, fatty acid dimethylamides; and mixtures thereof.

Suitable solid carriers or fillers are mineral earths, e.g. silicates, silica gels, talc, kaolins, limestone, lime, chalk, clays, dolomite, diatomaceous earth, bentonite, calcium sulfate, magnesium sulfate, magnesium oxide; polysaccharides, e.g. cellulose, starch; fertilizers, e.g. ammonium sulfate, ammonium phosphate, ammonium nitrate, ureas; products of vegetable origin, e.g. cereal meal, tree bark meal, wood meal, nutshell meal, and mixtures thereof.

Suitable surfactants are surface-active compounds, such as anionic, cationic, nonionic and amphoteric surfactants, block polymers, polyelectrolytes, and mixtures thereof. Such surfactants can be used as emulsifier, dispersant, solubilizer, wetter, penetration enhancer, protective colloid, or adjuvant. Examples of surfactants are listed in McCutcheon's, Vol.1: Emulsifiers & Detergents, McCutcheon's Directories, Glen Rock, USA, 2008 (International Ed. or North American Ed.).

Suitable anionic surfactants are alkali, alkaline earth or ammonium salts of sulfonates, sulfates, phosphates, carboxylates, and mixtures thereof. Examples of sulfonates are alkylaryl-sulfonates, diphenylsulfonates, alpha-olefin sulfonates, lignine sulfonates, sulfonates of fatty acids and oils, sulfonates of ethoxylated alkylphenols, sulfonates of alkoxyated arylphenols, sulfonates of condensed naphthalenes, sulfonates of dodecyl- and tridecylbenzenes, sulfonates of naphthalenes and alkylnaphthalenes, sulfosuccinates or sulfosuccinamates. Examples of sulfates are sulfates of fatty acids and oils, of ethoxylated alkylphenols, of alcohols, of ethoxylated alcohols, or of fatty acid esters. Examples of phosphates are phosphate esters. Examples of carboxylates are alkyl carboxylates, and carboxylated alcohol or alkylphenol ethoxylates.

Suitable nonionic surfactants are alkoxyates, N-substituted fatty acid amides, amine oxides, esters, sugar-based surfactants, polymeric surfactants, and mixtures thereof. Examples of alkoxyates are compounds such as alcohols, alkylphenols, amines, amides, arylphenols, fatty acids or fatty acid esters which have been alkoxyated with 1 to 50 equivalents. Ethylene oxide

and/or propylene oxide may be employed for the alkoxylation, preferably ethylene oxide. Examples of N-substituted fatty acid amides are fatty acid glucamides or fatty acid alkanolamides. Examples of esters are fatty acid esters, glycerol esters or monoglycerides. Examples of sugar-based surfactants are sorbitans, ethoxylated sorbitans, sucrose and glucose esters or alkylpolyglucosides. Examples of polymeric surfactants are homo- or copolymers of vinylpyrrolidone, vinylalcohols, or vinylacetate.

Suitable cationic surfactants are quaternary surfactants, for example quaternary ammonium compounds with one or two hydrophobic groups, or salts of long-chain primary amines. Suitable amphoteric surfactants are alkylbetains and imidazolines. Suitable block polymers are block polymers of the A-B or A-B-A type comprising blocks of polyethylene oxide and polypropylene oxide, or of the A-B-C type comprising alkanol, polyethylene oxide and polypropylene oxide. Suitable polyelectrolytes are polyacids or polybases. Examples of polyacids are alkali salts of polyacrylic acid or polyacid comb polymers. Examples of polybases are polyvinylamines or polyethyleneamines.

Suitable adjuvants are compounds, which have a neglectable or even no pesticidal activity themselves, and which improve the biological performance of the inventive mixtures on the target. Examples are surfactants, mineral or vegetable oils, and other auxiliaries. Further examples are listed by Knowles, Adjuvants and additives, Agrow Reports DS256, T&F Informa UK, 2006, chapter 5.

Suitable thickeners are polysaccharides (e.g. xanthan gum, carboxymethylcellulose), anorganic clays (organically modified or unmodified), polycarboxylates, and silicates.

Suitable bactericides are bronopol and isothiazolinone derivatives such as alkylisothiazolinones and benzisothiazolinones.

Suitable anti-freezing agents are ethylene glycol, propylene glycol, urea and glycerin.

Suitable anti-foaming agents are silicones, long chain alcohols, and salts of fatty acids.

Suitable colorants (e.g. in red, blue, or green) are pigments of low water solubility and water-soluble dyes. Examples are inorganic colorants (e.g. iron oxide, titan oxide, iron hexacyanoferrate) and organic colorants (e.g. alizarin-, azo- and phthalocyanine colorants).

Suitable tackifiers or binders are polyvinylpyrrolidons, polyvinylacetates, polyvinyl alcohols, polyacrylates, biological or synthetic waxes, and cellulose ethers.

Examples for composition types and their preparation are:

i) Water-soluble concentrates (SL, LS)

10-60 wt% of an inventive mixture and 5-15 wt% wetting agent (e.g. alcohol alkoxyates) are dissolved in water and/or in a water-soluble solvent (e.g. alcohols) ad 100 wt%. The active substance dissolves upon dilution with water.

ii) Dispersible concentrates (DC)

5-25 wt% of an inventive mixture and 1-10 wt% dispersant (e. g. polyvinylpyrrolidone) are dissolved in organic solvent (e.g. cyclohexanone) ad 100 wt%. Dilution with water gives a dispersion.

iii) Emulsifiable concentrates (EC)

15-70 wt% of an inventive mixture and 5-10 wt% emulsifiers (e.g. calcium dodecylbenzenesulfonate and castor oil ethoxylate) are dissolved in water-insoluble organic solvent (e.g. aromatic hydrocarbon) ad 100 wt%. Dilution with water gives an emulsion.

iv) Emulsions (EW, EO, ES)

5-40 wt% of an inventive mixture and 1-10 wt% emulsifiers (e.g. calcium dodecylbenzene-sulfonate and castor oil ethoxylate) are dissolved in 20-40 wt% water-insoluble organic solvent (e.g. aromatic hydrocarbon). This mixture is introduced into water ad 100 wt% by means of an emulsifying machine and made into a homogeneous emulsion. Dilution with water gives an emulsion.

v) Suspensions (SC, OD, FS)

In an agitated ball mill, 20-60 wt% of an inventive mixture are comminuted with addition of 2-10 wt% dispersants and wetting agents (e.g. sodium lignosulfonate and alcohol ethoxylate), 0.1-2 wt% thickener (e.g. xanthan gum) and water ad 100 wt% to give a fine active substance suspension. Dilution with water gives a stable suspension of the active substance. For FS type composition up to 40 wt% binder (e.g. polyvinylalcohol) is added.

vi) Water-dispersible granules and water-soluble granules (WG, SG)

50-80 wt% of an inventive mixture are ground finely with addition of dispersants and wetting agents (e.g. sodium lignosulfonate and alcohol ethoxylate) ad 100 wt% and prepared as water-dispersible or water-soluble granules by means of technical appliances (e. g. extrusion, spray tower, fluidized bed). Dilution with water gives a stable dispersion or solution of the active substance.

vii) Water-dispersible powders and water-soluble powders (WP, SP, WS)

50-80 wt% of an inventive mixture are ground in a rotor-stator mill with addition of 1-5 wt% dispersants (e.g. sodium lignosulfonate), 1-3 wt% wetting agents (e.g. alcohol ethoxylate) and solid carrier (e.g. silica gel) ad 100 wt%. Dilution with water gives a stable dispersion or solution of the active substance.

viii) Gel (GW, GF)

In an agitated ball mill, 5-25 wt% of an inventive mixture are comminuted with addition of 3-10 wt% dispersants (e.g. sodium lignosulfonate), 1-5 wt% thickener (e.g. carboxymethylcellulose) and water ad 100 wt% to give a fine suspension of the active substance. Dilution with water gives a stable suspension of the active substance.

ix) Microemulsion (ME)

5-20 wt% of an inventive mixture are added to 5-30 wt% organic solvent blend (e.g. fatty acid dimethylamide and cyclohexanone), 10-25 wt% surfactant blend (e.g. alcohol ethoxylate and arylphenol ethoxylate), and water ad 100 %. This mixture is stirred for 1 h to produce spontaneously a thermodynamically stable microemulsion.

x) Microcapsules (CS)

An oil phase comprising 5-50 wt% of an inventive mixture, 0-40 wt% water insoluble organic solvent (e.g. aromatic hydrocarbon), 2-15 wt% acrylic monomers (e.g. methylmethacrylate, methacrylic acid and a di- or triacrylate) are dispersed into an aqueous solution of a protective colloid (e.g. polyvinyl alcohol). Radical polymerization initiated by a radical initiator results in the formation of poly(meth)acrylate microcapsules. Alternatively, an oil phase comprising 5-50 wt% of an inventive mixture according to the invention, 0-40 wt% water insoluble organic solvent (e.g. aromatic hydrocarbon), and an isocyanate monomer (e.g. diphenylmethene-4,4'-diisocyanate) are dispersed into an aqueous solution of a protective colloid (e.g. polyvinyl alcohol). The addition of a polyamine (e.g. hexamethylenediamine) results in the formation of poly-

yurea microcapsules. The monomers amount to 1-10 wt%. The wt% relate to the total CS composition.

ix) Dustable powders (DP, DS)

1-10 wt% of an inventive mixture are ground finely and mixed intimately with solid carrier (e.g. finely divided kaolin) ad 100 wt%.

x) Granules (GR, FG)

0.5-30 wt% of an inventive mixture is ground finely and associated with solid carrier (e.g. silicate) ad 100 wt%. Granulation is achieved by extrusion, spray-drying or fluidized bed.

xi) Ultra-low volume liquids (UL)

1-50 wt% of an inventive mixture are dissolved in organic solvent (e.g. aromatic hydrocarbon) ad 100 wt%.

The compositions types i) to xi) may optionally comprise further auxiliaries, such as 0.1-1 wt% bactericides, 5-15 wt% anti-freezing agents, 0.1-1 wt% anti-foaming agents, and 0.1-1 wt% colorants.

The resulting agrochemical compositions generally comprise between 0.01 and 95%, preferably between 0.1 and 90%, and in particular between 0.5 and 75%, by weight of active substance. The active substances are employed in a purity of from 90% to 100%, preferably from 95% to 100% (according to NMR spectrum).

Solutions for seed treatment (LS), Suspoemulsions (SE), flowable concentrates (FS), powders for dry treatment (DS), water-dispersible powders for slurry treatment (WS), water-soluble powders (SS), emulsions (ES), emulsifiable concentrates (EC) and gels (GF) are usually employed for the purposes of treatment of plant propagation materials, particularly seeds. The compositions in question give, after two-to-tenfold dilution, active substance concentrations of from 0.01 to 60% by weight, preferably from 0.1 to 40%, in the ready-to-use preparations. Application can be carried out before or during sowing. Methods for applying the inventive mixtures and compositions thereof, respectively, on to plant propagation material, especially seeds include dressing, coating, pelleting, dusting, soaking and in-furrow application methods of the propagation material. Preferably, the inventive mixtures or the compositions thereof, respectively, are applied on to the plant propagation material by a method such that germination is not induced, e. g. by seed dressing, pelleting, coating and dusting.

When employed in plant protection, the amounts of active substances applied are, depending on the kind of effect desired, from 0.001 to 2 kg per ha, preferably from 0.005 to 2 kg per ha, more preferably from 0.01 to 1.0 kg per ha, and in particular from 0.05 to 0.75 kg per ha.

In treatment of plant propagation materials such as seeds, e. g. by dusting, coating or drenching seed, amounts of active substance of from 0.01-10kg, preferably from 0.1-1000 g, more preferably from 1-100 g per 100 kilogram of plant propagation material (preferably seeds) are generally required.

When used in the protection of materials or stored products, the amount of active substance applied depends on the kind of application area and on the desired effect. Amounts customarily applied in the protection of materials are 0.001 g to 2 kg, preferably 0.005 g to 1 kg, of active substance per cubic meter of treated material.

Various types of oils, wetters, adjuvants, fertilizer, or micronutrients, and further pesticides (e.g. herbicides, insecticides, fungicides, growth regulators, safeners) may be added to the ac-

tive substances or the compositions comprising them as premix or, if appropriate not until immediately prior to use (tank mix). These agents can be admixed with the compositions according to the invention in a weight ratio of 1:100 to 100:1, preferably 1:10 to 10:1.

The user applies the composition according to the invention usually from a predosage device, a knapsack sprayer, a spray tank, a spray plane, or an irrigation system. Usually, the agrochemical composition is made up with water, buffer, and/or further auxiliaries to the desired application concentration and the ready-to-use spray liquor or the agrochemical composition according to the invention is thus obtained. Usually, 20 to 2000 liters, preferably 50 to 400 liters, of the ready-to-use spray liquor are applied per hectare of agricultural useful area.

According to one embodiment, individual components of the composition according to the invention such as parts of a kit or parts of a binary or ternary mixture may be mixed by the user himself in a spray tank or any other kind of vessel used for applications (e. g. seed treater drums, seed pelleting machinery, knapsack sprayer) and further auxiliaries may be added, if appropriate.

Consequently, one embodiment of the invention is a kit for preparing a usable pesticidal composition, the kit comprising a) a composition comprising component 1) as defined herein and at least one auxiliary; and b) a composition comprising component 2) as defined herein and at least one auxiliary; and optionally c) a composition comprising at least one auxiliary and optionally a further active component 3) as defined herein.

As said above, the inventive mixtures are employed by treating the pests or the plants, plant propagation materials (preferably seeds), materials or soil to be protected from fungal attack with a pesticidally effective amount of the active compounds. The application can be carried out both before and after the infection of the materials, plants or plant propagation materials (preferably seeds) by the pests.

Herein, the present invention comprises a method for controlling harmful fungi, wherein the pest, their habitat, breeding grounds, their locus or the plants to be protected against pest attack, the soil or plant propagation material (preferably seed) are treated with an pesticidally effective amount of a inventive mixture.

Advantageously, the inventive mixtures are suitable for controlling the following fungal plant diseases:

Albugo spp. (white rust) on ornamentals, vegetables (e. g. *A. candida*) and sunflowers (e. g. *A. tragopogonis*); *Alternaria* spp. (*Alternaria* leaf spot) on vegetables, rape (*A. brassicola* or *brassicae*), sugar beets (*A. tenuis*), fruits, rice, soybeans, potatoes (e. g. *A. solani* or *A. alternata*), tomatoes (e. g. *A. solani* or *A. alternata*) and wheat; *Aphanomyces* spp. on sugar beets and vegetables; *Ascochyta* spp. on cereals and vegetables, e. g. *A. tritici* (anthracnose) on wheat and *A. hordei* on barley; *Bipolaris* and *Drechslera* spp. (teleomorph: *Cochliobolus* spp.), e. g. Southern leaf blight (*D. maydis*) or Northern leaf blight (*B. zeicola*) on corn, e. g. spot blotch (*B. sorokiniana*) on cereals and e. g. *B. oryzae* on rice and turfs; *Blumeria* (formerly *Erysiphe*) *graminis* (powdery mildew) on cereals (e. g. on wheat or barley); *Botrytis cinerea* (teleomorph: *Botryotinia fuckeliana*: grey mold) on fruits and berries (e. g. strawberries), vegetables (e. g. lettuce, carrots, celery and cabbages), rape, flowers, vines, forestry plants and wheat; *Bremia*

lactucae (downy mildew) on lettuce; *Ceratocystis* (syn. *Ophiostoma*) spp. (rot or wilt) on broad-leaved trees and evergreens, e. g. *C. ulmi* (Dutch elm disease) on elms; *Cercospora* spp. (*Cercospora* leaf spots) on corn (e. g. Gray leaf spot: *C. zea-maydis*), rice, sugar beets (e. g. *C. beticola*), sugar cane, vegetables, coffee, soybeans (e. g. *C. sojae* or *C. kikuchii*) and rice;

5 *Cladosporium* spp. on tomatoes (e. g. *C. fulvum*: leaf mold) and cereals, e. g. *C. herbarum* (black ear) on wheat; *Claviceps purpurea* (ergot) on cereals; *Cochliobolus* (anamorph: *Helminthosporium* of *Bipolaris*) spp. (leaf spots) on corn (*C. carbonum*), cereals (e. g. *C. sativus*, anamorph: *B. sorokiniana*) and rice (e. g. *C. miyabeanus*, anamorph: *H. oryzae*); *Colletotrichum* (teleomorph: *Glomerella*) spp. (anthracnose) on cotton (e. g. *C. gossypii*), corn (e. g. *C. graminicola*: Anthracnose stalk rot), soft fruits, potatoes (e. g. *C. coccodes*: black dot), beans (e. g. *C. lindemuthianum*) and soybeans (e. g. *C. truncatum* or *C. gloeosporioides*); *Corticium* spp., e. g. *C. sasakii* (sheath blight) on rice; *Corynespora cassiicola* (leaf spots) on soybeans and ornamentals; *Cycloconium* spp., e. g. *C. oleaginum* on olive trees; *Cylindrocarpon* spp. (e. g. fruit tree canker or young vine decline, teleomorph: *Nectria* or *Neonectria* spp.) on fruit trees, vines

15 (e. g. *C. liriodendri*, teleomorph: *Neonectria liriodendri*. Black Foot Disease) and ornamentals; *Dematophora* (teleomorph: *Rosellinia*) *necatrix* (root and stem rot) on soybeans; *Diaporthe* spp., e. g. *D. phaseolorum* (damping off) on soybeans; *Drechslera* (syn. *Helminthosporium*, teleomorph: *Pyrenophora*) spp. on corn, cereals, such as barley (e. g. *D. teres*, net blotch) and wheat (e. g. *D. tritici-repentis*: tan spot), rice and turf; Esca (dieback, apoplexy) on vines, caused by

20 *Formitiporia* (syn. *Phellinus*) *punctata*, *F. mediterranea*, *Phaeoconiella chlamydospora* (earlier *Phaeoacremonium chlamydosporum*), *Phaeoacremonium aleophilum* and/or *Botryosphaeria obtusa*; *Elsinoe* spp. on pome fruits (*E. pyri*), soft fruits (*E. veneta*: anthracnose) and vines (*E. ampelina*: anthracnose); *Entyloma oryzae* (leaf smut) on rice; *Epicoccum* spp. (black mold) on wheat; *Erysiphe* spp. (powdery mildew) on sugar beets (*E. betae*), vegetables (e. g. *E. pisi*),

25 such as cucurbits (e. g. *E. cichoracearum*), cabbages, rape (e. g. *E. cruciferarum*); *Eutypa lata* (*Eutypa* canker or dieback, anamorph: *Cytosporina lata*, syn. *Libertella blepharis*) on fruit trees, vines and ornamental woods; *Exserohilum* (syn. *Helminthosporium*) spp. on corn (e. g. *E. turcicum*); *Fusarium* (teleomorph: *Gibberella*) spp. (wilt, root or stem rot) on various plants, such as *F. graminearum* or *F. culmorum* (root rot, scab or head blight) on cereals (e. g. wheat or barley),

30 *F. oxysporum* on tomatoes, *F. solani* (f. sp. *glycines* now syn. *F. virguliforme*) and *F. tucumaniae* and *F. brasiliense* each causing sudden death syndrome on soybeans, and *F. verticillioides* on corn; *Gaeumannomyces graminis* (take-all) on cereals (e. g. wheat or barley) and corn; *Gibberella* spp. on cereals (e. g. *G. zea*) and rice (e. g. *G. fujikuroi*: Bakanae disease); *Glomerella cingulata* on vines, pome fruits and other plants and *G. gossypii* on cotton; Grainstaining complex on rice; *Guignardia bidwellii* (black rot) on vines; *Gymnosporangium* spp. on rosaceous plants and junipers, e. g. *G. sabinae* (rust) on pears; *Helminthosporium* spp. (syn. *Drechslera*, teleomorph: *Cochliobolus*) on corn, cereals and rice; *Hemileia* spp., e. g. *H. vastatrix* (coffee leaf rust) on coffee; *Isariopsis clavispora* (syn. *Cladosporium vitis*) on vines; *Macrophomina phaseolina* (syn. *phaseoli*) (root and stem rot) on soybeans and cotton; *Microdochium* (syn.

40 *Fusarium*) *nivale* (pink snow mold) on cereals (e. g. wheat or barley); *Microsphaera diffusa* (powdery mildew) on soybeans; *Monilinia* spp., e. g. *M.A. laxa*, *M.A. fructicola* and *M.A. fructigena* (bloom and twig blight, brown rot) on stone fruits and other rosaceous plants; *Mycosphaerella* spp. on cereals, bananas, soft fruits and ground nuts, such as e. g. *M.A. graminicola* (anamorph: *Septoria tritici*, *Septoria* blotch) on wheat or *M.A. fijiensis* (black Sigatoka disease)

on bananas; *Peronospora* spp. (downy mildew) on cabbage (e. g. *P. brassicae*), rape (e. g. *P. parasitica*), onions (e. g. *P. destructor*), tobacco (*P. tabacina*) and soybeans (e. g. *P. manshurica*); *Phakopsora pachyrhizi* and *P. meibomia* (soybean rust) on soybeans; *Phialophora* spp. e. g. on vines (e. g. *P. tracheiphila* and *P. tetraspora*) and soybeans (e. g. *P. gregata*: stem rot);

5 *Phoma lingam* (root and stem rot) on rape and cabbage and *P. betae* (root rot, leaf spot and damping-off) on sugar beets; *Phomopsis* spp. on sunflowers, vines (e. g. *P. viticola*: can and leaf spot) and soybeans (e. g. stem rot: *P. phaseoli*, teleomorph: *Diaporthe phaseolorum*); *Phy-*

10 *soderma maydis* (brown spots) on corn; *Phytophthora* spp. (wilt, root, leaf, fruit and stem root) on various plants, such as paprika and cucurbits (e. g. *P. capsici*), soybeans (e. g. *P. megasperma*, syn. *P. sojae*), potatoes and tomatoes (e. g. *P. infestans*: late blight) and broad-

15 leaved trees (e. g. *P. ramorum*: sudden oak death); *Plasmodiophora brassicae* (club root) on cabbage, rape, radish and other plants; *Plasmopara* spp., e. g. *P. viticola* (grapevine downy mildew) on vines and *P. halstedii* on sunflowers; *Podosphaera* spp. (powdery mildew) on rosa-

20 ceous plants, hop, pome and soft fruits, e. g. *P. leucotricha* on apples; *Polymyxa* spp., e. g. on cereals, such as barley and wheat (*P. graminis*) and sugar beets (*P. betae*) and thereby trans-

25 mitted viral diseases; *Pseudocercospora herpotrichoides* (eyespot, teleomorph: *Tapesia yal-lundae*) on cereals, e. g. wheat or barley; *Pseudoperonospora* (downy mildew) on various plants, e. g. *P. cubensis* on cucurbits or *P. humili* on hop; *Pseudopezizcula tracheiphila* (red fire disease or 'rotbrenner', anamorph: *Phialophora*) on vines; *Puccinia* spp. (rusts) on various

30 plants, e. g. *P. triticea* (brown or leaf rust), *P. striiformis* (stripe or yellow rust), *P. hordei* (dwarf rust), *P. graminis* (stem or black rust) or *P. recondita* (brown or leaf rust) on cereals, such as e. g. wheat, barley or rye, *P. kuehnii* (orange rust) on sugar cane and *P. asparagi* on asparagus; *Pyrenophora* (anamorph: *Drechslera*) *tritici-repentis* (tan spot) on wheat or *P. teres* (net blotch)

35 on barley; *Pyricularia* spp., e. g. *P. oryzae* (teleomorph: *Magnaporthe grisea*, rice blast) on rice and *P. grisea* on turf and cereals; *Pythium* spp. (damping-off) on turf, rice, corn, wheat, cotton, rape, sunflowers, soybeans, sugar beets, vegetables and various other plants (e. g. *P. ultimum*

40 or *P. aphanidermatum*); *Ramularia* spp., e. g. *R. collo-cygni* (Ramularia leaf spots, Physiological leaf spots) on barley and *R. beticola* on sugar beets; *Rhizoctonia* spp. on cotton, rice, potatoes, turf, corn, rape, potatoes, sugar beets, vegetables and various other plants, e. g. *R. solani* (root and stem rot) on soybeans, *R. solani* (sheath blight) on rice or *R. cerealis* (Rhizoctonia spring blight) on wheat or barley; *Rhizopus stolonifer* (black mold, soft rot) on strawberries, carrots, cabbage, vines and tomatoes; *Rhynchosporium secalis* (scald) on barley, rye and triticale; *Sa-*

45 *rocladium oryzae* and *S. attenuatum* (sheath rot) on rice; *Sclerotinia* spp. (stem rot or white mold) on vegetables and field crops, such as rape, sunflowers (e. g. *S. sclerotiorum*) and soy-

50 beans (e. g. *S. rolfsii* or *S. sclerotiorum*); *Septoria* spp. on various plants, e. g. *S. glycines* (brown spot) on soybeans, *S. tritici* (Septoria blotch) on wheat and *S.* (syn. *Stagonospora*) *nodorum* (Stagonospora blotch) on cereals; *Uncinula* (syn. *Erysiphe*) *necator* (powdery mildew, anamorph: *Oidium tuckeri*) on vines; *Setosphaeria* spp. (leaf blight) on corn (e. g. *S. turcicum*, syn. *Helminthosporium turcicum*) and turf; *Sphacelotheca* spp. (smut) on corn, (e. g. *S. reiliana*:

55 head smut), sorghum und sugar cane; *Sphaerotheca fuliginea* (powdery mildew) on cucurbits; *Spongospora subterranea* (powdery scab) on potatoes and thereby transmitted viral diseases; *Stagonospora* spp. on cereals, e. g. *S. nodorum* (Stagonospora blotch, teleomorph: *Lepto-*

60 *sphaeria* [syn. *Phaeosphaeria*] *nodorum*) on wheat; *Synchytrium endobioticum* on potatoes (potato wart disease); *Taphrina* spp., e. g. *T. deformans* (leaf curl disease) on peaches and *T. pruni*

(plum pocket) on plums; *Thielaviopsis* spp. (black root rot) on tobacco, pome fruits, vegetables, soybeans and cotton, e. g. *T. basicola* (syn. *Chalara elegans*); *Tilletia* spp. (common bunt or stinking smut) on cereals, such as e. g. *T. tritici* (syn. *T. caries*, wheat bunt) and *T. controversa* (dwarf bunt) on wheat; *Typhula incarnata* (grey snow mold) on barley or wheat; *Urocystis* spp.,
5 e. g. *U. occulta* (stem smut) on rye; *Uromyces* spp. (rust) on vegetables, such as beans (e. g. *U. appendiculatus*, syn. *U. phaseoli*) and sugar beets (e. g. *U. betae*); *Ustilago* spp. (loose smut) on cereals (e. g. *U. nuda* and *U. avenae*), corn (e. g. *U. maydis*: corn smut) and sugar cane; *Venturia* spp. (scab) on apples (e. g. *V. inaequalis*) and pears; and *Verticillium* spp. (wilt) on various plants, such as fruits and ornamentals, vines, soft fruits, vegetables and field crops,
10 e. g. *V. dahliae* on strawberries, rape, potatoes and tomatoes.

The mixtures according to the present invention, respectively, are also suitable for controlling harmful fungi in the protection of stored products or harvest and in the protection of materials.

The term "protection of materials" is to be understood to denote the protection of technical and non-living materials, such as adhesives, glues, wood, paper and paperboard, textiles,
15 leather, paint dispersions, plastics, cooling lubricants, fiber or fabrics, against the infestation and destruction by harmful microorganisms, such as fungi and bacteria. As to the protection of wood and other materials, the particular attention is paid to the following harmful fungi: Ascomycetes such as *Ophiostoma* spp., *Ceratocystis* spp., *Aureobasidium pullulans*, *Sclerophoma* spp., *Chaetomium* spp., *Humicola* spp., *Petriella* spp., *Trichurus* spp.; Basidiomycetes such as *Coni-*
20 *ophora* spp., *Coriolus* spp., *Gloeophyllum* spp., *Lentinus* spp., *Pleurotus* spp., *Poria* spp., *Serpula* spp. and *Tyromyces* spp., Deuteromycetes such as *Aspergillus* spp., *Cladosporium* spp., *Penicillium* spp., *Trichoderma* spp., *Alternaria* spp., *Paecilomyces* spp. and Zygomycetes such as *Mucor* spp., and in addition in the protection of stored products and harvest the following yeast fungi are worthy of note: *Candida* spp. and *Saccharomyces cerevisiae*.

25 They are particularly important for controlling a multitude of fungi on various cultivated plants, such as bananas, cotton, vegetable species (for example cucumbers, beans and cucurbits), cereals such as wheat, rye, barley, rice, oats; grass coffee, potatoes, corn, fruit species, soya, tomatoes, grapevines, ornamental plants,
30 sugar cane and also on a large number of seeds.

Furthermore, as outlined above, the present invention comprises a method for controlling harmful animal pest, wherein the pest, their habitat, breeding grounds, their locus or the plants to be protected against pest attack, the soil or plant propagation material (preferably seed) are treated
35 with an pesticidally effective amount of a inventive mixture.

Herein, the mixtures according to the present publication exhibit also outstanding action against animal pests from the following orders:

40 insects from the order of **Lepidoptera**, for example *Achroia grisella*, *Acleris* spp. such as *A. fimbriana*, *A. gloverana*, *A. variana*; *Acrolepiopsis assectella*, *Acronicta major*, *Adoxophyes* spp. such as *A. cyrtosema*, *A. orana*; *Aedia leucomelas*, *Agrotis* spp. such as *A. exclamationis*, *A. fucosa*, *A. ipsilon*, *A. orthogoma*, *A. segetum*, *A. subterranea*; *Alabama argillacea*, *Aleurodicus dispersus*, *Alsophila pometaria*, *Ampelophaga rubiginosa*, *Amyelois transitella*, *Anacampsis*

sarcitella, *Anagasta kuehniella*, *Anarsia lineatella*, *Anisota senatoria*, *Antheraea pernyi*, *Anticarsia* (= *Thermesia*) spp. such as *A. gemmatalis*; *Apamea* spp., *Aproaerema modicella*, *Archips* spp. such as *A. argyrospila*, *A. fuscocupreanus*, *A. rosana*, *A. xyloseanus*; *Argyresthia conjugella*, *Argyroproce* spp., *Argyrotaenia* spp. such as *A. velutinana*; *Athetis mindara*, *Austroasca viridigrisea*, *Autographa gamma*, *Autographa nigrisigna*, *Barathra brassicae*, *Bedellia* spp., *Bonagota salubricola*, *Borbo cinnara*, *Bucculatrix thurberiella*, *Bupalus piniarius*, *Busseola* spp., *Cacoecia* spp. such as *C. murinana*, *C. podana*; *Cactoblastis cactorum*, *Cadra cautella*, *Calingo braziliensis*, *Caloptilis theivora*, *Capua reticulana*, *Carposina* spp. such as *C. niponensis*, *C. sasakii*; *Cephus* spp., *Chaetocnema aridula*, *Cheimatobia brumata*, *Chilo* spp. such as *C. Indicus*, *C. suppressalis*, *C. partellus*; *Choreutis pariana*, *Choristoneura* spp. such as *C. conflictana*, *C. fumiferana*, *C. longicellana*, *C. murinana*, *C. occidentalis*, *C. rosaceana*; *Chrysodeixis* (= *Pseudoplusia*) spp. such as *C. eriosoma*, *C. includens*; *Cirphis unipuncta*, *Clysia ambiguella*, *Cnaphalocerus* spp., *Cnaphalocrocis medinalis*, *Cnephasia* spp., *Cochylis hospes*, *Coleophora* spp., *Colias eurytheme*, *Conopomorpha* spp., *Conotrachelus* spp., *Copitarsia* spp., *Corcyra cephalonica*, *Crambus caliginosellus*, *Crambus teterrellus*, *Crociosema* (= *Epinotia*) *aporema*, *Cydalima* (= *Diaphania*) *perspectalis*, *Cydia* (= *Carpocapsa*) spp. such as *C. pomonella*, *C. latiferreana*; *Dalaca noctuides*, *Datana integerrima*, *Dasychira pinicola*, *Dendrolimus* spp. such as *D. pini*, *D. spectabilis*, *D. sibiricus*; *Desmia funeralis*, *Diaphania* spp. such as *D. nitidalis*, *D. hyalinata*; *Diatraea grandiosella*, *Diatraea saccharalis*, *Diphthera festiva*, *Earias* spp. such as *E. insulana*, *E. vittella*; *Ecdytolopa aurantianu*, *Egira* (= *Xylomyges*) *curialis*, *Elasmopalpus lignosellus*, *Eldana saccharina*, *Endopiza viteana*, *Ennomos subsignaria*, *Eoreuma loftini*, *Ephestia* spp. such as *E. cautella*, *E. elutella*, *E. kuehniella*; *Epinotia aporema*, *Epiphyas postvittana*, *Erannis tiliaria*, *Erionota thrax*, *Etiella* spp., *Eulia* spp., *Eupoecilia ambiguella*, *Euproctis chryssorrhoea*, *Euxoa* spp., *Evetria bouliana*, *Faronta albilinea*, *Feltia* spp. such as *F. subterranean*; *Galleria mellonella*, *Gracillaria* spp., *Grapholita* spp. such as *G. funebrana*, *G. molesta*, *G. inopinata*; *Halysidota* spp., *Harrisina americana*, *Hedylepta* spp., *Helicoverpa* spp. such as *H. armigera* (= *Heliiothis armigera*), *H. zea* (= *Heliiothis zea*); *Heliiothis* spp. such as *H. assulta*, *H. subflexa*, *H. virescens*; *Hellula* spp. such as *H. undalis*, *H. rogatalis*; *Helocoverpa gelotopoeon*, *Hemileuca oliviae*, *Herpetogramma licarsisalis*, *Hibernia defoliaria*, *Hofmannophila pseudopretella*, *Homoeosoma electellum*, *Homona magnanima*, *Hypena scabra*, *Hyphantria cunea*, *Hyponomeuta padella*, *Hyponomeuta malinellus*, *Kakivoria flavofasciata*, *Keiferia lycopersicella*, *Lambdina fiscellaria fiscellaria*, *Lambdina fiscellaria lugubrosa*, *Lamprosema indicata*, *Laspeyresia molesta*, *Leguminivora glycinivorella*, *Lerodea eufala*, *Leucinodes orbonalis*, *Leucoma salicis*, *Leucoptera* spp. such as *L. coffeella*, *L. scitella*; *Leuminivora lycinivorella*, *Lithocolletis blancardella*, *Lithophane antennata*, *Llattia octo* (= *Amyna axis*), *Lobesia botrana*, *Lophocampa* spp., *Loxagrotis albicosta*, *Loxostege* spp. such as *L. sticticalis*, *L. cereralis*; *Lymantria* spp. such as *L. dispar*, *L. monacha*; *Lyonetia clerkella*, *Lyonetia prunifoliella*, *Malacosoma* spp. such as *M. americanum*, *M. californicum*, *M. constrictum*, *M. neustria*; *Mamestra* spp. such as *M. brassicae*, *M. configurata*; *Mamstra brassicae*, *Manduca* spp. such as *M. quinquemaculata*, *M. sexta*; *Marasmia* spp., *Marmara* spp., *Maruca testulalis*, *Megalopyge lanata*, *Melanchra picta*, *Melanitis leda*, *Mocis* spp. such as *M. lapites*, *M. repanda*; *Mocis latipes*, *Monochroa fragariae*, *Mythimna separata*, *Nemapogon cloacella*, *Neoleucinodes elegantalis*, *Nepytia* spp., *Nymphula* spp., *Oiketicus* spp., *Omiodes indicata*, *Omphisa anastomosalis*, *Operophtera brumata*, *Orgyia pseudotsugata*, *Oria* spp., *Orthaga thyrisalis*, *Ostrinia* spp. such as *O.*

nubilalis; *Oulema oryzae*, *Paleacrita vernata*, *Panolis flammea*, *Parnara* spp., *Papaipema nebris*, *Papilio cresphontes*, *Paramyelois transitella*, *Paranthrene regalis*, *Paysandisia archon*, *Pectinophora* spp. such as *P. gossypiella*; *Peridroma saucia*, *Perileucoptera* spp., such as *P. coffeella*; *Phalera bucephala*, *Phryganidia californica*, *Phthorimaea* spp. such as *P. operculella*;
5 *Phyllocnistis citrella*, *Phyllonorycter* spp. such as *P. blancardella*, *P. crataegella*, *P. issikii*, *P. ringoniella*; *Pieris* spp. such as *P. brassicae*, *P. rapae*, *P. napi*; *Pilocrocis tripunctata*, *Plathypena scabra*, *Platynota* spp. such as *P. flavedana*, *P. idaeusalis*, *P. stultana*; *Platyptilia carduidactyla*, *Plebejus argus*, *Plodia interpunctella*, *Plusia* spp, *Plutella maculipennis*, *Plutella xylostella*, *Pontia protodica*, *Prays* spp., *Prodenia* spp., *Proxenus lepigone*, *Pseudaletia* spp. such
10 as *P. sequax*, *P. unipuncta*; *Pyrausta nubilalis*, *Rachiplusia nu*, *Richia albicosta*, *Rhizobius ventralis*, *Rhyacionia frustrana*, *Sabulodes aegrotata*, *Schizura concinna*, *Schoenobius* spp., *Schreckensteinia festaliella*, *Scirpophaga* spp. such as *S. incertulas*, *S. innotata*; *Scotia segetum*, *Sesamia* spp. such as *S. inferens*, *Seudyra subflava*, *Sitotroga cerealella*, *Sparganthis pilleriana*, *Spilonota lechriaspis*, *S. ocellana*, *Spodoptera (=Lamphygma)* spp. such as *S.*
15 *cosmoides*, *S. eridania*, *S. exigua*, *S. frugiperda*, *S. latifascia*, *S. littoralis*, *S. litura*, *S. omithogalli*; *Stigmella* spp., *Stomopteryx subsecivella*, *Strymon bazochii*, *Sylepta derogata*, *Synanthedon* spp. such as *S. exitiosa*, *Tecia solanivora*, *Telehin licus*, *Thaumatopoea pityocampa*, *Thaumatotibia (=Cryptophlebia) leucotreta*, *Thaumatopoea pityocampa*, *Thecla* spp., *Theresimima ampelophaga*, *Thyrintina* spp, *Tildenia inconspicua*, *Tinea* spp. such as *T. cloacella*,
20 *T. pellionella*; *Tineola bisselliella*, *Tortrix* spp. such as *T. viridana*; *Trichophaga tapetzella*, *Trichoplusia* spp. such as *T. ni*; *Tuta (=Scrobipalpula) absoluta*, *Udea* spp. such as *U. rubigalis*, *U. rubigalis*; *Virachola* spp., *Yponomeuta padella*, and *Zeiraphera canadensis*;
insects from the order of **Coleoptera**, for example *Acalymma vittatum*, *Acanthoscehdes obtectus*, *Adoretus* spp., *Agelastica alni*, *Agrilus* spp. such as *A. anxius*, *A. planipennis*, *A. sinuatus*;
25 *Agriotes* spp. such as *A. fuscicollis*, *A. lineatus*, *A. obscurus*; *Alphitobius diaperinus*, *Amphimallus solstitialis*, *Anisandrus dispar*, *Anisoplia austriaca*, *Anobium punctatum*, *Anomala corpulenta*, *Anomala rufocuprea*, *Anoplophora* spp. such as *A. glabripennis*; *Anthonomus* spp. such as *A. eugenii*, *A. grandis*, *A. pomorum*; *Anthrenus* spp., *Aphthona euphoridae*, *Apion* spp., *Apogonia* spp., *Athous haemorrhoidalis*, *Atomaria* spp. such as *A. linearis*; *Attagenus* spp., *Aulacophora femoralis*, *Blastophagus piniperda*, *Blitophaga undata*, *Bruchidius obtectus*, *Bruchus*
30 spp. such as *B. lentis*, *B. pisorum*, *B. rufimanus*; *Byctiscus betulae*, *Callidiellum rufipenne*, *Callopietria floridensis*, *Callosobruchus chinensis*, *Cameraria ohridella*, *Cassida nebulosa*, *Cerotoma trifurcata*, *Cetonia aurata*, *Ceuthorhynchus* spp. such as *C. assimilis*, *C. napi*; *Chaetocnema tibialis*, *Cleonus mendicus*, *Conoderus* spp. such as *C. vespertinus*; *Conotrachelus nenuphar*,
35 *Cosmopolites* spp., *Costelytra zealandica*, *Crioceris asparagi*, *Cryptolestes ferrugineus*, *Cryptorhynchus lapathi*, *Ctenicera* spp. such as *C. destructor*; *Curculio* spp., *Cylindrocopturus* spp., *Cyclocephala* spp., *Dactylispa balyi*, *Dectes texanus*, *Dermestes* spp., *Diabrotica* spp. such as *D. undecimpunctata*, *D. speciosa*, *D. longicornis*, *D. semipunctata*, *D. virgifera*; *Diaprepes abbreviatus*, *Dichocrocis* spp., *Di cladispa armigera*, *Diloboderus abderus*, *Diocalandra frumenti*
40 (*Diocalandra stigmaticollis*), *Enaphalodes rufulus*, *Epilachna* spp. such as *E. varivestis*, *E. vigintioctomaculata*; *Epitrix* spp. such as *E. hirtipennis*, *E. similaris*; *Eutheola humilis*, *Eutinobothrus brasiliensis*, *Faustinus cubae*, *Gibbium psylloides*, *Gnathocerus cornutus*, *Hellula undalis*, *Heteronychus arator*, *Hylamorpha elegans*, *Hylobius abietis*, *Hylotrupes bajulus*, *Hypera* spp. such as *H. brunneipennis*, *H. postica*; *Hypomeces squamosus*, *Hypothenemus* spp.,

Ips typographus, *Lachnosterna consanguinea*, *Lasioderma serricorne*, *Latheticus oryzae*, *Lathridius* spp., *Lema* spp. such as *L. bilineata*, *L. melanopus*; *Leptinotarsa* spp. such as *L. decemlineata*; *Leptispa pygmaea*, *Limonius californicus*, *Lissorhoptrus oryzophilus*, *Lixus* spp., *Luperodes* spp., *Lyctus* spp. such as *L. bruneus*; *Liogenys fuscus*, *Macroductylus* spp. such as *M. subspinosus*; *Maladera matrida*, *Megaplatypus mutates*, *Megascelis* spp., *Melanotus communis*, *Meligethes* spp. such as *M. aeneus*; *Melolontha* spp. such as *M. hippocastani*, *M. melolontha*; *Metamasius hemipterus*, *Microtheca* spp., *Migdolus* spp. such as *M. fryanus*, *Monochamus* spp. such as *M. alternatus*; *Naupactus xanthographus*, *Niptus hololeucus*, *Oberia brevis*, *Oemona hirta*, *Oryctes rhinoceros*, *Oryzaephilus surinamensis*, *Oryzaphagus oryzae*, *Otiorrhynchus sulcatus*, *Otiorrhynchus ovatus*, *Otiorrhynchus sulcatus*, *Oulema melanopus*, *Oulema oryzae*, *Oxycetonia jucunda*, *Phaedon* spp. such as *P. brassicae*, *P. cochleariae*; *Phoracantha recurva*, *Phyllobius pyri*, *Phyllopertha horticola*, *Phyllophaga* spp. such as *P. helleri*; *Phyllotreta* spp. such as *P. chrysocephala*, *P. nemorum*, *P. striolata*, *P. vittula*; *Phyllopertha horticola*, *Popillia japonica*, *Premnotrypes* spp., *Psacotha hilaris*, *Psylliodes chrysocephala*, *Prostephanus truncates*, *Psylliodes* spp., *Ptinus* spp., *Pulga saltona*, *Rhizopertha dominica*, *Rhynchophorus* spp. such as *R. billineatus*, *R. ferrugineus*, *R. palmarum*, *R. phoenicis*, *R. vulneratus*; *Saperda candida*, *Scolytus schevyrewi*, *Scyphophorus acupunctatus*, *Sitona lineatus*, *Sitophilus* spp. such as *S. granaria*, *S. oryzae*, *S. zeamais*; *Sphenophorus* spp. such as *S. levis*; *Stegobium paniceum*, *Sternechus* spp. such as *S. subsignatus*; *Strophomorphus ctenotus*, *Symphyletes* spp., *Tanymecus* spp., *Tenebrio molitor*, *Tenebrioides mauretanicus*, *Tribolium* spp. such as *T. castaneum*; *Trogoderma* spp., *Tychius* spp., *Xylotrechus* spp. such as *X. pyrrhoderus*; and, *Zabrus* spp. such as *Z. tenebrioides*;

insects from the order of **Diptera** for example *Aedes* spp. such as *A. aegypti*, *A. albopictus*, *A. vexans*; *Anastrepha ludens*, *Anopheles* spp. such as *A. albimanus*, *A. crucians*, *A. freeborni*, *A. gambiae*, *A. leucosphyrus*, *A. maculipennis*, *A. minimus*, *A. quadrimaculatus*, *A. sinensis*; *Bactrocera invadens*, *Bibio hortulanus*, *Calliphora erythrocephala*, *Calliphora vicina*, *Ceratitis capitata*, *Chrysomyia* spp. such as *C. bezziana*, *C. hominivorax*, *C. macellaria*; *Chrysops atlanticus*, *Chrysops discalis*, *Chrysops silacea*, *Cochliomyia* spp. such as *C. hominivorax*; *Contarinia* spp. such as *C. sorghicola*; *Cordylobia anthropophaga*, *Culex* spp. such as *C. nigripalpus*, *C. pipiens*, *C. quinquefasciatus*, *C. tarsalis*, *C. tritaeniorhynchus*; *Culicoides furens*, *Culiseta inornata*, *Culiseta melanura*, *Cuterebra* spp., *Dacus cucurbitae*, *Dacus oleae*, *Dasineura brassicae*, *Dasineura oxycoccana*, *Delia* spp. such as *D. antique*, *D. coarctata*, *D. platura*, *D. radicum*; *Dermatobia hominis*, *Drosophila* spp. such as *D. sukukii*, *Fannia* spp. such as *F. canicularis*; *Gastrophilus* spp. such as *G. intestinalis*; *Geomyza tipunctata*, *Glossina* spp. such as *G. fuscipes*, *G. morsitans*, *G. palpalis*, *G. tachinoides*; *Haematobia irritans*, *Haplodiplosis equestris*, *Hippelates* spp., *Hylemyia* spp. such as *H. platura*; *Hypoderma* spp. such as *H. lineata*; *Hyppobosca* spp., *Hydrellia philippina*, *Leptoconops torrens*, *Liriomyza* spp. such as *L. sativae*, *L. trifolii*; *Lucilia* spp. such as *L. caprina*, *L. cuprina*, *L. sericata*; *Lycoria pectoralis*, *Mansonia titillanus*, *Mayetiola* spp. such as *M. destructor*; *Musca* spp. such as *M. autumnalis*, *M. domestica*; *Muscina stabulans*, *Oestrus* spp. such as *O. ovis*; *Opomyza florum*, *Oscinella* spp. such as *O. frit*; *Orseolia oryzae*, *Pegomya hysocyami*, *Phlebotomus argentipes*, *Phorbia* spp. such as *P. antiqua*, *P. brassicae*, *P. coarctata*; *Phytomyza gymnostoma*, *Prosimulium mixtum*, *Psila rosae*, *Psorophora columbiae*, *Psorophora discolor*, *Rhagoletis* spp. such as *R. cerasi*, *R. cingulate*, *R. indifferens*, *R. mendax*, *R. pomonella*; *Rivellia quadrifasciata*, *Sarcophaga* spp. such as *S.*

haemorrhoidalis; *Simulium vittatum*, *Sitodiplosis mosellana*, *Stomoxys* spp. such as *S. calcitrans*; *Tabanus* spp. such as *T. atratus*, *T. bovinus*, *T. lineola*, *T. similis*; *Tannia* spp., *Thecodiplosis japonensis*, *Tipula oleracea*, *Tipula paludosa*, and *Wohlfahrtia* spp;

insects from the order of **Thysanoptera** for example, *Baliothrips biformis*, *Dichromothrips corbetti*, *Dichromothrips* ssp., *Echinothrips americanus*, *Enneothrips flavens*, *Frankliniella* spp.

5 such as *F. fusca*, *F. occidentalis*, *F. tritici*; *Heliothrips* spp., *Hercinothrips femoralis*, *Kakothrips* spp., *Microcephalothrips abdominalis*, *Neohydatothrips samayunkur*, *Pezothrips kellyanus*, *Rhipiphorotherips cruentatus*, *Scirtothrips* spp. such as *S. citri*, *S. dorsalis*, *S. perseae*; *Stenchaetothrips* spp, *Taeniothrips cardamoni*, *Taeniothrips inconsequens*, *Thrips* spp. such as *T. imagines*, *T. hawaiiensis*, *T. oryzae*, *T. palmi*, *T. parvispinus*, *T. tabaci*;

10 insects from the order of **Hemiptera** for example, *Acizzia jamatonica*, *Acrosternum* spp. such as *A. hilare*; *Acyrtosipon* spp. such as *A. onobrychis*, *A. pisum*; *Adelges laricis*, *Adelges tsugae*, *Adelphocoris* spp., such as *A. rapidus*, *A. superbus*; *Aeneolamia* spp., *Agonoscena* spp., *Aulacorthum solani*, *Aleurocanthus woglumi*, *Aleurodes* spp., *Aleurodicus disperses*, *Aleurolobus barodensis*, *Aleurothrixus* spp., *Amrasca* spp., *Anasa tristis*, *Antestiopsis* spp., *Anuraphis cardui*, *Aonidiella* spp., *Aphanostigma piri*, *Aphidula nasturtii*, *Aphis* spp. such as *A. craccivora*, *A. fabae*, *A. forbesi*, *A. gossypii*, *A. grossulariae*, *A. maidiradicis*, *A. pomi*, *A. sambuci*, *A. schneideri*, *A. spiraeicola*; *Arboridia apicalis*, *Arilus critatus*, *Aspidiella* spp., *Aspidiotus* spp., *Atanus* spp., *Aulacaspis yasumatsui*, *Aulacorthum solani*, *Bactericera cockerelli* (*Paratrioza cockerelli*),

20 *Bemisia* spp. such as *B. argentifolii*, *B. tabaci* (*Aleurodes tabaci*); *Blissus* spp. such as *B. leucopterus*; *Brachycaudus* spp. such as *B. cardui*, *B. helichrysi*, *B. persicae*, *B. prunicola*; *Brachycolus* spp., *Brachycorynella asparagi*, *Brevicoryne brassicae*, *Cacopsylla* spp. such as *C. fulguralis*, *C. pyricola* (*Psylla piri*); *Calligypona marginata*, *Calocoris* spp., *Campylomma livida*, *Capitophorus horni*, *Carneocephala fulgida*, *Cavelerius* spp., *Ceraplastes* spp., *Ceratovacuna lanigera*, *Ceroplastes ceriferus*, *Cerosiphia gossypii*, *Chaetosiphon fragaefolii*, *Chionaspis tegalensis*, *Chlorita onukii*, *Chromaphis juglandicola*, *Chrysomphalus ficus*, *Cicadulina mbila*, *Cimex* spp.

25 such as *C. hemipterus*, *C. lectularius*; *Coccomytilus halli*, *Coccus* spp. such as *C. hesperidum*, *C. pseudomagnoliarum*, *Corythucha arcuata*, *Creontiades dilutus*, *Cryptomyzus ribis*, *Chrysomphalus aonidum*, *Cryptomyzus ribis*, *Ctenarytaina spatulata*, *Cyrtopeltis notatus*, *Dalbulus* spp., *Dasyneus piperis*, *Dialeurodes* spp. such as *D. citrifolii*; *Dalbulus maidis*, *Diaphorina* spp. such as *D. citri*; *Diaspis* spp. such as *D. bromeliae*; *Dichelops furcatus*, *Diconocoris hewetti*, *Doralis* spp., *Dreyfusia nordmanniana*, *Dreyfusia piceae*, *Drosicha* spp., *Dysaphis* spp. such as *D. plantaginea*, *D. pyri*, *D. radicola*; *Dysaulacorthum pseudosolani*, *Dysdercus* spp. such as *D. cingulatus*, *D. intermedius*; *Dysmicoccus* spp., *Edessa* spp., *Geocoris* spp., *Empoasca* spp.

30 such as *E. fabae*, *E. solana*; *Epidiaspis leperii*, *Eriosoma* spp. such as *E. lanigerum*, *E. pyricola*; *Erythroneura* spp., *Eurygaster* spp. such as *E. integriceps*; *Euscelis bilobatus*, *Euschistus* spp. such as *E. heros*, *E. impictiventris*, *E. servus*; *Fiorinia theae*, *Geococcus coffeae*, *Glycaspis brimblecombei*, *Halyomorpha* spp. such as *H. halys*; *Heliozeltis* spp., *Homalodisca vitripennis* (= *H. coagulata*), *Horcias nobilellus*, *Hyalopterus pruni*, *Hyperomyzus lactucae*, *Icerya* spp. such as *I. purchae*; *Idiocerus* spp., *Idioscopus* spp., *Laodelphax striatellus*, *Lecanium* spp., *Lecanoideus floccissimus*, *Lepidosaphes* spp. such as *L. ulmi*; *Leptocorisa* spp., *Leptoglossus phyllopus*, *Lipaphis erysimi*, *Lygus* spp. such as *L. hesperus*, *L. lineolaris*, *L. pratensis*; *Maconellicoccus hirsutus*, *Marchalina hellenica*, *Macropes excavatus*, *Macrosiphum* spp. such as *M. rosae*, *M. avenae*, *M. euphorbiae*; *Macrosteles quadrilineatus*, *Mahanarva fimbriolata*, *Meg-*

acopta cribraria, *Megoura viciae*, *Melanaphis pyrarius*, *Melanaphis sacchari*, *Melanocallis*
 (= *Tinocallis*) *caryaefoliae*, *Metcafiella* spp., *Metopolophium dirhodum*, *Monellia costalis*, *Monel-*
liopsis pecanis, *Myzocallis coryli*, *Murgantia* spp., *Myzus* spp. such as *M. ascalonicus*, *M.*
cerasi, *M. nicotianae*, *M. persicae*, *M. varians*; *Nasonovia ribis-nigri*, *Neotoxoptera formosana*,
 5 *Neomegalotomus* spp, *Nephotettix* spp. such as *N. malayanus*, *N. nigropictus*, *N. parvus*, *N.*
virescens; *Nezara* spp. such as *N. viridula*; *Nilaparvata lugens*, *Nysius huttoni*, *Oebalus* spp.
 such as *O. pugnax*; *Oncometopia* spp., *Orthezia praelonga*, *Oxycaraenus hyalinipennis*, *Para-*
bemisia myricae, *Parlatoria* spp., *Parthenolecanium* spp. such as *P. corni*, *P. persicae*; *Pemphi-*
gus spp. such as *P. bursarius*, *P. populivenae*; *Peregrinus maidis*, *Perkinsiella saccharicida*,
 10 *Phenacoccus* spp. such as *P. aceris*, *P. gossypii*; *Phloeomyzus passerinii*, *Phorodon humuli*,
Phylloxera spp. such as *P. devastatrix*, *Piesma quadrata*, *Piezodorus* spp. such as *P. guildinii*;
Pinnaspis aspidistrae, *Planococcus* spp. such as *P. citri*, *P. ficus*; *Prosapia bicincta*, *Protopulvi-*
naria pyriformis, *Psallus seriatus*, *Pseudacysta perseae*, *Pseudaulacaspis pentagona*, *Pseudo-*
coccus spp. such as *P. comstocki*; *Psylla* spp. such as *P. mali*; *Pteromalus* spp., *Pulvinaria*
 15 *amygdali*, *Pyrilla* spp., *Quadraspidotus* spp., such as *Q. perniciosus*; *Quesada gigas*, *Rastro-*
coccus spp., *Reduvius senilis*, *Rhizoecus americanus*, *Rhodnius* spp., *Rhopalomyzus ascaloni-*
cus, *Rhopalosiphum* spp. such as *R. pseudobrassicarum*, *R. insertum*, *R. maidis*, *R. padi*; *Saga-*
todes spp., *Sahlbergella singularis*, *Saissetia* spp., *Sappaphis mala*, *Sappaphis mali*, *Scapto-*
coris spp., *Scaphoides titanus*, *Schizaphis graminum*, *Schizoneura lanuginosa*, *Scotinophora*
 20 spp., *Selenaspis articulatus*, *Sitobion avenae*, *Sogata* spp., *Sogatella furcifera*, *Solubea insu-*
laris, *Spissistilus festinus* (= *Stictocephala festina*), *Stephanitis nashi*, *Stephanitis pyrioides*,
Stephanitis takeyai, *Tenalaphara malayensis*, *Tetraleurodes perseae*, *Therioaphis maculate*,
Thyanta spp. such as *T. accerra*, *T. perditor*; *Tibraca* spp., *Tomaspis* spp., *Toxoptera* spp. such
 as *T. aurantii*; *Trialeurodes* spp. such as *T. abutilonea*, *T. ricini*, *T. vaporariorum*; *Triatoma* spp.,
 25 *Trioza* spp., *Typhlocyba* spp., *Unaspis* spp. such as *U. citri*, *U. yanonensis*; and *Viteus vitifolii*,
 Insects from the order **Hymenoptera** for example *Acanthomyops interjectus*, *Athalia rosae*, *Atta*
 spp. such as *A. capiguara*, *A. cephalotes*, *A. cephalotes*, *A. laevigata*, *A. robusta*, *A. sexdens*,
A. texana, *Bombus* spp., *Brachymyrmex* spp., *Camponotus* spp. such as *C. floridanus*, *C.*
pennsylvanicus, *C. modoc*; *Cardiocondyla nuda*, *Chalibion* sp, *Crematogaster* spp., *Dasymutilla*
 30 *occidentalis*, *Diprion* spp., *Dolichovespula maculata*, *Dorymyrmex* spp., *Dryocosmus kuriphilus*,
Formica spp., *Hoplocampa* spp. such as *H. minuta*, *H. testudinea*; *Iridomyrmex humilis*, *Lasius*
 spp. such as *L. niger*, *Linepithema humile*, *Liometopum* spp., *Leptocybe invasa*, *Monomorium*
 spp. such as *M. pharaonis*, *Monomorium*, *Nylandria fulva*, *Pachycondyla chinensis*, *Paratre-*
china longicornis, *Paravespula* spp., such as *P. germanica*, *P. pennsylvanica*, *P. vulgaris*; *Phei-*
 35 *dole* spp. such as *P. megacephala*; *Pogonomyrmex* spp. such as *P. barbatus*, *P. californicus*,
Polistes rubiginosa, *Prenolepis imparis*, *Pseudomyrmex gracilis*, *Schelipron* spp., *Sirex cya-*
neus, *Solenopsis* spp. such as *S. geminata*, *S. invicta*, *S. molesta*, *S. richteri*, *S. xyloni*, *Sphe-*
cius speciosus, *Sphex* spp., *Tapinoma* spp. such as *T. melanocephalum*, *T. sessile*; *Tetramori-*
um spp. such as *T. caespitum*, *T. bicarinatum*, *Vespa* spp. such as *V. crabro*; *Vespula* spp.
 40 such as *V. squamosa*; *Wasmannia auropunctata*, *Xylocopa* sp;
 Insects from the order **Orthoptera** for example *Acheta domesticus*, *Calliptamus italicus*, *Chor-*
toicetes terminifera, *Ceuthophilus* spp., *Diastramma asynamora*, *Dociostaurus maroccanus*,
Gryllotalpa spp. such as *G. africana*, *G. gryllotalpa*; *Gryllus* spp., *Hieroglyphus daganensis*,
Kraussaria angulifera, *Locusta* spp. such as *L. migratoria*, *L. pardalina*; *Melanoplus* spp. such

as *M. bivittatus*, *M. femurrubrum*, *M. mexicanus*, *M. sanguinipes*, *M. spretus*; *Nomadacris septemfasciata*, *Oedaleus senegalensis*, *Scapteriscus* spp., *Schistocerca* spp. such as *S. americana*, *S. gregaria*, *Stemopelmatus* spp., *Tachycines asynamorus*, and *Zonozerus variegatus*;

Pests from the Class **Arachnida** for example **Acari**, e.g. of the families Argasidae, Ixodidae and

5 Sarcoptidae, such as *Amblyomma* spp. (e.g. *A. americanum*, *A. variegatum*, *A. maculatum*), Argas spp. such as *A. persicu*), *Boophilus* spp. such as *B. annulatus*, *B. decoloratus*, *B. microplus*, *Dermacentor* spp. such as *D. silvarum*, *D. andersoni*, *D. variabilis*, *Hyalomma* spp. such as *H. truncatum*, *Ixodes* spp. such as *I. ricinus*, *I. rubicundus*, *I. scapularis*, *I. holocyclus*, *I. pacificus*, *Rhipicephalus sanguineus*, *Ornithodoros* spp. such as *O. moubata*, *O. hermsi*, *O.*

10 *turicata*, *Ornithonyssus bacoti*, *Otobius megnini*, *Dermanyssus gallinae*, *Psoroptes* spp. such as *P. ovis*, *Rhipicephalus* spp. such as *R. sanguineus*, *R. appendiculatus*, *Rhipicephalus evertsi*, *Rhizoglyphus* spp., *Sarcoptes* spp. such as *S. Scabiei*, and Family **Eriophyidae** including *Aceria* spp. such as *A. sheldoni*, *A. anthocoptes*, *Acallitus* spp., *Aculops* spp. such as *A. lycopersici*, *A. pelekassi*, *Aculus* spp. such as *A. schlechtendali*; *Colomerus vitis*, *Epitrimerus pyri*, *Phyllo-*

15 *coptruta oleivora*; *Eriophytes ribis* and *Eriophyes* spp. such as *Eriophyes sheldoni*, Family **Tarsonemidae** including *Hemitarsonemus* spp., *Phytonemus pallidus* and *Polyphagotarsonemus latus*, *Stenotarsonemus* spp. *Steneotarsonemus spinki*, Family **Tenuipalpidae** including *Brevipalpus* spp. such as *B. phoenicis*; Family **Tetranychidae** including *Eotetranychus* spp., *Eutetranychus* spp., *Oligonychus* spp., *Petrobia latens*, *Tetranychus* spp. such as *T. cinnabarinus*, *T.*

20 *evansi*, *T. kanzawai*, *T. pacificus*, *T. phaseolus*, *T. telarius* and *T. urticae*; *Bryobia praetiosa*; *Panonychus* spp. such as *P. ulmi*, *P. citri*, *Metatetranychus* spp. and *Oligonychus* spp. such as *O. pratensis*, *O. perseae*, *Vasates lycopersici*, *Raoiella indica*, Family **Carpoglyphidae** including *Carpoglyphus* spp.; *Penthaleidae* spp. such as *Halotydeus destructor*, Family **Demodicidae** with species such as *Demodex* spp.; Family **Trombicidea** including *Trombicula* spp.; Family **Macro-**

25 **nyssidae** including *Ornithonyssus* spp.; Family **Pyemotidae** including *Pyemotes tritici*, *Tyrophagus putrescentiae*; Family **Acaridae** including *Acarus siro*; Family **Araneida** including *Latrodectus mactans*, *Tegenaria agrestis*, *Chiracanthium sp*, *Lycosa sp* *Achaearanea tepidariorum* and *Loxosceles reclusa*;

Pests from the Phylum **Nematoda**, for example, plant parasitic nematodes such as root-knot

30 nematodes, *Meloidogyne* spp. such as *M. hapla*, *M. incognita*, *M. javanica*; cyst-forming nematodes, *Globodera* spp. such as *G. rostochiensis*; *Heterodera* spp. such as *H. avenae*, *H. glycines*, *H. schachtii*, *H. trifolii*; Seed gall nematodes, *Anguina* spp.; Stem and foliar nematodes, *Aphelenchoides* spp. such as *A. besseyi*; Sting nematodes, *Belonolaimus* spp. such as *B. longicaudatus*; Pine nematodes, *Bursaphelenchus* spp. such as *B. lignicolus*, *B. xylophilus*; Ring

35 nematodes, *Criconema* spp., *Criconemella* spp. such as *C. xenoplax* and *C. ornata*; and, *Criconemoides* spp. such as *Criconemoides informis*; *Mesocriconema* spp.; Stem and bulb nematodes, *Ditylenchus* spp. such as *D. destructor*, *D. dipsaci*; Awl nematodes, *Dolichodorus* spp.; Spiral nematodes, *Heliocotylenchus multincinctus*; Sheath and sheathoid nematodes, *Hemicycliophora* spp. and *Hemicriconemoides* spp.; *Hirshmanniella* spp.; Lance nematodes, *Hop-*

40 *loaimus* spp.; False rootknot nematodes, *Nacobbus* spp.; Needle nematodes, *Longidorus* spp. such as *L. elongatus*; Lesion nematodes, *Pratylenchus* spp. such as *P. brachyurus*, *P. neglectus*, *P. penetrans*, *P. curvatus*, *P. goodeyi*; Burrowing nematodes, *Radopholus* spp. such as *R. similis*; *Rhadopholus* spp.; *Rhadopholus* spp.; Reniform nematodes, *Rotylenchus* spp. such as *R. robustus*, *R. reniformis*; *Scutellonema* spp.; Stubby-root nematode, *Trichodorus* spp. such as

- T. obtusus*, *T. primitivus*; *Paratrichodoros* spp. such as *P. minor*; Stunt nematodes, *Tylenchorhynchus* spp. such as *T. claytoni*, *T. dubius*; Citrus nematodes, *Tylenchulus* spp. such as *T. semipenetrans*; Dagger nematodes, *Xiphinema* spp.; and other plant parasitic nematode species;
- 5 Insects from the order **Isoptera** for example *Calotermes flavicollis*, *Coptotermes* spp. such as *C. formosanus*, *C. gestroi*, *C. acinaciformis*; *Cornitermes cumulans*, *Cryptotermes* spp. such as *C. brevis*, *C. cavifrons*; *Globitermes sulfureus*, *Heterotermes* spp. such as *H. aureus*, *H. longiceps*, *H. tenuis*; *Leucotermes flavipes*, *Odontotermes* spp., *Incisitermes* spp. such as *I. minor*, *I. Snyder*, *Marginitermes hubbardi*, *Mastotermes* spp. such as *M. darwiniensis* *Neocapritermes*
- 10 spp. such as *N. opacus*, *N. parvus*; *Neotermes* spp., *Procornitermes* spp., *Zootermopsis* spp. such as *Z. angusticollis*, *Z. nevadensis*, *Reticulitermes* spp. such as *R. hesperus*, *R. tibialis*, *R. speratus*, *R. flavipes*, *R. grassei*, *R. lucifugus*, *R. santonensis*, *R. virginicus*; *Termes natalensis*, Insects from the order **Blattaria** for example *Blatta* spp. such as *B. orientalis*, *B. lateralis*; *Blattella* spp. such as *B. asahinae*, *B. germanica*; *Leucophaea maderae*, *Panchlora nivea*, *Periplaneta*
- 15 spp. such as *P. americana*, *P. australasiae*, *P. brunnea*, *P. fuliginosa*, *P. japonica*; *Supella longipalpa*, *Parcoblatta pennsylvanica*, *Eurycotis floridana*, *Pycnoscelus surinamensis*, Insects from the order **Siphonoptera** for example *Cediopsylla simplex*, *Ceratophyllus* spp., *Ctenocephalides* spp. such as *C. felis*, *C. canis*, *Xenopsylla cheopis*, *Pulex irritans*, *Trichodectes canis*, *Tunga penetrans*, and *Nosopsyllus fasciatus*,
- 20 Insects from the order **Thysanura** for example *Lepisma saccharina*, *Ctenolepisma urbana*, and *Thermobia domestica*, Pests from the class **Chilopoda** for example *Geophilus* spp., *Scutigera* spp. such as *Scutigera coleoptrata*;
- Pests from the class **Diplopoda** for example *Blaniulus guttulatus*, *Julus* spp., *Narceus* spp.,
- 25 Pests from the class **Symphyla** for example *Scutigera immaculata*, Insects from the order **Dermoptera**, for example *Forficula auricularia*, Insects from the order **Collembola**, for example *Onychiurus* spp., such as *Onychiurus armatus*, Pests from the order **Isopoda** for example, *Armadillidium vulgare*, *Oniscus asellus*, *Porcellio scaber*,
- 30 Insects from the order **Phthiraptera**, for example *Damalinia* spp., *Pediculus* spp. such as *Pediculus humanus capitis*, *Pediculus humanus corporis*, *Pediculus humanus humanus*; *Pthirus pubis*, *Haematopinus* spp. such as *Haematopinus eurytenuis*, *Haematopinus suis*; *Linognathus* spp. such as *Linognathus vituli*; *Bovicola bovis*, *Menopon gallinae*, *Menacanthus stramineus* and *Solenopotes capillatus*, *Trichodectes* spp.,
- 35 Examples of further pest species which may be controlled by compounds of formula (I) include: from the Phylum **Mollusca**, class **Bivalvia**, for example, *Dreissena* spp.; class **Gastropoda**, for example, *Arion* spp., *Biomphalaria* spp., *Bulinus* spp., *Deroceas* spp., *Galba* spp., *Lymnaea* spp., *Oncomelania* spp., *Pomacea canaliculata*, *Succinea* spp.; from the class of the **helminths**, for example, *Ancylostoma duodenale*, *Ancylostoma ceylanicum*, *Ancylostoma braziliensis*, *Ancylostoma*
- 40 spp., *Ascaris lubricoides*, *Ascaris* spp., *Brugia malayi*, *Brugia timori*, *Bunostomum* spp., *Chabertia* spp., *Clonorchis* spp., *Cooperia* spp., *Dicrocoelium* spp., *Dictyocaulus filaria*, *Diphyllobothrium latum*, *Dracunculus medinensis*, *Echinococcus granulosus*, *Echinococcus multilocularis*, *Enterobius vermicularis*, *Faciola* spp., *Haemonchus* spp. such as *Haemonchus contortus*; *Heterakis* spp., *Hymenolepis nana*, *Hyostrogylus* spp., *Loa Loa*, *Nematodirus* spp., *Oesoph-*

agostomum spp., *Opisthorchis* spp., *Onchocerca volvulus*, *Ostertagia* spp., *Paragonimus* spp., *Schistosomen* spp., *Strongyloides fuelleborni*, *Strongyloides stercoralis*, *Strongyloides* spp., *Taenia saginata*, *Taenia solium*, *Trichinella spiralis*, *Trichinella nativa*, *Trichinella britovi*, *Trichinella nelsoni*, *Trichinella pseudospiralis*, *Trichostrongylus* spp., *Trichuris trichuria*, *Wuchereria bancrofti*.

The inventive mixture can be applied to any and all developmental stages of pests, such as egg, larva, pupa, and adult. The pests may be controlled by contacting the target pest, its food supply, habitat, breeding ground or its locus with a pesticidally effective amount of the inventive mixtures or of compositions comprising the mixtures.

"Locus" means a plant, plant propagation material (preferably seed), soil, area, material or environment in which a pest is growing or may grow.

In general, "pesticidally effective amount" means the amount of the inventive mixtures or of compositions comprising the mixtures needed to achieve an observable effect on growth, including the effects of necrosis, death, retardation, prevention, and removal, destruction, or otherwise diminishing the occurrence and activity of the target organism. The pesticidally effective amount can vary for the various mixtures / compositions used in the invention. A pesticidally effective amount of the mixtures / compositions will also vary according to the prevailing conditions such as desired pesticidal effect and duration, weather, target species, locus, mode of application, and the like.

As said above, the present invention comprises a method for improving the health of plants, wherein the plant, the locus where the plant is growing or is expected to grow or plant propagation material, from which the plant grows, is treated with an plant health effective amount of an inventive mixture.

The term "plant effective amount" denotes an amount of the inventive mixtures, which is sufficient for achieving plant health effects as defined herein below. More exemplary information about amounts, ways of application and suitable ratios to be used is given below. Anyway, the skilled artisan is well aware of the fact that such an amount can vary in a broad range and is dependent on various factors, e.g. the treated cultivated plant or material and the climatic conditions.

When preparing the mixtures, it is preferred to employ the pure active compounds, to which further active compounds against pests, such as insecticides, herbicides, fungicides or else herbicidal or growth-regulating active compounds or fertilizers can be added as further active components according to need.

The inventive mixtures are employed by treating the pests or the plants, plant propagation materials (preferably seeds), materials or soil to be protected from fungal attack with a pesticidally effective amount of the active compounds. The application can be carried out both before and after the infection of the materials, plants or plant propagation materials (preferably seeds) by the pests.

In the context of the present invention, the term plant refers to an entire plant, a part of the plant or the propagation material of the plant.

5 The inventive mixtures and compositions thereof are particularly important in the control of a multitude of phytopathogenic fungi and/or harmful insects on various cultivated plants, such as cereals, e. g. wheat, rye, barley, triticale, oats or rice; beet, e. g. sugar beet or fodder beet; fruits, such as pomes, stone fruits or soft fruits, e. g. apples, pears, plums, peaches, almonds, cherries, strawberries, raspberries, blackberries or gooseberries; leguminous plants, such as
10 lentils, peas, alfalfa or soybeans; oil plants, such as rape, mustard, olives, sunflowers, coconut, cocoa beans, castor oil plants, oil palms, ground nuts or soybeans; cucurbits, such as squashes, cucumber or melons; fiber plants, such as cotton, flax, hemp or jute; citrus fruit, such as oranges, lemons, grapefruits or mandarins; vegetables, such as spinach, lettuce, asparagus, cabbages, carrots, onions, tomatoes, potatoes, cucurbits or paprika; lauraceous plants, such as
15 avocados, cinnamon or camphor; energy and raw material plants, such as corn, soybean, rape, sugar cane or oil palm; corn; tobacco; nuts; coffee; tea; bananas; vines (Table Agrapes and grape juice grape vines); hop; turf; sweet leaf (also called Stevia); natural rubber plants or ornamental and forestry plants, such as flowers, shrubs, broad-leaved trees or evergreens, e. g. conifers; and on the plant propagation material, such as seeds, and the crop material of these
20 plants.

Preferably, the inventive mixtures and compositions thereof, respectively are used for controlling a multitude of fungi on field crops, such as potatoes, sugar beets, tobacco, wheat, rye, barley, oats, rice, corn, cotton, soybeans, rape, legumes, sunflowers, coffee or sugar cane; fruits; vines; ornamentals; or vegetables, such as cucumbers, tomatoes, beans or squashes.

25 Preferably, treatment of plant propagation materials with the inventive mixtures and compositions thereof, respectively, is used for controlling a multitude of fungi on cereals, such as wheat, rye, barley and oats; potatoes, tomatoes, vines, rice, corn, cotton and soybeans.

The term "cultivated plants" is to be understood as including plants which have been modified by breeding, mutagenesis or genetic engineering including but not limiting to agricultural
30 biotech products on the market or in development (cf. <http://cera-gmc.org/>, see GM crop database therein). Genetically modified plants are plants, which genetic material has been so modified by the use of recombinant DNA techniques that under natural circumstances cannot readily be obtained by cross breeding, mutations or natural recombination. Typically, one or more genes have been integrated into the genetic material of a genetically modified plant in order to
35 improve certain properties of the plant. Such genetic modifications also include but are not limited to targeted post-translational modification of protein(s), oligo- or polypeptides e. g. by glycosylation or polymer additions such as prenylated, acetylated or farnesylated moieties or PEG moieties.

40 Plants that have been modified by breeding, mutagenesis or genetic engineering, e. g. have been rendered tolerant to applications of specific classes of herbicides, such as auxin herbicides such as dicamba or 2,4-D; bleacher herbicides such as hydroxyphenylpyruvate dioxygenase (HPPD) inhibitors or phytoene desaturase (PDS) inhibitors; acetolactate synthase (ALS) inhibitors such as sulfonyl ureas or imidazolinones; enolpyruvylshikimate-3-phosphate synthase (EPSPS) inhibitors, such as glyphosate; glutamine synthetase (GS) inhibitors such as

glufosinate; protoporphyrinogen-IX oxidase inhibitors; lipid biosynthesis inhibitors such as acetyl CoA carboxylase (ACCase) inhibitors; or oxynil (i. e. bromoxynil or ioxynil) herbicides as a result of conventional methods of breeding or genetic engineering. Furthermore, plants have been made resistant to multiple classes of herbicides through multiple genetic modifications, such as

5 resistance to both glyphosate and glufosinate or to both glyphosate and a herbicide from another class such as ALS inhibitors, HPPD inhibitors, auxin herbicides, or ACCase inhibitors. These herbicide resistance technologies are e. g. described in Pest Management Sci. 61, 2005, 246; 61, 2005, 258; 61, 2005, 277; 61, 2005, 269; 61, 2005, 286; 64, 2008, 326; 64, 2008, 332; Weed Sci. 57, 2009, 108; Austral. J. Agricult. Res. 58, 2007, 708; Science 316, 2007, 1185; and

10 references quoted therein. Several cultivated plants have been rendered tolerant to herbicides by conventional methods of breeding (mutagenesis), e. g. Clearfield® summer rape (Canola, BASF SE, Germany) being tolerant to imidazolinones, e. g. imazamox, or ExpressSun® sunflowers (DuPont, USA) being tolerant to sulfonyl ureas, e. g. tribenuron. Genetic engineering methods have been used to render cultivated plants such as soybean, cotton, corn, beets and

15 rape, tolerant to herbicides such as glyphosate and glufosinate, some of which are commercially available under the trade names RoundupReady® (glyphosate-tolerant, Monsanto, U.S.A.), Cultivance® (imidazolinone tolerant, BASF SE, Germany) and LibertyLink® (glufosinate-tolerant, Bayer CropScience, Germany).

Furthermore, plants are also covered that are by the use of recombinant DNA techniques

20 capable to synthesize one or more insecticidal proteins, especially those known from the bacterial genus *Bacillus*, particularly from *Bacillus thuringiensis*, such as δ -endotoxins, e. g. CryIA(b), CryIA(c), CryIF, CryIF(a2), CryIIA(b), CryIIIA, CryIIIB(b1) or Cry9c; vegetative insecticidal proteins (VIP), e. g. VIP1, VIP2, VIP3 or VIP3A; insecticidal proteins of bacteria colonizing nematodes, e. g. *Photorhabdus* spp. or *Xenorhabdus* spp.; toxins produced by animals, such as

25 scorpion toxins, arachnid toxins, wasp toxins, or other insect-specific neurotoxins; toxins produced by fungi, such Streptomyces toxins, plant lectins, such as pea or barley lectins; agglutinins; proteinase inhibitors, such as trypsin inhibitors, serine protease inhibitors, patatin, cystatin or papain inhibitors; ribosome-inactivating proteins (RIP), such as ricin, maize-RIP, abrin, luffin, saporin or bryodin; steroid metabolism enzymes, such as 3-hydroxysteroid oxidase, ecdysteroid-IDP-glycosyl-transferase, cholesterol oxidases, ecdysone inhibitors or HMG-CoA-reductase;

30 ion channel blockers, such as blockers of sodium or calcium channels; juvenile hormone esterase; diuretic hormone receptors (helicokinin receptors); stilben synthase, bibenzyl synthase, chitinases or glucanases. In the context of the present invention these insecticidal proteins or toxins are to be understood expressly also as pre-toxins, hybrid proteins, truncated or otherwise

35 modified proteins. Hybrid proteins are characterized by a new combination of protein domains, (see, e. g. WO 02/015701). Further examples of such toxins or genetically modified plants capable of synthesizing such toxins are disclosed, e. g., in EP-A 374 753, WO 93/007278, WO 95/34656, EP-A 427 529, EP-A 451 878, WO 03/18810 und WO 03/52073. The methods for producing such genetically modified plants are generally known to the person skilled in the

40 art and are described, e. g. in the publications mentioned above. These insecticidal proteins contained in the genetically modified plants impart to the plants producing these proteins tolerance to harmful pests from all taxonomic groups of arthropods, especially to beetles (Coeloptera), two-winged insects (Diptera), and moths (Lepidoptera) and to nematodes (Nematoda). Genetically modified plants capable to synthesize one or more insecticidal proteins are, e. g.,

described in the publications mentioned above, and some of which are commercially available such as YieldGard® (corn cultivars producing the Cry1Ab toxin), YieldGard® Plus (corn cultivars producing Cry1Ab and Cry3Bb1 toxins), Starlink® (corn cultivars producing the Cry9c toxin), Herculex® RW (corn cultivars producing Cry34Ab1, Cry35Ab1 and the enzyme Phosphinothri-
5 c-in-N-Acetyltransferase [PAT]); NuCOTN® 33B (cotton cultivars producing the Cry1Ac toxin), Bollgard® I (cotton cultivars producing the Cry1Ac toxin), Bollgard® II (cotton cultivars producing Cry1Ac and Cry2Ab2 toxins); VIPCOT® (cotton cultivars producing a VIP-toxin); NewLeaf® (po-
tato cultivars producing the Cry3A toxin); Bt-Xtra®, NatureGard®, KnockOut®, BiteGard®, Pro-
10 tecta®, Bt11 (e. g. Agrisure® CB) and Bt176 from Syngenta Seeds SAS, France, (corn cultivars producing the Cry1Ab toxin and PAT enzyme), MIR604 from Syngenta Seeds SAS, France
(corn cultivars producing a modified version of the Cry3A toxin, c.f. WO 03/018810), MON 863
from Monsanto Europe S.A., Belgium (corn cultivars producing the Cry3Bb1 toxin), IPC 531
from Monsanto Europe S.A., Belgium (cotton cultivars producing a modified version of the
15 Cry1Ac toxin) and 1507 from Pioneer Overseas Corporation, Belgium (corn cultivars producing
the Cry1F toxin and PAT enzyme).

Furthermore, plants are also covered that are by the use of recombinant DNA techniques capable to synthesize one or more proteins to increase the resistance or tolerance of those plants to bacterial, viral or fungal pathogens. Examples of such proteins are the so-called “path-
20 ogenesis-related proteins” (PR proteins, see, e. g. EP-A 392 225), plant disease resistance genes (e. g. potato cultivars, which express resistance genes acting against *Phytophthora in-
festans* derived from the mexican wild potato *Solanum bulbocastanum*) or T4-lysozym (e. g. potato cultivars capable of synthesizing these proteins with increased resistance against bacte-
ria such as *Erwinia amylovora*). The methods for producing such genetically modified plants are
generally known to the person skilled in the art and are described, e. g. in the publications men-
25 tioned above.

Furthermore, plants are also covered that are by the use of recombinant DNA techniques capable to synthesize one or more proteins to increase the productivity (e. g. bio mass produc-
tion, grain yield, starch content, oil content or protein content), tolerance to drought, salinity or
other growth-limiting environmental factors or tolerance to pests and fungal, bacterial or viral
30 pathogens of those plants.

Furthermore, plants are also covered that contain by the use of recombinant DNA tech-
niques a modified amount of substances of content or new substances of content, specifically to
improve human or animal nutrition, e. g. oil crops that produce health-promoting long-chain
omega-3 fatty acids or unsaturated omega-9 fatty acids (e. g. Nexera® rape, DOW Agro Scienc-
35 es, Canada).

Furthermore, plants are also covered that contain by the use of recombinant DNA tech-
niques a modified amount of substances of content or new substances of content, specifically to
improve raw material production, e. g. potatoes that produce increased amounts of amylopectin
(e. g. Amflora® potato, BASF SE, Germany).

40 The separate or joint application of the compounds of the inventive mixtures is carried out by
spraying or dusting the seeds, the seedlings, the plants or the soils before or after sowing of the
plants or before or after emergence of the plants.

5 The inventive mixtures and the compositions comprising them can be used for protecting wooden materials such as trees, board fences, sleepers, etc. and buildings such as houses, out-houses, factories, but also construction materials, furniture, leathers, fibers, vinyl articles, electric wires and cables etc. from ants and/or termites, and for controlling ants and termites from doing harm to crops or human being (e.g. when the pests invade into houses and public facilities).


10 Customary application rates in the protection of materials are, for example, from 0.01 g to 1000 g of active compound per m² treated material, desirably from 0.1 g to 50 g per m².

For use in spray compositions, the content of the mixture of the active ingredients is from 0.001 to 80 weights %, preferably from 0.01 to 50 weight % and most preferably from 0.01 to 15 weight %.

Claims

1. Pesticidal mixtures comprising, as active components,

5 1) a compound of formula I



as compound I; wherein R¹ is hydrogen, methyl, methoxy or ethoxy; and

R² and R⁴ are methyl, R³ is methyl or chlorine

10 and

2) one fungicidal compound II, wherein compound II is selected from the group consisting of mefentrifluconazole, picarbutrazox, 2-[2-fluoro-6-[(8-fluoro-2-methyl-3-quinolyl)oxy]phenyl]propan-2-ol, 2-[2-[(7,8-difluoro-2-methyl-3-quinolyl)oxy]-6-fluoro-phenyl]propan-2-ol, 9-fluoro-2,2-dimethyl-5-(3-quinolyl)-3H 1,4-benzoxazepine.

15

2. A mixture as claimed in claim 1, wherein compound I is mefentrifluconazole .

3. mixture as claimed in claim 1, wherein compound II is selected from 2-[2-fluoro-6-[(8-fluoro-2-methyl-3-quinolyl)oxy]phenyl]propan-2-ol, 2-[2-[(7,8-difluoro-2-methyl-3-quinolyl)oxy]-6-fluoro-phenyl]propan-2-ol or 9-fluoro-2,2-dimethyl-5-(3-quinolyl)-3H 1,4-benzoxazepine.

20

4. A mixture as claimed in any of claims 1 to 3, wherein the ratio by weight of compound I and compound II is from 500:1 to 1:500.

25

5. A mixture as claimed in any of claims 1 to 3, wherein the ratio by weight of compound I and compound II is from 100:1 to 1:100.

6. A pesticidal composition, comprising a liquid or solid carrier and a mixture as defined in any of claims 1 to 5.

30

7. A method for controlling phytopathogenic pests, wherein the pest, their habitat, breeding grounds, their locus or the plants to be protected against pest attack, the soil or plant propagation material are treated with an effective amount of a mixture as defined in any of claims 1 to 5.

35

8. A method for improving the health of plants, wherein the plant, the locus where the plant is growing or is expected to grow or plant propagation material from which the plant grows are treated with an effective amount of a mixture as defined in any of claims 1 to 5.
- 5 9. A method for protection of plant propagation material from pests comprising contacting the plant propagation materials with a mixture as defined in any of claims 1 to 5 in pesticidally effective amounts.
- 10 10. A method as claimed in claim 9, wherein the mixture as defined in any of claims 1 to 5 is applied in an amount of from 0.01 g to 10 kg per 100 kg of plant propagation materials.
11. A method as claimed in any of claims 7 to 10, wherein the mixture as defined in any of claims 1 to 13 are applied simultaneously, that is jointly or separately, or in succession.
- 15 12. Plant propagation material, comprising the mixture as defined in any of claims 1 to 5 in an amount of from 0.01 g to 10 kg per 100 kg of plant propagation materials.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2017/075189

A. CLASSIFICATION OF SUBJECT MATTER
 INV. A01N43/90 A01N43/42 A01N43/653 A01N43/713 A01N43/72
 A01P3/00
 ADD.
 According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
 EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2009/049851 A1 (SYNGENTA PARTICIPATIONS AG [CH]; ZAMBACH WERNER [CH]; HUETER OTTMAR FR) 23 April 2009 (2009-04-23) cited in the application claim 1 claim 6 page 73, line 27 - page 74, line 35 -----	1-12
Y	EP 2 522 658 A1 (NIPPON SODA CO [JP]) 14 November 2012 (2012-11-14) cited in the application compounds 37, 125 claims 3, 4 ----- -/--	1-12

Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 14 November 2017	Date of mailing of the international search report 24/11/2017
---	--

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Götz, Gerhard
--	---

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2017/075189

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 2 762 002 A1 (NIPPON SODA CO [JP]) 6 August 2014 (2014-08-06) cited in the application tables 1-15 page 21: compound A claim 1 -----	1-12

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2017/075189

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2009049851	A1	23-04-2009	AP 2805 A 30-11-2013
			AR 068865 A1 09-12-2009
			AU 2008314078 A1 23-04-2009
			CA 2702445 A1 23-04-2009
			CN 101827815 A 08-09-2010
			CN 105061426 A 18-11-2015
			CO 6280442 A2 20-05-2011
			CR 11366 A 31-05-2010
			CY 1113631 T1 22-06-2016
			DK 2197844 T3 10-09-2012
			EA 201000544 A1 29-10-2010
			EP 2197844 A1 23-06-2010
			ES 2389230 T3 24-10-2012
			GE P20125572 B 10-07-2012
			HR P20120705 T1 30-09-2012
			JP 5502738 B2 28-05-2014
			JP 2011500522 A 06-01-2011
			KR 20100086479 A 30-07-2010
			MA 31770 B1 01-10-2010
			MY 150609 A 30-01-2014
			NZ 584178 A 30-03-2012
			PT 2197844 E 31-08-2012
			SI 2197844 T1 30-10-2012
			TN 2010000149 A1 11-11-2011
			TW 200922474 A 01-06-2009
			UA 98813 C2 25-06-2012
			US 2010311777 A1 09-12-2010
			UY 31392 A1 29-05-2009
			WO 2009049851 A1 23-04-2009

EP 2522658	A1	14-11-2012	AP 3294 A 31-05-2015
			AR 082046 A1 07-11-2012
			AU 2010339323 A1 19-07-2012
			BR 112012016111 A2 01-09-2015
			CA 2786089 A1 07-07-2011
			CA 2848577 A1 07-07-2011
			CN 102844304 A 26-12-2012
			CN 104170824 A 03-12-2014
			CR 20120321 A 24-09-2012
			CU 20120100 A7 31-07-2012
			EA 201290401 A1 30-01-2013
			EC SP12012002 A 31-08-2012
			EP 2522658 A1 14-11-2012
			GT 201200190 A 26-08-2014
			HN 2012001369 A 31-08-2015
			IL 220692 A 26-02-2015
			JP 5483485 B2 07-05-2014
			JP W02011081174 A1 13-05-2013
			KR 20120088000 A 07-08-2012
			KR 20140081897 A 01-07-2014
			MA 33851 B1 03-12-2012
			MY 160443 A 15-03-2017
			NZ 600926 A 26-07-2013
			PE 03562013 A1 06-04-2013
			SG 181976 A1 30-08-2012
			TW 201249339 A 16-12-2012
			UA 106253 C2 11-08-2014
			US 2012289702 A1 15-11-2012

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2017/075189

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		VN 32513 A1	25-02-2013
		WO 2011081174 A1	07-07-2011

EP 2762002	A1	06-08-2014	
		AU 2012317718 A1	27-03-2014
		BR 112014006574 A2	04-04-2017
		CA 2849425 A1	04-04-2013
		CN 103889229 A	25-06-2014
		CN 106973906 A	25-07-2017
		EA 201400247 A1	30-07-2014
		EP 2762002 A1	06-08-2014
		JP 6005652 B2	12-10-2016
		JP 2016169234 A	23-09-2016
		JP 2016199595 A	01-12-2016
		JP WO2013047441 A1	26-03-2015
		KR 20140049600 A	25-04-2014
		KR 20160075837 A	29-06-2016
		NZ 622017 A	31-07-2015
		UA 110657 C2	25-01-2016
		US 2014221298 A1	07-08-2014
		WO 2013047441 A1	04-04-2013
