

Gauss and the History of the Fast Fourier Transform

Michael T. Heideman
Don H. Johnson
C. Sidney Burrus

INTRODUCTION

THE fast Fourier transform (FFT) has become well known as a very efficient algorithm for calculating the discrete Fourier Transform (DFT) of a sequence of N numbers. The DFT is used in many disciplines to obtain the spectrum or frequency content of a signal, and to facilitate the computation of discrete convolution and correlation. Indeed, published work on the FFT algorithm as a means of calculating the DFT, by J. W. Cooley and J. W. Tukey in 1965 [1], was a turning point in digital signal processing and in certain areas of numerical analysis. They showed that the DFT, which was previously thought to require N^2 arithmetic operations, could be calculated by the new FFT algorithm using only $N \log N$ operations. This algorithm had a revolutionary effect on many digital processing methods, and remains the most widely used method of computing Fourier transforms [2].

In their original paper, Cooley and Tukey referred only to I. J. Good's work published in 1958 [3] as having influenced their development. However, it was soon discovered there are major differences between the Cooley-Tukey FFT and the algorithm described by Good, which is now commonly referred to as the prime factor algorithm (PFA). Soon after the appearance of the Cooley-Tukey paper, Rudnick [4] demonstrated a similar algorithm, based on the work of Danielson and Lanczos [5] which had appeared in 1942. This discovery prompted an investigation into the history of the FFT algorithm by Cooley, Lewis, and Welch [6]. They discovered that the Danielson-Lanczos paper referred to work by Runge published at the turn of the century [7, 8]. The algorithm developed by Cooley and Tukey clearly had its roots in, though perhaps not a *direct* influence from, the early twentieth century.

In a recently published history of numerical analysis [9], H. H. Goldstine attributes to Carl Friedrich Gauss, the eminent German mathematician, an algorithm similar to the FFT for the computation of the coefficients of a finite Fourier series. Gauss' treatise describing the algorithm was not published in his lifetime; it appeared only in his collected works [10] as an unpublished manuscript. The presumed year of the composition of this treatise is 1805, thereby suggesting that efficient algorithms for evaluating

coefficients of Fourier series were developed at least a century earlier than had been thought previously. If this year is accurate, it predates Fourier's 1807 work on harmonic analysis. A second reference to Gauss' algorithm was found in an article in the *Encyklopädie der Mathematischen Wissenschaften* [11], written by H. Burkhardt in 1904. It is interesting to note that Goldstine's and Burkhardt's work went almost as unnoticed as Gauss' work.

Because of Goldstine's discovery, the history of the FFT is again open to question. Is Gauss' method indeed equivalent to a modern FFT algorithm? If so, which type? And why was this work by a great mathematician not known to engineers and physicists even after the publication of Goldstine's book? What influenced Gauss' work and who developed the DFT? How firmly established is the date of writing? To answer these questions and to trace the history of Fourier series coefficient calculation into the eighteenth and nineteenth centuries, we undertook our own historical investigation¹, dealing primarily with original texts and concentrating on Gauss' work. What follows is a summary of our work, with historical references and evidence provided for readers to pursue the history as they wish. Another class of efficient DFT algorithms, called prime factor algorithms, which includes work from Thomas [13], Good [3], Winograd [14], and others, is not included in this investigation.

THE TWENTIETH CENTURY

Cooley, Lewis, and Welch [6] discovered that Danielson and Lanczos referred to the work of Carl David Tolmé Runge (1856–1927) [7, 8] as the inspiration for their algorithm. In these two papers and the book by Runge and König [15], a doubling algorithm is described which computes the Fourier transform of two N -point subsequences to obtain a $2N$ -point Fourier transform using approximately N auxiliary operations. This algorithm is not as general as the Cooley-Tukey FFT algorithm because it only allows doubling of the original sequence length, whereas the Cooley-Tukey approach efficiently computes the DFT for any multiple of the original length. The work of Runge also influenced Stumpff, who, in his book on harmonic analysis and periodograms [16], gives a doubling and tripling algorithm for the evaluation of harmonic series. Furthermore, on p. 142 of that book, he suggests a generalization to an arbitrary multiple. All of this historical information was exposed by Cooley, Lewis, and Welch [6] in much greater detail, and is mentioned here to provide

The authors are with the Electrical and Computer Engineering Department, Rice University. This research was supported by NSF grant ECS-83-14006, NASA grant NGT-44-006804, and ONR grant N00014-81-K-0565.

a background on the knowledge of the history of the FFT circa 1967.

LATE NINETEENTH AND EARLY TWENTIETH CENTURY

One interesting aspect to this historical information is that the work of Runge was well known in the early part of the twentieth century and is even referred to in the popular textbook written by Whittaker and Robinson [17], originally published in 1924. Whittaker and Robinson state that Runge's method had been widely published and cite a paper by Silvanus Phillips Thompson (1851–1916) [18]. Thompson—also a biographer of Sir William Thomson, Lord Kelvin (1824–1907)—was apparently trying to popularize Runge's method in Great Britain [18, 19]. Thompson's second paper [18] does not actually use an FFT method to obtain computational savings, but is interesting because of the discussion included at the end of the paper. This discussion includes comments by George Howard Darwin (1845–1912) (son of the more famous Charles), who claims to have used efficient techniques for the harmonic analysis of tides in 1883 [20] which he attributes to Archibald Smith (1813–1872) in 1874 [21], and to Sir Richard Strachey (1817–1908) in 1884 [22]. In Darwin's paper [20], reference is made to a paper by Joseph David Everett (1831–1904) published in 1860 [23], and credit is also given to Archibald Smith. Everett was working with Lord Kelvin on the harmonic analysis of daily temperature variations, and he gives a method for harmonic analysis using 12 samples. He claims this is an extension of the method used by Lord Kelvin in [24]. Lord Kelvin used a method based on 32 samples which was credited to Archibald Smith, originally published in 1846 [25] and presented in more detail in 1850 [26] and 1855 [27]. Apparently, S. P. Thompson was unaware of the use of efficient harmonic-analysis techniques in this paper.

The British discovery of efficient techniques for harmonic analysis can be reliably traced to Archibald Smith in 1846. Other algorithms had been developed independently by various researchers in the nineteenth century, and are tabulated on pp. 686–687 of [11]. The earliest method referenced in [11] is that of Francesco Carlini (1783–1862) from 1828 [28] for $n = 12$. The only other method which predates Archibald Smith's is that of Peter Andreas Hansen (1795–1874) from 1835 [29] for $n = 64$. Hansen was heavily influenced by Gauss in his astronomical work, but does not mention Gauss in the development of his algorithms for harmonic analysis, for reasons which shall be made clear later.

An important detail that should not be overlooked is that most of the methods preceding Runge were not intended for computing harmonics above the fourth. This was adequate for most applications of harmonic analysis in the nineteenth century, because the measurement quantization was generally on the same order of magnitude as the contributions of the higher-order harmonics. These methods are, therefore, similar to what are now called

"pruned" FFT's [30]. Most of these methods were described by computational tables for a fixed number of samples and were not presented as general techniques for computing harmonics for an arbitrary number of samples.

EARLY NINETEENTH CENTURY

Based only on the aforementioned evidence, one might conclude that FFT-type methods originated with Francesco Carlini in 1828. However, on p. 249 of *A History of Numerical Analysis from the 16th Through the 19th Century* by Herman H. Goldstine [9], the following footnote appears.

This fascinating work of Gauss was neglected and was rediscovered by Cooley and Tukey in an important paper in 1965.

This quotation refers to the treatise written by Gauss (1777–1855) entitled "Theoria Interpolationis Methodo Nova Tractata" [10]; this was published posthumously in volume 3 of his collected works in 1866, but was originally written, most likely, in 1805. Goldstine [9] gives, on p. 249–253, an English translation of parts of Gauss' paper related to trigonometric interpolation algorithms, which are Articles 25–28 of the original Latin text. Gauss wrote his important works and his personal mathematical diary in a nineteenth-century version of Latin which is now called neo-Latin. Unfortunately, neo-Latin is difficult for the casual student of classical Latin to translate accurately. Another source of difficulty for the modern reader is the notation adopted by Gauss to describe his method. Examples of this notation are the use of π as the length of a sequence (instead of N), the use of symbols $a, b, c, d, \dots; a', b', c', d', \dots; a'', b'', c'', d''$; etc. as the indices of the time series; and the use of capital letters to refer to the value of a function at a point whose index is the corresponding small letter (e.g., $f(a) = A$). Gauss' method was also derived using real trigonometric functions rather than complex exponentials, making it more difficult to relate his method to current FFT techniques.

At this point, there are three questions to be addressed. Was the method used by Gauss in [10] a form of what is now called an FFT? If so, what type and how general was it? Did Gauss realize he had developed a computationally efficient algorithm?

THE BACKGROUND OF GAUSS' WORK

The analysis of a trigonometric series goes back at least to the work of Leonhard Euler (1707–1783) [31–33]. He had no particular application in mind as far as we can tell. He dealt with infinite, cosine-only series, and did not concern himself with convergence issues. The stature of Euler in his own time meant that his work was read by his contemporaries, particularly the French mathematicians Clairaut, d'Alembert, and Lagrange. Alexis-Claude Clairaut (1713–1765) published in 1754 [34] what we currently believe to be the earliest formula for the DFT, but it was restricted to a cosine-only finite Fourier series. Joseph Louis Lagrange (1736–1813) published a DFT-like formula for sine-only series in 1762 [35]. Daniel Bernoulli (1700–1782) expressed the form of a vibrating string as a series of sine and cosine terms with arguments of both

¹This has resulted in a bibliography of over 2000 entries [12].

time and distance in 1753 [36], which implied that an arbitrary function could be expressed as an infinite sum of cosines. The most authoritative compilation of the early history of trigonometric series is a 536-page article by H. Burkhardt [37].

Clairaut and Lagrange were concerned with orbital mechanics and the problem of determining the details of an orbit from a finite set of observations. Consequently, their data was periodic and they used an interpolation approach to orbit determination: in modern terminology and notation, an even periodic function $f(x)$ which has a period of one is represented as a finite trigonometric series by

$$f(x) = \sum_{k=0}^{N-1} a_k \cos 2\pi kx, \quad 0 < x \leq 1. \quad (1)$$

The problem is to find the coefficients $\{a_k\}$ from the N values of $f(x)$ for values of $x_n = n/N$ with $n = 0, 1, \dots, N - 1$. By forcing $f(x)$ to equal the observed values at the abscissas $\{x_n\}$, one can easily show that the coefficients $\{a_k\}$ are given by the cosine-only DFT of the observed values of $f(x)$. Gauss knew of the works of Euler and Lagrange [38]: he borrowed their works from the library at Göttingen while a student from 1795 to 1798.²

GAUSS' ALGORITHM FOR COMPUTING THE DFT

Gauss extended this work on trigonometric interpolation to periodic functions, which are not necessarily odd or even. This was done while considering the problem of determining the orbit of certain asteroids from sample locations. These functions are expressed by a Fourier series of the form

$$f(x) = \sum_{k=0}^m a_k \cos 2\pi kx + \sum_{k=1}^m b_k \sin 2\pi kx \quad (2)$$

where $m = (N - 1)/2$ for N odd, or $m = N/2$ for N even. Gauss showed in Articles 19–20 of his interpolation treatise that if one were given the values of $f(x_n)$, $x_n = n/N$ ($n = 0, 1, \dots, N - 1$), the coefficients a_k and b_k are given by the now well-known formulas for the DFT [40]. This set of equations is the earliest explicit formula for the general DFT that we have found.

Gauss develops his efficient algorithm by using N_1 (or μ in his notation) equally spaced samples over one period of the signal. This set of N_1 samples is a subset of N total samples, where $N = N_1 N_2$ (or $\pi = \mu\nu$). Gauss computes the finite Fourier series which passes through these samples using m harmonics, where m is as defined in (2). He then assumes that another subset of N_1 equally spaced samples of the signal are measured which are offset from the original set of samples by a fraction, $1/N_2$, of the original sample interval where N_2 is a positive integer. A finite Fourier series with m harmonics is computed which passes through this new set of samples, and it is discovered that these coefficients are quite different from those

computed for the original N_1 samples. Gauss realized the problem and proceeded to develop a method for correcting the coefficients he had already calculated, and to determine additional coefficients for the higher-frequency harmonics. Using modern terminology, we would say that the waveform was undersampled, and that therefore the coefficients were in error because of aliasing of the high-frequency harmonics [2].

Gauss' solution to this problem was to measure a total of N_2 sets of N_1 equally spaced samples, which together form an overall set of $N = N_1 N_2$ equally spaced samples. The finite Fourier series for the entire set of N samples is computed by first computing the coefficients for each of the N_2 sets of length N_1 , all shifted relative to a common origin, and then computing coefficients of the N_1 series of length N_2 which are formed from the coefficients of corresponding terms in the N_2 sets of coefficients originally computed. A final trigonometric identity is used to convert these coefficients into the finite Fourier series coefficients for the N samples.

In modern terminology, the DFT of the samples of $f(x)$ is defined by

$$C(k) = \sum_{n=0}^{N-1} X(n) W_N^{nk} \quad (3)$$

where, if $f(x)$ has a period of one, $X(n) = f(n/N)$ are the N equally spaced samples, $W_N = e^{-j2\pi/N}$, and $k = 0, 1, \dots, N - 1$ are the indices of the Fourier coefficients. This DFT can be rewritten in terms of N_2 sets of N_1 subsamples by the change of index variables [41]

$$\begin{aligned} n &= N_2 n_1 + n_2 \\ k &= k_1 + N_1 k_2 \end{aligned}$$

for $n_1, k_1 = 0, 1, \dots, N_1 - 1$ and $n_2, k_2 = 0, 1, \dots, N_2 - 1$. Each subsequence is a function of n_1 ; n_2 denotes which subsequence it is. The DFT in (3) becomes

$$C(k_1 + N_1 k_2) = \sum_{n_2=0}^{N_2-1} \left[\sum_{n_1=0}^{N_1-1} X(N_2 n_1 + n_2) W_{N_1}^{n_1 k_1} W_N^{n_2 k_1} \right] W_{N_2}^{n_2 k_2} \quad (4)$$

where the inner sum calculates the N_2 length- N_1 DFT's corrected by W_N , and the outer sum calculates the N_1 length- N_2 DFT's. This is exactly the exponential form of Gauss' algorithm where the W_N term accounts for the shifts from the origin of the N_2 length- N_1 sequences. This is also exactly the FFT algorithm derived by Cooley and Tukey in 1965 [1] where the W_N is called a *twiddle factor* [2], a factor to correct the DFT of the inner sum for the shifted samples of $X(n)$. The equivalence of Gauss' algorithm and the Cooley-Tukey FFT is not obvious due to the notation and trigonometric formulation of Gauss. One can easily verify the results by calculating the inner sum of (4) and comparing the numerical results with the intermediate calculation in Article 28 of [10] after converting from exponential to trigonometric form and correcting for factors of $1/N_1$.

The example of $N = 12$ for the orbit of the asteroid

²Dunnington [39] has compiled a list of books borrowed by Gauss at Göttingen by searching the University library records.

DATING OF GAUSS' WORK ON THE FFT
THEORIA INTERPOLATIONIS METHODO NOVA TRACTATA
VOLUME III, WERKE

- April 30, 1777
Sept., 1795
Gauss is born in Brunswick.
Gauss arrives at Göttingen. Throughout his stay there, he checked out many books from the university library. In particular, he continually read *Miscellanea Taurinensia*, the proceedings of the academy located in Turin. When these proceedings were being published, Lagrange was there and this journal served as his exclusive outlet. In Volume III, Lagrange's DFT (sine only) appears.
- Nov. 25, 1796
Diary entry 44³, which reads *Formula interpolationis elegans*. Translated, this entry means "Elegant formula for interpolation." The editor of the diary connected this with the Lagrange interpolation formula. No specific library books can be readily connected to this entry.
- Dec. 1796
Diary entry 46, which reads *Formulae trigonometricae per series expressae*. Translated, this entry means "Trigonometric formulas expressed with series." The editor of the diary made no comment about this entry. On December 16, 1796, Gauss checked out both volumes of Euler's *Opuscula Analytica*. In this work, Euler relates algebraic series for trigonometric functions to the calculation of π . This diary entry probably refers to this kind of series.
- Sep. 28, 1798
Dec. 1804–Dec. 1805
Gauss returns home to Brunswick after finishing his studies at Göttingen.
Correspondence between Gauss and Bessel indicates their concern with the interpolation problem. No mention is made, however, of the trigonometric interpolation problem.⁴
- May 1805
Publication date of an issue of *Monatliche Correspondenz*, a collection of unreviewed notes containing astronomical observations and information. This note credits Gauss with further measurements of the orbit of the asteroid *Juno*, among which is the eccentricity value 0.254236.⁵ This number is used by Gauss in an example in his FFT writings. Thus, the treatise must have been completed after this date.
- Nov. 1805
Diary entry 124, which reads *Theoriam interpolationis ulterius excoluimus*. Translated, this entry means "We have worked out further a theory of interpolation." The editor takes this entry to mean that his treatise on interpolation could not have been written before November 1805. He refers to a notebook of Gauss consisting of short mathematical notes (*Mathematische Brouillons*), which was begun in October 1805. Volume 18 of the notebook contains an opening note on interpolation. The editor takes this note to be a first draft of the treatise. However, the collected work of Gauss does not contain this paper.
- July 30, 1806
Date attached to a letter sent from Gauss to Bode, in which the eccentricity value for *Juno* of 0.2549441 is given. Presumably, this means that the FFT treatise must have been written prior to this date. This letter appeared in *Monatliche Correspondenz* later in 1806.⁶
- Nov. 8, 1808
A letter from Schumacher, a former student of Gauss, to Gauss mentions that Schumacher's mother has a handwritten copy of his work on interpolation.⁷ It is unclear whether this letter is referring to *Mathematische Brouillons* or the *Theoria interpolationis*.
- June 8, 1816
Schumacher writes Gauss that he has a handwritten version of Gauss' work on interpolation, which he hopes Gauss will publish soon.⁷ Thus, Gauss did not keep this work secret, but presumably was not interested in publishing it.

³The entries of Gauss' mathematical diary are published with accompanying comments by the editors in volume X.1 of the *Werke*, pp. 488–571.

⁴*Briefwechsel zwischen Gauss und Bessel*, Leipzig, 1880.

⁵Volume VI, *Werke*, p. 262.

⁶Volume VI, *Werke*, p. 279.

⁷Volume X.2, *Werke*, p. 125.

Principal Discoveries of Efficient Methods of Computing the DFT

Researcher(s)	Date	Sequence Lengths	Number of DFT Values	Application
C. F. Gauss [10]	1805	Any composite integer	All	Interpolation of orbits of celestial bodies
F. Carlini [28]	1828	12	—	Harmonic analysis of barometric pressure
A. Smith [25]	1846	4, 8, 16, 32	5 or 9	Correcting deviations in compasses on ships
J. D. Everett [23]	1860	12	5	Modeling underground temperature deviations
C. Runge [7]	1903	2^nk	All	Harmonic analysis of functions
K. Stumpff [16]	1939	$2^nk, 3^nk$	All	Harmonic analysis of functions
Danielson and Lanczos [5]	1942	2^n	All	X-ray diffraction in crystals
L. H. Thomas [13]	1948	Any integer with relatively prime factors	All	Harmonic analysis of functions
I. J. Good [3]	1958	Any integer with relatively prime factors	All	Harmonic analysis of functions
Cooley and Tukey [1]	1965	Any composite integer	All	Harmonic analysis of functions
S. Winograd [14]	1976	Any integer with relatively prime factors	All	Use of complexity theory for harmonic analysis

Pallas was worked out in Article 28 of [10] for $N_1 = 4$, $N_2 = 3$ and for $N_1 = 3$, $N_2 = 4$, and an example was given in Article 41 for $N = 36$ with $N_1 = N_2 = 6$ and for the special case of odd symmetry.

In Article 27, Gauss states that his algorithm can be generalized to the case where N has more than two factors, although no examples are given. This and the observed efficiency are seen in the following translation from Article 27.

And so for this case, where most of the proposed values of the function X , an integral period of the arrangement, the number is composite and $= \pi = \mu\nu$, in articles 25, 26, we learned that through the division of that period into ν periods of μ terms, it produces, when all values are given, the same satisfactory function, which by the immediate application of the general theory applies to the whole period; truly, that method greatly reduces the tediousness of mechanical calculations, success will teach the one who tries it. Now the work will be no greater than the explanation, of how that division can be extended still further and can be applied to the case where the majority of all proposed values are composed of three or more factors, for example, if the number μ would again be composite, in that case clearly each period of μ terms can be subdivided into many lesser periods.

Gauss did not, however, go on to quantify the computational requirements of his method to obtain the now familiar $N \sum N_i$ or $N \log N$ expression for its computational complexity. From his short excerpt, Gauss clearly developed his procedure because it was computationally efficient and because it could be applied to a select, but

interesting, set of sequence lengths. Thus, Gauss' algorithm is as general and powerful as the Cooley-Tukey common-factor algorithm and is, in fact, equivalent to a decimation-in-frequency algorithm adapted to a real data sequence.

The hints used by Gauss' biographers [39, 42] and by us to establish a date for this work are summarized in the accompanying table. From these facts, we infer that Gauss wrote this treatise in October-November 1805. This work predates the 1807 work of Jean Baptiste Joseph Fourier (1768–1830) on representations of functions as infinite harmonic series. Fourier did not publish his results until 1822 [43] because his presentation to the Academy of Sciences in Paris on December 21, 1807 was not well received by Lagrange and was refused publication in the Memoirs of the Academy. One of his earlier manuscripts dates back to 1804–1805 and includes research which he may have started as early as 1802 [44].

The DFT approach to solving the orbital mechanics problem was one of several. Approaches related to Newtonian mechanics gave alternative solutions to the problem, and, in the end, came to be preferred even by Gauss. Mathematicians concerned with orbital mechanics who read his posthumous treatise in 1866 would probably not have found the technique described therein of much interest. Thus, the dated nature of the publication, its publication in Latin, and the lack of notice of Goldstine's and Burkhardt's work contributed to the "loss" of Gauss' FFT technique until now.

CONCLUSION

Although it appears that the discrete Fourier transform should really have been named after Gauss, it is obviously not practical to rename it. However, the term "Gauss-Fourier transform (GFT)" was coined by Hope [45] in 1965, and the term "discrete Gauss transform (DGT)" has also been previously used [46]. T. S. Huang was unknowingly accurate when he satirically remarked in 1971 that the FFT was Gauss' 1001st algorithm [47].

This investigation has demonstrated, once again, the virtuosity of Carl Friedrich Gauss. In addition, it has shown that certain problems can be timeless, but their solution rediscovered again and again. Burkhardt pointed out this algorithm in 1904 and Goldstine suggested the connection between Gauss and the FFT in 1977, but both of these went largely unnoticed, presumably because they were published in books dealing primarily with history. It was shown that various FFT-type algorithms were used in Great Britain and elsewhere in the nineteenth century, but were unrelated to the work of Gauss and were, in fact, not as general or well-formulated as Gauss' work. Almost one-hundred years passed between the publication of Gauss' algorithm and the modern rediscovery of this approach by Cooley and Tukey.

ACKNOWLEDGMENTS

Our work was greatly aided by the assistance of language scholars at Rice University. The Latin was translated by Dr. H. L. Eaker and Mr. W. D. Menefee. Relevant German references were translated by Dr. S. L. Clark of Rice and Professor H. W. Schüssler of the Universität Erlangen-Nürnberg.

REFERENCES

- [1] Cooley, J. W. and Tukey, J. W., "An algorithm for the machine calculation of complex Fourier series," *Math. Comput.*, vol. 19, no. 2, pp. 297-301, April 1965. Reprinted in *Digital Signal Processing*, L. R. Rabiner and C. M. Rader, Eds., pp. 223-227, New York: IEEE Press, 1972.
- [2] Brigham, E. O., *The Fast Fourier Transform*, Englewood, NJ: Prentice-Hall, Inc., 1974.
- [3] Good, I. J., "The interaction algorithm and practical Fourier analysis," *J. R. Statist. Soc. B*, vol. 20, no. 2, pp. 361-372, 1958. Addendum in *J. R. Statist. Soc. B*, vol. 22, no. 2, pp. 373-375, 1960.
- [4] Rudnick, P., "Note on the calculation of Fourier series," *Math. Comput.*, vol. 20, no. 3, pp. 429-430, July 1966.
- [5] Danielson, G. C. and Lanczos, C., "Some improvements in practical Fourier analysis and their application to X-ray scattering from liquids," *J. Franklin Inst.*, vol. 233, nos. 4 and 5, pp. 365-380 and 435-452, April and May 1942.
- [6] Cooley, J. W., Lewis, P. A. W., and Welch, P. D., "Historical notes on the fast Fourier transform," *IEEE Trans. Audio Electroacoust.*, AU-15, no. 2, pp. 76-79, June 1967. Reprinted in *Proc. IEEE*, vol. 55, no. 10, pp. 1675-1677, Oct. 1967. Also reprinted in *Digital Signal Processing*, L. R. Rabiner and C. M. Rader, Eds., pp. 260-262, New York: IEEE Press, 1972.
- [7] Runge, C., "Über die zerlegung empirisch gegebener periodischer funktionen in sinuswellen," *Z. Math. Phys.*, vol. 48, pp. 443-456, 1903.
- [8] Runge, C., "Über die zerlegung einer empirischen funktion in sinuswellen," *Z. Math. Phys.*, vol. 52, pp. 117-123, 1905.
- [9] Goldstine, H. H., *A History of Numerical Analysis from the 16th Through the 19th Century*, pp. 249-253, New York, Heidelberg, and Berlin: Springer-Verlag, 1977.
- [10] Gauss, C. F., "Nachlass: Theoria interpolationis methodo nova tractata," pp. 265-303, in *Carl Friedrich Gauss, Werke, Band 3*, Göttingen: Königlichen Gesellschaft der Wissenschaften, 1866.
- [11] Burkhardt, H., "Trigonometrische interpolation," in ch. 9, *Encyklopädie der Mathematischen Wissenschaften, vol. 2, part 1, 1st half*, pp. 643-693, Leipzig: B. G. Teubner, 1899-1916.
- [12] Heideman, M. T. and Burrus, C. S., "A Bibliography of Fast Transform and Convolution Algorithms II," Technical Report Number 8402, Electrical Engineering Dept., Rice University, Houston, TX 77251-1892, Feb. 1984.
- [13] Thomas, L. H., "Using a computer to solve problems in physics," in *Applications of Digital Computers*, Boston: Ginn, 1963.
- [14] Winograd, S., "On computing the discrete Fourier transform," *Proc. Nat. Acad. Sci. USA*, vol. 73, no. 4, pp. 1005-1006, April 1976.
- [15] Runge, C. and König, H., "Vorlesungen über numerisches rechnen," *Die Grundlehren der Mathematischen Wissenschaften*, vol. 11, pp. 211-237, Berlin: Julius Springer, 1924.
- [16] Stumpff, K., *Tafeln und Aufgaben zur Harmonischen Analyse und Periodogramrechnung*, Berlin: Julius Springer, 1939.
- [17] Whittaker, E. T. and Robinson, G., *The Calculus of Observations*, pp. 260-284, London: Blackie and Sons, Limited, 1924.
- [18] Thompson, S. P., "A new method of approximate harmonic analysis by selected ordinates," *Proc. Phys. Soc. London*, vol. 23, pp. 334-343, 1911.
- [19] Thompson, S. P., "Note on a rapid method of harmonic analysis," *Proc. Phys. Soc. London*, vol. 19, pp. 443-453, 1904.
- [20] Darwin, G. H. and Adams, J. C., "The harmonic analysis of tidal observations," *Brit. Assoc. Report*, pp. 49-118, 1883. Reprinted in *G. H. Darwin: Scientific Papers Volume I: Oceanic Tides and Lunar Disturbances of Gravity*, pp. 1-69, Cambridge: University Press, 1907.
- [21] Smith, A., *Admiralty Scientific Manual on "Deviations of the Compass"*, 5th ed., London: Hydrographic Office, Admiralty, 1874.

- [22] Strachey, R., "On the computation of the harmonic components of a series representing a phenomenon recurring in daily and yearly periods," *Proc. Roy. Soc. London*, vol. 42, no. 251, pp. 61–79, 1887.
- [23] Everett, J. D., "On a method of reducing observations of underground temperature, with its application to the monthly mean temperatures of underground thermometers at the Royal Edinburgh Observatory," *Trans. Roy. Soc. Edin.*, vol. 22, no. 2, pp. 429–439, 1860.
- [24] Thomson, W., "On the reduction of observations of underground temperature; with application to Professor Forbes' Edinburgh observations, and the continued Carlton Hill Series," *Trans. Roy. Soc. Edin.*, vol. 22, no. 2, pp. 405–427, 1860. Reprinted in *Mathematical and Physical Papers, Vol. 3* of William Thomson, Lord Kelvin, pp. 261–290, London: C. J. Clay and Sons, Cambridge Univ. Press, 1890.
- [25] Smith, A. and Sabine, E., "Contributions to terrestrial magnetism—no. VIII, memorandum on elements of reduction," *Phil. Trans. Roy. Soc. London*, vol. 136, no. 3, pp. 347–355, 1846.
- [26] Smith, A., *Instructions for the Computation of a Table of the Deviations of a Ship's Compass, From Deviations Observed on 4, 8, 16, and 32 Points, and for the Adjustment of the Table on a Change of Magnetic Latitude*, London: Gt. Brit. Hydrographic Office, 1850.
- [27] Smith, A., *Supplement to the Practical Rules for Ascertaining the Deviations of the Ship's Compass Caused by the Ship's Iron, Being Instructions for the Computation of a Table of the Deviation of the Ship's Compass from Observations Made on 4, 8, 16, or 32 Points (Second Edition) and a Graphic Method of Correcting the Deviations of a Ship's Compass*, London: J. D. Potter, 1855.
- [28] Carlini, F., *Sulla Legge Delle Variazioni Orarie del Barometro*, Modena: Presso la Tipographia Camerale, 1828. Also in *Societa Italiana della Scienze, Rome, Memorie di Matematica e di Fisica*, vol. 20, p. 198, 1828.
- [29] Hansen, P. A., *Mémoire Sur la Détermination des Perturbations Absolues dans des Ellipses d'une Excentricité et d'une Inclinaison Quelconques*. Paris: Bachelier, 1835.
- [30] Markel, J. D., "FFT pruning," *IEEE Trans. Audio Electroacoust.*, AU-19, no. 4, pp. 305–311, Dec. 1971.
- [31] Euler, L., *Introductio in Analysin Infinitorum*, Lausanne: Marc-Michel Bousquet et Comp., 1748. Reprinted in *Leonhardi Euleri Opera Omnia, Series I.*, vol. 8, A. Krazer and F. Rudio, Eds., Leipzig and Berlin: B. G. Teubner, 1922.
- [32] Euler, L., "Observationes generales circa series quarum termini secundum sinus vel cosinus angulorum multiporum progrediuntur," *Nova Acta Academiae Scientiarum Petropolitanae*, vol. 7, pp. 87–98, 1789. Reprinted in *Leonhardi Euleri Opera Omnia, Series I, vol. 16, Section 1, Commentationes Analyticae ad Theoriam Serierum Infinitarum Pertinentes*, vol. 3, Section 1, C. Boehm, Ed., pp. 163–177, Leipzig and Berlin: B. G. Teubner, 1933.
- [33] Euler, L., "Methodus facilis inveniendi series per sinus cosinusve angulorum multiporum procedentes quarum usus in universa theoria astronomiae est amplissimus," *Nova Acta Academiae Scientiarum Petropolitanae*, vol. 11, pp. 94–113, 1793. Reprinted in *Leonhardi Euleri Opera Omnia, Series I, Vol. 16, Section 1, Commentationes Analyticae ad Theoriam Serierum Infinitarum Pertinentes*, vol. 3, section 1, C. Boehm, Ed., pp. 311–332, Leipzig and Berlin: B. G. Teubner, 1933.
- [34] Clairaut, A.-C., "Mémoire sur l'orbite apparente du soleil autour de la terre, en ayant égard aux perturbations produites par les actions de la lune et des planètes principales," *Mémoires de Mathématique et de Physique de l'Académie Royale des Sciences*, no. 9, pp. 521–564, 1754.
- [35] Lagrange, J. L., "Solution de différents problèmes de calcul intégral," *Miscellanea Taurinensia ou Mélanges de Turin*, vol. III, 1762–1765. Reprinted in *Oeuvres de Lagrange*, vol. 1, J. A. Serret, Ed., pp. 469–668, Paris: Gauthier-Villars, 1867.
- [36] Bernoulli, D., "Réflexions et éclaircissemens sur les nouvelles vibrations des cordes," *Mémoires de l'Académie Royale des Sciences et Belles Lettres, Berlin*, 1753.
- [37] Burkhardt, H., "Trigonometrische reihen und integrale (bis etwa 1850)," in ch. 12, *Encyklopädie der Mathematischen Wissenschaften*, vol. 2, part 1, 2nd half, pp. 819–1354, Leipzig: B. G. Teubner, 1904–1916.
- [38] Lagrange, J. L., "Recherches sur la nature et la propagation du son," *Miscellanea Taurinensia ou Mélanges de Turin*, vol. I, no. I-X, pp. 1–112, 1759. Reprinted in *Oeuvres de Lagrange*, vol. 1, J. A. Serret, Ed., pp. 39–148, Paris: Gauthier-Villars, 1867.
- [39] Dunnington, G. W., *Carl Friedrich Gauss: Titan of Science*, pp. 398–404, New York: Exposition Press, 1955.
- [40] Bracewell, R. N., *The Fourier Transform and Its Applications*, p. 205, New York: McGraw-Hill, 1978.
- [41] Burrus, C. S., "Index mappings for multidimensional formulation of the DFT and convolution," *IEEE Trans. Acoust., Speech, Sig. Process.*, ASSP-25, no. 3, pp. 239–242, June 1977.
- [42] Merzbach, U. C., *Carl Friedrich Gauss: A Bibliography*, Wilmington, DE: Scholarly Resources Inc., 1984.
- [43] Fourier, J. B. J., *Théorie Analytique de la Chaleur*, Paris: F. Didot, 1822. English translation in *The Analytical Theory of Heat*, translated by Alexander Freeman, Cambridge: University Press, 1878.
- [44] Herivel, J., *Joseph Fourier: The Man and the Physicist*, p. 149, Oxford: Clarendon Press, 1975.
- [45] Hope, L. L., "A fast Gaussian method for Fourier transform evaluation," *Proc. IEEE*, vol. 63, no. 9, pp. 1353–1354, Sept. 1975.
- [46] Good, I. J., "Inversion of the discrete Gauss trans-

form," *Applic. Anal.*, vol. 9, no. 3, pp. 205-218, Oct. 1979.

[47] Huang, T. S., "How the fast Fourier transform got its name," *Computer*, vol. 4, no. 3, p. 15, May-June 1971.

Michael T. Heideman (S'82) was born in Medford, OR, on November 13, 1956. He received the B.S. and M.S. degrees in electrical engineering from Stanford University in 1978 and 1980, respectively. From 1978 to 1981, he was a Member of the Technical Staff of Lockheed Missiles and Space Company in Palo Alto, California, where he engaged in image processing research. From 1981 to the present, he has been with Lockheed Engineering and Management Services Company in Houston, Texas, where he is involved in the development of radar cross-section models for the space shuttle program. He is currently working toward a Ph.D. degree in electrical engineering at Rice University in Houston, TX, specializing in fast signal processing algorithms.

C. Sidney Burrus (S'55-M'61-SM'75-F'81) was born in Abilene, TX, on October 9, 1934. He received the BA, BSEE, and MS degrees from Rice University, Houston, TX, in 1957, 1958, and 1960 respectively, and the Ph.D. degree from Stanford University, Stanford, CA, in 1965.

From 1960 to 1962, he served in the Navy Reserve by teaching at the Nuclear Power School in New London, CT, and during the summer of 1964 and 1965, he was a Lecturer in Electrical Engineering at Stanford

University. In 1965, he joined the faculty at Rice University where he is now Professor of Electrical Engineering. From 1968 to 1975, he was a visiting faculty member at Baylor College of Medicine and a consultant at the VA Hospital Research Center in Houston, TX. From 1972 to 1978, he was Master of Lovett College at Rice University and in 1975, and again in 1979 he was a Visiting Professor at the Institut für Nachrichtentechnik, Universität Erlangen-Nürnberg, Germany.

Dr. Burrus is a member of Tau Beta Pi and Sigma Xi. He has received teaching awards from Rice in 1969, 74, 75, 76 and 80, received an IEEE S-ASSP Senior Award in 1974, a Senior Alexander von Humboldt Award in 1975, and a Senior Fulbright Fellowship in 1979. He was elected an IEEE Fellow in 1981.

Don H. Johnson (M'78) was born in Mt. Pleasant, TX, on July 9, 1946. He received the S.B. and S.M. degrees in 1970 and the Ph.D. degree in 1974, all from the Department of Electrical Engineering at the Massachusetts Institute of Technology, Cambridge.

He joined the MIT Lincoln Laboratory, Lexington, MA, as a Staff Member in 1974 working on digital speech systems, and is currently a Consultant there. In 1977, he joined the faculty of the Department of Electrical Engineering at Rice University, Houston, TX, where he is currently an Associate Professor. He received an American Society of Engineering Education Summer Study Fellowship in 1980 and studied passive sonar systems at the Naval Ocean Systems Center, San Diego, CA. His present research activities are in the areas of statistical communication theory as applied to digital signal processing.

Dr. Johnson is a member of Eta Kappa Nu and Tau Beta Pi.

Call for Papers International Conference on MATHEMATICS IN SIGNAL PROCESSING

at University of Bath, Claverton Down, Bath, UK
September 17th-19th, 1985

Signal processing represents one of the major growth areas for the application of advanced mathematical concepts that have their well-springs in approximation theory, number theory, linear algebra, time series analysis, optimisation theory, applied probability and statistics, amongst others.

The objective of the conference is to identify some of the unifying mathematics used in signal processing and to address important unresolved mathematical problems in signal processing. The aim is to pull together the common strands arising in various applications of the subject and to provide a focal point for attracting researchers into this important field.

There will be a number of distinguished invited speakers including Professor T. Kailath (Information Systems Laboratory, Stanford University, USA), who will deliver the keynote address, and Dr. E. R. Pike FRS FIMA (RSRE Malvern).

Contributed papers of up to thirty minutes presentation time are invited. These will be accepted on the basis of a 300-500 word abstract which should be sent by *December 1st, 1984*. Authors will be advised of acceptance by *March 1st, 1985*.

The following list of topics and applications are of rele-

vance to the conference: Signal modelling, Spectral estimation techniques, Inverse problems in signal processing, Least squares techniques and adaptive methods, Singular value decomposition approaches, Lattice methods and fast algorithms, Number theoretical techniques, Multidimensional filtering, Image processing and reconstruction, Detection and estimation, Encryption and coding technique, Knowledge-based signal processing, Algorithms for parallel processing, Applications to speech, communications, radar, sonar, seismology, economics, geophysics.

Members of the Technical and Executive Committee: Professor T. S. Durrani (Strathclyde); Dr. J. B. Abbiss AFIMA (RAE Farnborough); Mr. J. H. McDonnell FIMA (Brit Aerospace); Dr. J. G. McWhirter AFIMA (RSRE Malvern); Mr. T. A. Moore (BBC Research Department); Professor H. O. Berktaay (Bath).

Abstracts and all other enquiries should be sent to: (Conference: Mathematics in Signal Processing), The Deputy Secretary, The Institute of Mathematics and its Applications, Maitland House, Warrior Square, Southend-on-Sea, Essex SS1 2JY, Telephone: (0702) 612177.