



Bearded Reedling. Adult. Male (08-X).

BEARDED REEDLING (Panurus biarmicus)

IDENTIFICATION

16-17 cm. **Adult** with chesnut upperparts, dark in **juveniles**; underparts pinkish; long tail with light outer feathers; **male** with grey head and a long moustache.



B e a r d e d Reedling. Adult. Pattern of head: t o p male; bottom female.



SIMILAR SPECIES

This species is unmistakable.

SEXING

Adult male with grey head, black moustache and black undertail coverts. Adult female with similar upperparts, without black moustache; pale undertail coverts. In old juveniles, male with bright yellow bill; female with dull yellow bill.



B e a r d e d Reedling. Adult. Sexing. Pattern of head: top male; bottom female.





Bearded Reedling. Adult. Sexing. Pattern of undertail coverts: left male; right female.





B e a r d e d R e e d l i n g . J u v e n i l e . Sexing. Colour of bill: top male; bottom female.



AGEING

2 types of age can be recognized:

Juvenile with dark upperparts; dark outer tail feathers; 10th primary with rounded tip and longer than primary coverts.

Adult without black on upperparts or with only soft streaks in some **females**; all tail feathers similar; 10th primary with pointed tip and the same length than primary coverts.

After the **postbreeding/postjuvenile** moult, ageing is not possible using plumage characters.






Bearded Reedling. Ageing. Pattern of upperparts: left adult female; right juvenile.





Bearded Reedling. Ageing. Pattern of the outermost tail feathers: left adult; right juvenile.





Bearded Reedling. Ageing. Length of the 10th primary: top adult; bottom juvenile.

MOUL

Complete **postbreeding** and **postjuvenile** moult, usually finished late October.

PHENOLOGY



STATUS IN ARAGON

Resident species related with Reed fields. Breeds in Cinco Villas and Sariñena.



Bearded Reedling. Adult. Female (8-X).





Bearded Reedling. Juvenile. Female (30-VI).






B e a r d e d R e e d l i n g . Juvenile. Head pattern and bill colour: top male (08 - V I I); bottom female (30-VI).



Bearded Reedling. Juvenile. Male (08-VII).









B e a r d e d Reedling. Adult. Head pattern and bill colour: top male (08-X); bottom female (08-X).











Bearded Reedling. Pattern of tail undercoverts: top left adult male (08-X); top right adult female (08-X); left juvenile (08-VII).





Bearded Reedling. Tail pattern: left adult (08-X); right juvenile (08-VII)





Bearded Reedling. 10th primary length: left adult (08-VII); right juvenile (08-VII)



Bearded Reedling. Autumn: pattern of wing (08-X)



Bearded Reedling. Summer. Adult: : pattern of wing (08-VII)



Bearded Reedling. Summer. Juvenile: pattern of wing (08-VII).