iberCaja Obra Social





Barn Owl. Male (26-I).

BARN OWL (Tyto alba)

IDENTIFICATION

33-39 cm. Brown golden upperparts, with grey feathers; white or buff underparts: white and heart-shape face; dark eyes,





Barn Owl. Pattern of heade and facial mask

SIMILAR SPECIES

This species is unmistakable.

SEXING

Male lacks, or has only a little, spots on breast and underwing coverts; usually has a pale band with variable width on neck; dark tinge on upperparts pale. Female usually with very spotted breast and underwing coverts; with a small or without a pale band on neck; dark tinge on upperparts usually dark. Juveniles can be sexed with the same characters than adults. (CAUTION: many birds cannot be sexed due to overlap, so it is necessary to have the three exposed characters for a confident sexed).





Barn Owl. Sexing. Pattern of the white band on neck: left male; right female.





Barn Owl. Sexing. Pattern of spotted on breast: left male; right female.





Barn Owl. Sexing. Pattern of spotted on underwing coverts: top male; bottom female.

AGEING

4 types of age can be recognized:

Juvenile with fresh feathers and without moult limit.

2nd year autumn/3rd year spring with some outer primaries and some inner secondaries moulted; usually moulted the central pair of tail feathers.

3nd year autumn/4rd year spring with another group of outer primaries and secondaries, so they have three ages of feathers: outer and innermost **juveniles** and two more generations of moulted adult pattern feathers.

Adult with three generation of feathers, but all with **adult** pattern.

CAUTION: pattern on flightfeathers in **adult** and **juvenile** is very similar, being very difficult to find differences.

iberCaja Obra Social



MOULT

Partial **postbreeding** and **postjuvenile** moults. Moult of primaries and secondaries take place as described in **AGEING**.

PHENOLOGY

I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XII

STATUS IN ARAGON

Resident. Widely distributed throughout the Region, absent only from the highest areas of Sistema Iberico and Pyrenees.



Barn Owl. Female (26-I).





Barn Owl. Adult. Pattern of the white band on neck: left male (20-XI); right female (20-XI).





Barn Owl. Juvenile. Pattern of the white band on neck: left male (10-II); right female (25-VI)





Barn Owl. Adult. Breast pattern: left male (25-VI); right female (13-X).





Barn Owl. Juvenile. Breast pattern: left male (10-II); right female (26-I).









Barn Owl. Adult. Upperpart pattern: left male (20-XI); right female (20-XI).





Barn Owl. Juvenile. Pattern Upperpart pattern: left male (10-II); right female (25-VI)





Barn Owl. Tail pattern: left adult (25-VI); right 4th year spring (03-I).





Barn Owl. Tail pattern: left 2nd year autumn (20-XI); right juvenile (21-VI)






Barn Owl. Adult. Pattern mottled on underwing coverts: top male (20-XI); bottom female (20-XI)





Barn Owl. Juvenile. Pattern of mottled on underwing coverts: top male (26-I); b o t t o mfemale (18-VI)







Barn Owl. Adult: pattern of primaries (1 to 3 different ages of feathers where 1 is the older and 3 the newer) (25-II).





Barn Owl. 2nd year autumn: pattern of primaries (1 juvenile feathers; 2 2nd year feathers) (20-XI).





Barn Owl. 4th year spring: pattern of primaries (1 juvenile feathers; 2 2nd year feathers; 3 3rd year feathers) (03-I).





Barn Owl. Juvenile: pattern of primaries (26-I).





Barn Owl. Adult: pattern of secondaries (1 to 3 different ages of feathers where 1 is the older and 3 the newer) (25-II).



Barn Owl. 2nd year autumn: pattern of secondaries (1 juvenile feathers; 2 2nd year feathers) (20-XI).





Barn Owl. 4th year spring: pattern of secondaries (1 juvenile feather; 2 2nd year feathers; 3 3rd year feathers) (03-I).





Barn Owl. Juvenile: pattern of secondaries (26-I).







Barn Owl. Adult: pattern of wing (1 to 3 different ages of feathers where 1 is the older and 3 the newer) (25-II).





Barn Owl. 2nd year autumn: pattern of wing (1 juvenile feathers; 2 2nd year feathers) (20-XI).





Barn Owl. 4th year spring: pattern of wing (1 juvenile feathers; 2 2nd year feathers; 3 3rd year feathers) (03-I).





Barn Owl. Juvenile: pattern of wing (26-I).