

This is a repository copy of *Palaeoproteomics resolves sloth relationships*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/147061/>

Version: Accepted Version

Article:

Presslee, Samantha Louise, Penkman, Kirsty Elizabeth Helena
orcid.org/0000-0002-6226-9799, Collins, Matthew James orcid.org/0000-0003-4226-5501
et al. (1 more author) (2019) Palaeoproteomics resolves sloth relationships. Nature
Ecology and Evolution. ISSN 2397-334X

<https://doi.org/10.1038/s41559-019-0909-z>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Palaeoproteomics resolves sloth relationships

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

Samantha Presslee^{1,2,3,¶}, Graham J. Slater^{4,¶}, François Pujos⁵, Analía M. Forasiepi⁵, Roman Fischer⁶, Kelly Molloy⁷, Meaghan Mackie^{3,8}, Jesper V. Olsen⁸, Alejandro Kramarz⁹, Matías Taglioretti¹⁰, Fernando Scaglia¹⁰, Maximiliano Lezcano¹¹, José Luis Lanata¹¹, John Southon¹², Robert Feranec¹³, Jonathan Bloch¹⁴, Adam Hajduk¹⁵, Fabiana M. Martin¹⁶, Rodolfo Salas Gismondi¹⁷, Marcelo Reguero¹⁸, Christian de Muizon¹⁹, Alex Greenwood^{20,21}, Brian T. Chait⁷, Kirsty Penkman²², Matthew Collins^{1,3}, Ross D. E. MacPhee^{2,*}

Authors' Affiliations:

¹Department of Archaeology and BioArCh, University of York, Wentworth Way, Heslington, York YO10 5NG, UK

²Department of Mammalogy, American Museum of Natural History, New York, NY 10024 USA

³Paleoproteomics Group, Natural History Museum of Denmark and University of Copenhagen 1353 Copenhagen, Denmark

⁴Department of the Geophysical Sciences, University of Chicago, 5734 S. Ellis Avenue, Chicago, IL 60637 USA

⁵Instituto Argentino de Nivología, Glaciología y Ciencias Ambientales, CCT-CONICET-Mendoza, Parque Gral San Martín, 5500 Mendoza, Argentina

⁶Target Discovery Institute, Nuffield Department of Medicine, University of Oxford, Roosevelt Drive, Oxford OX3 7FZ, UK

⁷Chait Laboratory and National Resource for the Mass Spectrometric Analysis of Biological Macromolecules, The Rockefeller University, 1230 York Avenue, New York, NY 10065 USA

24 ⁸Novo Nordisk Foundation Center for Protein Research, Blegdamsvej 3B, 2200, Copenhagen,
25 Denmark

26 ⁹Sección Paleovertebrados, Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”,
27 Av. Angel Gallardo 470, C1405DJR, Buenos Aires, Argentina.

28 ¹⁰Museo Municipal de Ciencias Naturales “Lorenzo Scaglia”, Plaza España s/n, 7600 Mar del
29 Plata, Argentina

30 ¹¹Instituto de Investigaciones en Diversidad Cultural y Procesos de Cambio, CONICET and
31 Universidad Nacional de Río Negro, 8400 Bariloche, Argentina

32 ¹²Keck-CCAMS Group, Earth System Science Department, B321 Croul Hall, University of
33 California, Irvine, Irvine, CA 92697-3100, USA

34 ¹³New York State Museum, Albany, NY 12230 USA

35 ¹⁴Florida Museum of Natural History, University of Florida, Gainesville, FL 32611-7800 USA

36 ¹⁵Museo de la Patagonia “F. P. Moreno”, Bariloche 8400, Argentina

37 ¹⁶Centro de Estudios del Hombre Austral, Instituto de la Patagonia, Universidad de Magallanes,
38 Punta Arenas, Chile

39 ¹⁷BioGeoCiencias Lab, Facultad de Ciencias y Filosofía/CIDIS, Universidad Peruana Cayetano
40 Heredia, Lima, Perú

41 ¹⁸CONICET and División Paleontología de Vertebrados, Museo de La Plata. Facultad de
42 Ciencias Naturales, Universidad Nacional de La Plata, Paseo del Bosque s/n, B1900FWA, La
43 Plata, Argentina

44 ¹⁹Centre de Recherches sur la Paléobiodiversité et les Paléoenvironnements (CR2P), UMR
45 CNRS 7207, Muséum national d'Histoire naturelle, 8, rue Buffon 75231 Paris, France

46 ²⁰Leibniz Institute for Zoo and Wildlife Research, Alfred-Kowalke-Straße 17, 10315 Berlin,
47 Germany

48 ²¹Department of Veterinary Medicine, Freie Universität Berlin, 14163 Berlin, Germany

49 ²²Department of Chemistry, University of York, Wentworth Way, Heslington, York YO10 5DD,
50 UK

51

52 ¶¶These authors contributed equally to this work.

53

54 *Author for correspondence: macphee@amnh.org

55

56 **Abstract**

57 The living tree sloths *Choloepus* and *Bradypus* are the only remaining members of Folivora, a
58 major xenarthran radiation that occupied a wide range of habitats in many parts of the western
59 hemisphere during the Cenozoic, including both continents and the West Indies. Ancient DNA
60 evidence has played only a minor role in folivoran systematics, as most sloths lived in places not
61 conducive to genomic preservation. Here we utilise collagen sequence information, both
62 separately and in combination with published mtDNA evidence, to assess the relationships of
63 tree sloths and their extinct relatives. Results from phylogenetic analysis of these datasets differ
64 substantially from morphology-based concepts: *Choloepus* groups with Mylodontidae, not
65 Megalonychidae; *Bradypus* and *Megalonyx* pair together as megatherioids, while monophyletic
66 Antillean sloths may be sister to all other folivorans. Divergence estimates are consistent with
67 fossil evidence for mid-Cenozoic presence of sloths in the West Indies and an early Miocene
68 radiation in South America.

69

70

71 The sloths (Xenarthra, Folivora), nowadays a taxonomically narrow (6 species in 2 genera)
72 component of the fauna of South and Central America^{1,2} were once a highly successful clade of
73 placental mammals as measured by higher-level diversity (Fig. 1). Diverging sometime in the
74 Palaeogene from their closest relatives, the anteaters (Vermilingua), folivorans greatly expanded
75 their diversity and range, eventually reaching North America as well as the West Indies³⁻⁸.
76 During the late Cenozoic sloth lineage diversity may have expanded and contracted several
77 times⁹. Final collapse occurred in the late Quaternary (end-Pleistocene on the continents, mid-

78 Holocene in the West Indies), leaving only the lineages that culminated in the extant two-toed
79 (*Choloepus*) and three-toed (*Bradypus*) tree sloths.

80 Radically differing from other sloth taxa in their manifold adaptations for “inverted” suspensory
81 locomotion, tree sloths have an obscure evolutionary history¹⁰. Despite their overall similarity in
82 body plans, tree sloths probably acquired their remarkable locomotor adaptations separately, one
83 of many indications that the course of folivoran evolution has been marked by detailed
84 convergences among evolutionarily distinct clades¹¹⁻¹⁹. The current consensus^{8-10,16,17} in
85 morphology-based phylogenetic treatments is to place the three-toed sloth as sister to all other
86 folivorans (Fig. 1, “eutardigrades”), while *Choloepus* is typically nested within the otherwise
87 extinct family Megalonychidae, either proximate to or actually within the group that radiated in
88 the West Indies^{3,7,11,13,16,21,22}. Although this arrangement recognizes the existence of
89 convergence in the origins of arboreality in tree sloths, it has proven difficult to effectively test.

90 Sloth palaeontology is an active field of inquiry (e.g., refs 10, 17, 22-31), but the placement of a
91 number of early Neogene clades is uncertain or disputed³² (e.g., “unallocated basal
92 megatherioids” in Fig. 1), and the nature of their relationships with the tree sloths is accordingly
93 indeterminate. This has an obvious impact on our ability to make macroevolutionary inferences¹⁴
94 (e.g., ancestral modes of locomotion) for tree sloth species, which have no known pre-
95 Quaternary fossil record¹⁰.

96 Genomic evidence, now routinely used in mammalian systematic research and phylogenetic
97 reconstruction, has so far been of limited use in evaluating these issues. Mitochondrial and at
98 least some nuclear sequence data are available for most well-defined species of living tree sloths,
99 but published ancient DNA (aDNA) evidence exists for only two late Pleistocene species³³⁻³⁶.

100 Lack of aDNA evidence is not surprising, given that the vast majority of sloth species lived in

101 temperate or tropical environments not conducive to aDNA preservation. Yet despite these
102 limitations, aDNA analyses have tentatively pointed to a set of relationships between extant
103 sloths and their extinct relatives that are very different from those implied by morphological
104 data: the three-toed sloth is consistently recovered in association with the North American
105 megatheriid *Nothrotheriops shastensis*^{34,38,39}, a position reflected in some older classifications^{13,}
106 ^{20, 21} while the two-toed sloth is firmly established as sister to the South American mylodontoid
107 *Mylodon darwini*³⁴⁻⁴⁰. This, however, is not enough information to rigorously test, with
108 molecular evidence, cladistic relationships established solely on morphological grounds.
109 There is another potential source of ancient biomolecular evidence: sequence information
110 derived from proteins⁴¹⁻⁴⁴. Because an organism's proteins are coded by its DNA, amino acid
111 sequences in a protein are directly controlled by the gene sequences which specify them.
112 Importantly, proteins—especially structural proteins like collagen and myosin—
113 characteristically degrade at a slower rate than DNA⁴⁵⁻⁴⁷. Using tandem mass spectrometry
114 coupled with high-performance liquid chromatography, it has proven possible to recover
115 authentic collagen sequence information from mammalian fossils as old as mid-Pliocene (3.5-3.8
116 Ma)⁴⁸, which exceeds the current aDNA record (560–780 kyr BP) by a substantial interval^{49, 50}.
117 Another advantage is that proteomic data can be potentially recovered from specimens from a
118 wide range of taphonomic contexts, including ones generally inimical to aDNA preservation⁵¹.
119 There are of course limitations. Bones and teeth are typically the only parts of vertebrate bodies
120 that preserve as fossils, which restricts the choice of proteins to ones that occur in significant
121 amounts in such tissues. Type 1 collagen comprises ~90% of the organic fraction of vertebrate
122 bone⁵² and is the only bone protein⁴⁶ that is well represented in taxonomically extensive libraries
123 such as the National Center for Biotechnology Information (NCBI). Since type 1 collagen is

124 coded by only 2 genes, COL 1A1 and COL 1A2, only a small fraction of a species' genome can
125 be accessed with this probe. In the context of palaeontology, phylogenetic analyses of type 1
126 collagen have been shown to yield results that are highly congruent with those produced by
127 aDNA, especially at higher taxonomic levels^{43,53}.

128 One such application is testing morphology-based hypotheses of higher-level relationships where
129 there is a strong possibility that pervasive homoplasy among and between target groups has
130 affected morphological character analysis and therefore classification, as in the case of
131 incorrectly homologized caniniform tooth loci in living tree sloths⁵⁴. Because dental features
132 have always played a large role in folivoran systematics,^{7, 10, 12, 13 16, 31} such fundamental
133 reinterpretations are likely to have a significant impact. Clearly, it is desirable to use as many
134 sources of inference as possible in reconstructing phylogeny. Also, molecular data lend
135 themselves well to estimating divergence timing of major clades—another critical problem in
136 folivoran systematics^{29, 34, 35}.

137

138 **RESULTS**

139 To address some of the questions raised in the previous section, as well as to add to the available
140 molecular database for folivorans, we utilised proteomic data collected from fossil and living
141 sloths in order to focus on three fundamental issues: (1) relationships of tree sloths to each other
142 and to other folivorans; (2) composition of folivoran superfamilies Megatherioidea and
143 Mylodontoidea; and (3) divergence dating of major sloth ingroups. Results were tested against
144 datasets that additionally incorporated published genomic and phenomic information.

145 **Samples.** A total of 120 xenarthran samples comprising 24 different genus-level taxa (see
146 Supplementary Information, Table S1 and Fig. S1) were screened for protein survival using both
147 AAR (Amino Acid Racemization) and MALDI-ToF (Matrix-Assisted Laser
148 Desorption/Ionization Time-of-Flight) mass spectrometry. Three additional xenarthran sequences
149 were taken from the literature (see Methods, Proteomic Analysis). Of these, 34 or 28.3% of the
150 total number of samples (including 31.0% of 103 folivoran samples) produced promising results
151 for both AAR and MALDI-ToF MS. From these, the best sample per taxon was selected for LC-
152 MS/MS (Liquid Chromatography-Tandem Mass Spectrometry) analysis to derive protein
153 sequences, with some additions to maximize taxonomic coverage (Fig. 2, Table 1). We
154 resampled the specimen of *Megatherium* previously utilised by ref. 44; the results presented here
155 are *de novo*. The samples of *Neocnus dousman* and *Megalocnus zile* did not pass both MALDI-
156 ToF and AAR screening criteria, but it was decided to analyse them because they were the best
157 representatives of their species. However, because coverage for the *Megalocnus* sample was
158 particularly poor, recovered sequence being mostly contaminants, it was not used in the
159 phylogenetic analyses. To provide modern comparisons, samples of *Bradypus variegatus*
160 (AMNH 20820) and *Choloepus hoffmanni* (AMNH 139772) were also subjected to LC-MS/MS
161 analysis. For further details on all samples, see Supplemental Information, especially Table S1.
162 Relevant procedures for recovering sequence information and estimating phylogenetic
163 relationships are presented in Methods.

164 Samples ranged in assigned age from late Miocene to mid-Holocene (Supplementary
165 Information Table S1), but the 19 successfully-screened samples are all Quaternary (Table 1). Of
166 these, 15 were selected for radiocarbon dating, and 10 returned finite ^{14}C ages (Supplementary
167 Information Table S2). The oldest specimen that yielded sequence information, *Glossotherium*

168 *robustum* MACN-PV 2652, is catalogued as Bonaerian SALMA (South American Land
169 Mammal Age, 128-400 ka⁵⁵), but this age assignment cannot be independently confirmed.
170 To keep nomenclature manageable, we make frequent reference to the relatively simple
171 traditional taxonomic scheme presented in Fig. 1, which is in turn based on a large simultaneous
172 analysis of folivoran relationships^{8, 16}. Significant departures from traditional frameworks will be
173 denoted where necessary by an asterisk, but only for formal taxonomic names (e.g.,
174 *Mylodontoidea, i.e., clade redefined to include *Choloepus*, not a traditional member).

175 **Phylogenetic reconstruction.** Parsimony and Bayesian topology searches resulted in largely
176 congruent topologies. Bootstrap Support (BS) under parsimony was generally low, as might be
177 expected given few variable sites, while Bayesian Posterior Probabilities (PP), which make full
178 use of the data, resulted in somewhat higher clade support (Fig. 3; see Supplementary
179 Information, Fig. S2). Although Antillean sloth relationships are not meaningfully resolved,
180 other folivorans assort into two reciprocally monophyletic clades (PP = 0.99) that are consistent
181 with aDNA results^{34, 35}. The first includes the three-toed sloth and various extinct taxa
182 traditionally considered megatherioid (PP = 0.97). The sister group relationship of *Megatherium*
183 and *Nothrotheriops* (PP = 0.93) is noncontroversial (Fig. 1), but in the Bayesian consensus we
184 unexpectedly recovered a previously unreported and moderately well-supported pairing of
185 *Megalonyx* with *Bradypus* (PP = 0.89) (see Discussion). The second monophyletic clade (BS =
186 73, PP = 1.00) consists of traditional mylodontoids plus *Choloepus*. Because inclusion of
187 *Choloepus* in this group markedly contrasts with results achieved using morphological datasets,
188 we designate this clade as *Mylodontoidea. Here, *Scelidotherium* + *Scelidodon* is the earliest
189 diverging branch and *Choloepus* is recovered as part of a clade (PP = 0.83) consistent with
190 accepted mylodontid interrelationships^{16, 31, 55}.

191 To further interrogate the reliability of our proteomic topologies, we concatenated our collagen
192 sequences with previously published mitochondrial genome sequences (hereafter, “proteomic +
193 genomic data”) for all extant folivorans (2 species of *Choloepus*, 4 species of *Bradypus*), two
194 extinct folivorans (*Myiodon darwinii* and *Nothrotheriops shastensis*) and the two extant outgroup
195 taxa^{34,35}. Bayesian analysis (Fig. S3) of the combined dataset yielded a nearly identical topology
196 to that recovered using proteomic data alone, but in this instance *Megatherioidea (including
197 *Bradypus*) and *Mylodontoidea (including *Choloepus*) were unambiguously recovered as
198 reciprocally monophyletic clades (PP = 1). Recovery of a paraphyletic *Bradypus* (with respect to
199 *Megalonyx*) is almost certainly due to a long genomic branch and lack of proteomic data for *B.*
200 *torquatus*, combined with a comparable lack of genomic data for *Megalonyx*. As the monophyly
201 of *Bradypus* has never been questioned and this result is based exclusively on relative branch
202 lengths, we constrained *Bradypus* monophyly for subsequent analyses, though analyses without a
203 constraint were not noticeably different.

204 **Molecular clock considerations and divergence time estimates.** Incorporating time as an
205 analytical component in analysis of the combined dataset yielded a well-supported and
206 monophyletic Antillean clade (PP > 0.99), although within-clade relationships were not
207 satisfactorily resolved. More unexpectedly in light of traditional taxonomic concepts, BEAST
208 placed the Antillean clade as a well-supported sister to *Megatherioidea plus *Mylodontoidea
209 (PP = 0.97) rather than pairing it with the one or the other. Support for megatherioid (PP > 0.99)
210 and mylodontoid (PP > 0.99) monophyly remained strong, but variable for constituent sub-
211 clades.

212 The relatively permissive constraints employed for calculating divergences make it difficult to
213 draw detailed conclusions regarding the tempo of sloth diversification, although mean ages in the

214 combined analysis are reasonably consistent with inferences based on genomic^{34, 35} as well as
215 morphological²⁹ data (Fig. 4; Table 2). Posterior mean node ages suggest an early Oligocene
216 origin for folivorans, with megatherioids and mylodontoids diverging in the middle to late
217 Oligocene (Deseadan SALMA) and the generally-recognized families originating within the
218 middle Miocene (Colloncuran-Laventan SALMAs). The combined analysis indicates that the last
219 time *Choloepus* and *Bradypus* shared a common ancestor was ~ 26.9 Ma (95% HPD interval,
220 17.2 - 34.4), which is notably earlier than the estimate ~ 22.36 Ma (95% HPD interval, 16.87 -
221 28.64 Ma; Figs. S5, S6) based on proteomic evidence only and more in line with some recent
222 morphological assessments (e.g., ref. 29).

223

224 **DISCUSSION**

225 In most respects, our higher-level results for Folivora are consistent with recently-published
226 morphology-only phylogenies, but the few ways in which they differ are critical because they
227 have profound implications for macroevolutionary and biogeographical inference. Harmonizing
228 morphological and molecular datasets is complicated, as the molecular results imply that
229 traditional clades exhibit a massive amount of unrecognized homoplasy—or equally
230 unrecognized plesiomorphies, incorrectly interpreted as (syn)apomorphies. Molecular analyses
231 are of course subject to the same challenges, especially in contexts like the present in which
232 samples sizes and information content are limited. It is already widely appreciated that genomic
233 information is exceptionally useful for testing phylogenetic hypotheses; so is proteomic
234 information, especially when it can be shown to be highly congruent with genetic indicators of
235 relationship⁵³. Together, as illustrated here, they provide a strong basis for formulating
236 evolutionary hypotheses:

237 ***Choloepus* is a mylodontoid.** That the two-toed sloth may be closer to traditional mylodontids
238 than to megalonychids, a possibility occasionally raised in morphological studies^{16, 24, 57}, has
239 been consistently found in recent aDNA investigations^{34, 35, 37-40}. Due to the limited number of
240 extinct taxa included in those investigations the exact nature of their relationship has remained
241 indeterminate. However, the multiple tests of phylogenetic relationships and broad taxonomic
242 sampling used in the present study substantiates the conclusion that *Choloepus* is indeed a
243 mylodontoid.

244 Given the recent ages of all of the taxa investigated, coupled with low rates of sequence
245 evolution, it is unsurprising that divergence estimates based on proteins alone suggest an
246 early/middle Miocene origin for Scelidotheriidae + Mylodontidae (including *Choloepus*).
247 Inclusion of genomic data helps to push these estimates back to the earliest Miocene, but it
248 should be noted that a number of mylodontoid sloths of late Oligocene to late Miocene/early
249 Pliocene age do not fit neatly into better-defined clades. In the past, these taxa were occasionally
250 gathered^{9, 20, 21} into the probably nonmonophyletic grouping Orophodontidae. It would be
251 interesting to know on the basis of molecular evidence whether the inclusion of a putative
252 orophodontid would affect the placement of *Choloepus*, possibly moving it stemward (Fig. S7)
253 or help refine divergence time estimates at the base of *Mylodontoidea. At present there is no
254 evidence on point; however, the youngest of these ambiguously-placed taxa, *Octodontobradys*, is
255 late Miocene/early Pliocene in age⁵⁸—young enough to stand a chance of coming within the
256 range of proteomic methods as these continue to improve.

257 **Megalocnid sloths are monophyletic, and are not part of traditional Megalonychidae.**

258 Antillean sloths have had a complex taxonomic history⁷. In the past, this geographical grouping
259 of folivorans was sometimes regarded as diphyletic, with different island taxa having diverged

260 from different mainland antecedents^{3, 7, 20, 58}. Diphyletism now seems unlikely on the basis of our
261 molecular clock results (Fig. 4; see also Supplemental Information, Figs. S5, S6) as well as
262 recent morphology-based studies^{16, 25}. Although within-clade relationships are poorly resolved
263 (cf. paraphyletic *Neocnus*), the Antillean clade as a whole resolves as strongly monophyletic (PP
264 >0.99). In light of this fact, as well as clade antiquity, it is appropriate to remove Megalocninae
265 from traditional Megalonychidae and raise it to family level (*Megalocnidae).

266 ***Megalonyx* and *Bradypus* are megatherioids.** Although recent morphology-oriented cladistic
267 studies have usually recovered *Bradypus* as sister to all other folivorans^{8-10, 16}, genomic
268 approaches^{34, 35, 39} have consistently paired the three-toed sloth with the extinct North American
269 Pleistocene megatherioid *Nothrotheriops*. On this point the proteomic data presented here are
270 fully compliant with the genomic evidence and support rejection of the inference^{9, 16} that
271 Bradypodoidea (i.e., *Bradypus*) is sister to traditional Megatherioidea + Mylodontoidea, as tested
272 by both parsimony (13 additional steps) and Bayesian inference (2*lnBayes Factor = 6.72,
273 support = Strong). Equally controversial is the sister group relationship detected between
274 *Bradypus* + *Megalonyx* (PP = 0.89 - 0.98; Fig. 4; see Supplementary Information, Fig. S3).
275 Although well supported in analyses of both collagen-only and combined proteomic + genomic
276 data, this remains a surprising finding, inasmuch as such an association has never been reported
277 in any taxon-rich phylogenetic study emphasizing morphology. While both the three-toed sloth
278 and *Megalonyx* are likely to be megatherioids cladistically, settling their deeper relationships will
279 require substantially more data than is currently available.

280 That none of the Antillean sloths used in this study showed any proteomic affinity for *Megalonyx*
281 is also surprising, because much of what has been understood to morphologically characterize
282 non-South American Megalonychidae was based on Antillean species, the fossils of which tend

283 to be far more complete than those of most other taxa conventionally included in this family^{12, 16,}
284 ¹⁷. To resolve this conflict, additional high-quality data will be required, genomic and proteomic
285 as well as phenomic. The only certainty at present is that, if *Choloepus* is excluded,
286 Megalonychidae must now be relegated to the list of formerly diverse but now completely
287 extinct folivoran families.

288 **The West Indies may have been colonized early.** An early appearance of megalocnid sloths in
289 the West Indies has been proposed on general palaeobiogeographical grounds^{3, 10, 17, 24, 60}, but at
290 present the only pre-Quaternary fossil evidence for Antillean folivorans consists of a
291 morphologically inconclusive partial femur from the early Oligocene (~31 Ma) Yauco Formation
292 of Puerto Rico⁶¹ and unassociated remains attributable to a folivoran, *Imagocnus zazaе*, from the
293 late early Miocene (~17.5 Ma) Lagunitas Formation of Cuba⁵. Although “megalonychid”
294 affinities have been assumed for both on biogeographical grounds, now no longer applicable,
295 neither has been included in formal phylogenetic analyses and their placement within Folivora
296 remains uncertain.

297 The presence of sloths in the West Indies at least as early as the early Miocene is congruent with
298 our mean age estimate (31.2 Ma; Fig. 4, Table 2) for the last common ancestor of sloths sampled
299 in this study. This inference is also roughly consistent with the GAARlandia dispersal
300 hypothesis^{5, 62}, which holds that northwestern South America and the Greater Antilles were
301 briefly in land connection during the Eocene-Oligocene transition. Without going beyond the
302 very slim body of molecular evidence currently available, there is now at least some basis for
303 hypothesizing that *Megalocnidae might represent an *in situ* Antillean radiation that was
304 emplaced on the islands during the earliest phases of the evolution of the folivoran crown-
305 group—much earlier than previously thought and inconsistent with the hypothesis of a

306 Patagonian origin for Folivora as a whole⁹. If it proves possible to acquire genomic information
307 from Greater Antillean sloth taxa known to have survived into the mid-Holocene⁶³, we may
308 expect more light to be shed on megalocnid origins.

309 Systematic repositioning of *Bradypus*, *Choloepus* and megalocnid sloths also permits a better
310 understanding of how often “extreme” arboreality arose during folivoran evolution. The living
311 tree sloths are uniquely defined among extant vertebrates by a combination of relatively rigid
312 hooklike hands and feet, marked limb mobility, extremely long arms, and powerful flexion
313 capabilities in proximal limb joints¹⁹. None of the West Indian sloths possessed all of these
314 osteological traits, but, importantly, some came close—notably the Puerco Rican species
315 *Acratocnus odontrigonus*, which may have been technically capable of hand- and foot-
316 suspension but probably did not perform the “upside-down” form of locomotion characteristic of
317 extant sloths^{7,14}. Remains assigned to the early Miocene Patagonian sloth *Eucholoeops*, possibly
318 part of a clade ancestral to the Antillean radiation, also display many features consistent with
319 highly-developed arboreality^{14, 18}. Our phylogenetic results suggest that evolutionary
320 experiments connected with life in the trees probably occurred multiple times, and early on, in
321 folivore evolution. If so, it is puzzling that small-bodied sloths with highly mobile limbs and
322 other arboreal adaptations are as yet unknown for the interval between the early Miocene (e.g.,
323 *Eucholoeops*) and the Quaternary (e.g., *Diabolotherium*)¹⁸. It is possible that their absence is
324 only apparent, if they lived in heavily forested tropical environments that do not favour
325 fossilization (e.g., mid-Cenozoic proto-Amazonia^{64, 65}).

326 The advent of molecular resources providing novel information on both extinct and extant
327 species offers new ways of testing hypotheses about relationships that, in the past, were by
328 necessity based on morphological data alone. Thanks to ongoing improvements in

329 instrumentation and applicable software, the future for palaeoproteomics should be bright if it
330 can continue to make significant contributions to solving difficult questions like the ones
331 explored here.

332 A new aDNA study⁸⁷ of folivoran phylogeny, published as this paper was going to press, reaches
333 conclusions almost identical to ours regarding the evolutionary relationships of living tree sloths
334 and the phylogenetic distinctiveness of the West Indian radiation. Because the taxonomic
335 distribution of sampled species is not identical in the two studies, there are some minor
336 differences in lower-level relationships and estimated divergence times. However, their detailed
337 agreement overall supports the argument that high-quality protein sequence information is a
338 reliable source of evidence for reconstructing phylogenetic relationships.

339

340

341 **METHODS**

342 **Proteomic Analyses**

343 The 5-number codes following taxon names in this section refer to lab sample ID numbers
344 referenced in Table 1.

345 **AAR.** Samples were prepared using a slightly modified version of the protocol in ref. 66. A
346 small sub-sample of bone (~1 mg) was hydrolysed in 7M HCL (100 µl per mg) under N₂ for 18
347 hours at 110°C. After hydrolysis, the samples were dried down overnight before being re-
348 hydrated in 0.01mM L-homo-arginine as an internal standard. The samples were analysed using
349 reversed phase high pressure liquid chromatography (RP-HPLC) following a slightly modified

350 version of the protocol developed by ref. 67. Amino acid composition and extent of racemization
351 was used to assess promising samples for sequencing.

352 **Sample preparation for MS.** The majority of samples (see Supplementary Information, Table
353 S1) were prepared using a slightly modified version of the ZooMS protocol for bone reported by
354 ref. 43. Bone samples (15-30 mg) were demineralized in 250 μ l 0.6M HCl for a minimum of 3
355 weeks at -20°C. This allowed for a gentler demineralization and helped to protect any remaining
356 collagen. After demineralization, the samples were rinsed once in 200 μ l 0.01M NaOH, and
357 three times in 200 μ l 50mM ammonium bicarbonate (Ambic). The samples were gelatinized by
358 being resuspended in 100 μ l 50mM Ambic and heated at 65°C for 1 hour before being digested
359 overnight at 37°C; 50 μ l of the heated sample was digested using 1 μ l of 0.5 μ g/ μ l porcine
360 trypsin in trypsin resuspension buffer (Promega, UK) and the other 50 μ l was dried down and
361 resuspended in 50 μ l 100mM Tris solution to be digested with elastase (Worthington; USA) at
362 the same concentration in 10% Tris solution. Two different enzymes were used to increase the
363 protein sequence coverage for LC-MS/MS^{43, 68}. Digestion was stopped by the addition of
364 trifluoroacetic acid (TFA) at a concentration of 0.5-1% of the total solution. Peptides were
365 desalted using zip-tips⁶⁴ and eluted in 100 μ l of 50% acetonitrile (ACN)/0.1% TFA (v/v).

366 **SDS-PAGE.** Selected samples were analysed using SDS PAGE (Table 1). This method was used
367 on certain samples as the standard ZooMS protocol had not yielded positive results on certain
368 samples that were deemed potentially important phylogenetically. Bone samples were crushed to
369 ~1 μ m sized particles using a Retsch PM100 ball mill cooled with liquid nitrogen. The ball mill
370 was cleaned with distilled water and methanol before and after each sample⁶⁹. Nanoscale
371 crushing allowed for the highest potential retrieval of proteomic information. 50 mg of powdered
372 sample was heated at 70°C for 10 minutes in 200 μ l SDS solubilizing buffer (0.5M Tris base, 5%

373 SDS, 130mM DTT), cysteines were alkylated by the addition of 6 μ l 1M IAA at room
374 temperature in the dark for 30 minutes before the addition of 200 μ l of dye solution (0.05%
375 bromophenol blue, 5% glycerol). 20 μ l of the samples were run on a Bis-Tris gel (NuPAGE) for
376 10 minutes to concentrate the samples into a gel plug which was briefly washed in a fixing
377 solution (16% methanol, 10% acetic acid), before being washed twice in boiling water. The gel
378 was stained using Coomassie stain.

379 The gel plug was cut into approximately 1mm sized cubes in a fume hood with a scalpel and the
380 gel cubes for each sample placed in a separate Eppendorf. The gel pieces were washed in a de-
381 staining solution (66% ammonium bicarbonate 33% acetonitrile) until no more dye could be seen
382 before being washed in the following solvents for 10 minutes per solvent; ACN, HPLC grade
383 water, ACN and 50mM ammonium bicarbonate⁷⁰. The samples were digested overnight with 100
384 μ l 3.125 μ g/ μ l trypsin in 50mM ammonium bicarbonate at 37°C and then the tryptic digest was
385 pipetted into a cleaned Eppendorf tube. 100 μ l of 70%ACN/1.7% formic acid/0.1% TFA was
386 added to the gel pieces and the gel was heated at 37°C for 1 hour with the supernatant being
387 collected and added to the tryptic digest. This step was repeated sequentially with 100mM
388 triethyl ammonium bicarbonate (TEAB) and ACN. The extracted peptides were dried down and
389 then resuspended in 5% Formic acid/0.1% TFA desalted and purified on C18 membranes
390 (Empore) before being eluted in 80% ACN/0.5% acetic acid. The purified peptides were spun to
391 dryness ready for LC-MS/MS analysis.

392 **MALDI-ToF MS.** 1 μ l of sample was spotted in triplicate onto an MTP384 Bruker ground steel
393 MALDI target plate. 1 μ l of α -cyano-4-hydroxycinnamic acid matrix solution (1% in 50%
394 Acetonitrile/0.1% Trifluoroacetic acid (v/v/v)) was added to each sample spot and mixed with

395 the sample⁴³. All samples were analysed on a Bruker Ultraflex MALDI-ToF mass spectrometer
396 in triplicate.

397 **LC-MS/MS.** Most samples were analysed at the Discovery Proteomic Facility (DPF) at Oxford
398 (Table 1). *Choloepus* 17009 and *Myiodon* 16222 were analysed at the Novo Nordisk Foundation
399 Centre for Protein Research (NNFCPR), University of Copenhagen. The *Megalonyx* sample (ID
400 16849) was run at the Laboratory of Mass Spectrometry and Gaseous Ion Chemistry, Rockefeller
401 University.

402 At DPF, sample batches were analysed on an Orbitrap Fusion Lumos or Q-Exactive with
403 identical front-end separation, employing an Easyspray column (ES803, 500mmx75µm,
404 Thermo) and a gradient of 2%-35% ACN in 0.1% FA/5%DMSO over 60 minutes. On the Fusion
405 Lumos, MS1 resolution was set to 120,000 with an AGC target of 400,000. MS2 spectra were
406 acquired in TopSpeed mode (3 seconds duty cycle) in the linear ion trap (rapid scan mode) for up
407 to 250ms, with an AGC target of 4,000 and fragmentation in CID mode (35% normalized
408 collision energy). The MS1 resolution on the Q-Exactive was set to 70,000 with an AGC target
409 of 3E6. MS2 spectra for up to 15 precursors were acquired with a resolution of 17,500 and an
410 AGC target of 1E5 for up to 128ms and 28% normalized collision energy (higher-energy
411 collision dissociation). On both instruments, precursors were excluded for 27 seconds from re-
412 selection.

413 At NNFCPR, dried peptides were resuspended in 50µl of 80% ACN and 0.1% formic acid before
414 being transferred to a 96 well plate and placed in a vacuum centrifuge at 40°C until
415 approximately 3 µL of solution was left. The samples were rehydrated with 5 or 10 µL (*Myiodon*
416 16222 and *Choloepus* 17009 respectively) of 0.1% TFA, 5% ACN. Samples were separated on a

417 15 cm column (75 μm inner diameter) in-house laser pulled and packed with 1.9 μm C18 beads
418 (Dr. Maisch, Germany) on an EASY-nLC 1000 (Proxeon, Odense, Denmark) connected to a Q-
419 Exactive HF (Thermo Scientific, Bremen, Germany) on a 77 min gradient. 5 μl of sample was
420 injected. Buffer A was milliQ water. The peptides were separated with increasing buffer B (80%
421 ACN and 0.1% formic acid), going from 5% to 80% over an 80 minute gradient and a flow rate
422 of 250 nL/min. In addition, a wash-blank injecting 2 μl 0.1% TFA, 5% ACN was run in-between
423 each sample to hinder cross-contamination.

424 The Q-Exactive HF was operated in data dependent top 10 mode. Full scan mass spectra (350-
425 1400 m/z) were recorded at a resolution of 120,000 at m/z 200 with a target value of $3e6$ and a
426 maximum injection time of 25 ms for *Choloepus* 17009 and 45ms for *Myiodon* 16222. Fragment
427 ions were recorded with a maximum ion injection time set to 108 ms and a target value set to $2e5$
428 and recorded at a resolution of 60,000 for *Choloepus* 17009 and 30,000 for *Myiodon* 16222.
429 Normalized collision energy was set at 28% and the isolation window was 1.2 m/z with the
430 dynamic exclusion set to 20 s.

431 At Rockefeller University, peptides were resuspended in 20 μL 5% methanol, 0.2% formic acid.
432 10 μL were loaded onto an EASY-Spray column (Thermo Fisher Scientific ES800: 15 cm \times 75
433 μm ID, PepMap C18, 3 μm) via an EASY-nLC 1200 and separated over a 120 minute gradient of
434 2-32% Solvent B (Solvent A = 0.1% formic acid in water, Solvent B = 0.1% formic acid, 95%
435 acetonitrile) during online ESI-MS and MS/MS analyses with a Q Exactive Plus mass
436 spectrometer (Thermo Fisher Scientific). MS/MS analyses of the top 25 precursors in each full
437 scan (300 to 1700 m/z) used the following parameters: resolution: 17,500 (at 200 Th); AGC
438 target: 2×10^5 ; maximum injection time: 200ms; isolation width: 2.0 m/z; normalized collision
439 energy: 24%.

440 **Protein sequence analysis.** The LC-MS/MS raw files were converted to MGF files using
441 Proteowizard⁷¹ and searched against a mammal collagen database which included common
442 contaminants (<http://www.thegpm.org/crap/>) in PEAKS v7.5. Mass tolerances were set at 0.5Da
443 for the fragment ions and 10ppm for precursor ions and up to 3 missed cleavages were permitted.
444 Searches allowed various post translational modifications (PTMs) including oxidation (MHW)
445 and hydroxylation of proline (both +15.99), deamidation (NQ; +0.98) and pyro-glu from E (-
446 18.01) as well as a fixed PTM of carbamidomethylation (+57.02) which occurs as part of the
447 sample preparation. A maximum of 3 PTMs were allowed per peptide. Protein tolerances were
448 set at 0.5% false discovery rate (FDR), >50% average local confidence (ALC; *de novo* only) and
449 $-10\lg P$ score ≥ 20 .

450 Sequences of both COL 1A1 and COL 1A2 were concatenated using previously published
451 mammal collagen consensus sequences taken from NCBI, including sequences for the
452 xenarthrans *Dasybus novemcinctus* (nine-banded armadillo; GenBank: XP_004470764),
453 *Cyclopes didactylus* (silky anteater; Uniprot: COHJP1/COHJP2), and *Lestodon armatus* (extinct
454 mylodontoid sloth, ref. 44). Telopeptides very rarely survive in fossil samples and so these were
455 removed from all sequences. Isoleucine and leucine cannot be differentiated using low energy
456 tandem mass spectrometry and *de novo* sequencing as both amino acids are isobaric. Therefore,
457 the identification of leucine/isoleucine was consistent throughout the sequence analyses
458 concatenated in this study. Our approach is in line with previous phylogenetic studies using
459 collagen as probe⁴³, under the assumption that MS/MS sequence variation was not interpreted as
460 significant phylogenetic change (see below, Phylogenetic Analyses).

461 Once a potential collagen sequence was compiled for a given sloth taxon, the sequence was
462 added to the collagen database and the sample was re-run through PEAKS to check for coverage

463 and sequence substitutions. Any differences noted in either the consensus sequences or between
464 different species of sloths were inspected manually. In order for a difference to be considered
465 authentic, it had to occur in more than 1 product ion spectrum and be covered by both b and y
466 ions. For additional discussion, see Supplemental Information and Table S4.

467

468 **Phylogenetic Analyses**

469 Sequences developed from the MS/MS analyses were aligned in Geneious v. 9.1.7⁷² using the
470 MUSCLE algorithm⁷³ with default settings and then checked by eye. Mitochondrial sequence
471 data for extant folivorans and *Mylodon darwini* were obtained from ref. 35 and supplemented
472 with protein coding sequences for *Nothrotheriops shastensis* from ref. 34. Because the order of
473 genes differs between these two alignments, we extracted and aligned genes for *Nothrotheriops*
474 individually using MUSCLE in Geneious, checking each by eye to ensure accuracy. Of the 2096
475 amino acids in our alignment of the type 1 collagen molecule, 134 (6.4%) were variable and 76
476 (56 % of variable sites, 3.6% of total) were parsimony informative for the taxa represented.

477 We conducted three sets of phylogenetic analyses on the resulting protein alignment (see
478 Results). We first performed a Strict Parsimony (SP) analysis using PAUP v. 4.0a (build 157)⁷⁴.
479 We employed a branch and bound search with all sites treated as unordered and equally
480 weighted. To assess clade support, we performed 10,000 bootstrap replicates using full heuristic
481 tree searches and generated a weighted 50% majority rule (MR) consensus tree from the
482 resulting sample of most-parsimonious bootstrapped trees.

483 We performed two forms of model-based phylogenetic analyses, both in a Bayesian framework.
484 We used PartitionFinder v. 2.1.1^{75, 76} to determine the most appropriate model(s) of amino-acid

485 substitution and partitioning scheme for our concatenated alignment, resulting in selection of
486 separate Dayhoff models⁷⁷ with gamma-distributed rates for COL 1A1 and COL 1A2. The first
487 set of Bayesian phylogenetic analyses used MrBayes v 3.2.5⁷⁸. We performed two Markov Chain
488 Monte Carlo (MCMC) runs, each of four chains (one cold, three heated), for 10,000,000
489 generations, sampling from the chain every 5000 generations. After checking for convergence of
490 the two chains based on Gelman-Rubin statistics and ensuring that effective sample sizes for all
491 parameters were sufficient (> 200), we discarded the first 50% of each chain as burn-in,
492 combined the remaining posterior samples and summarized them as a 50% majority rule
493 consensus tree, with clade frequencies interpreted as posterior probabilities for a given clade. To
494 determine whether our unconstrained topology provided a better explanation of the data than a
495 previously proposed morphological topology¹⁶ in which *Bradypus* is the sister lineage to all other
496 folivorans and *Choloepus*, Megalocnidae, and *Megalonyx* form a monophyletic Megalonychidae
497 (including other taxa not referenced here), we estimated the marginal likelihood of the data on
498 unconstrained and constrained topologies using the stepping stone algorithm in MrBayes. We
499 performed two runs, each with four chains (three heated, one cold) for 10,000,000 generations
500 over 50 steps, with default settings for the Alpha parameter of the Beta distribution (0.4) and
501 burn-in (-1). We calculated $2 * \ln(\ln Lk_{\text{unconstrained}} - \ln Lk_{\text{constrained}})$ from the resulting estimates and
502 assessed support using the scale in ref. 79.

503 The fact that we cannot differentiate between isoleucine and leucine using low energy tandem
504 mass spectrometry creates a unique problem for model-based phylogenetic inference procedures.
505 The standard approach in ancient protein studies⁴³ is to designate all sites with a molecular mass
506 of 131.17 g/mol as leucine, but this has the potential to bias estimates of the instantaneous rate
507 matrix, branch lengths and, possibly, topology by entirely excluding one amino acid. We

508 investigated this by replacing all peptides coded as leucine with ambiguous codings {IL} and
509 repeating Bayesian estimation of topology and branch lengths using MrBayes. The resulting 50%
510 majority rule consensus tree was identical across coding schemes, and comparison of branch
511 length estimates among analyses show no significant deviation from 1:1 ($\text{branch length}_{[\text{Leucine}]} = -$
512 $0.00009 + \text{branch length}_{[\text{ambiguous}]} * 0.96$, $R^2 = 0.995$, $p \ll 0.001$), indicating that the use of leucine
513 is appropriate. We repeated Bayesian analyses of the combined proteomic + genomic dataset
514 using the same settings but with partitioning schemes and substitution models for genetic data
515 following ref. 35.

516 We attempted to integrate our combined molecular dataset with a large, recently-published
517 morphological dataset (ref. 9). The resulting majority rule consensus tree (Fig S7) is congruent in
518 some respects with our molecular topologies (e.g., *Choloepus* was recovered as a mylodontoid
519 and *Bradypus* as a megatherioid) but other results repeatedly found in molecular analyses were
520 not obtained. In particular, we recovered a strong (PP = 1.0) traditional Megalonychidae nested
521 within Megatherioidea that included Antillean sloths minus *Choloepus*. Although the Antillean
522 species were represented in the total dataset by proteomic sequences, genomic data were
523 unavailable. This result suggests that the large number of morphological characters, some known
524 to be highly homoplastic⁵⁴, were able to swamp the signal arising from the smaller proteomic
525 dataset. While combined analysis of morphological and molecular data will ultimately be
526 necessary to fully resolve folivoran phylogeny, this exercise suggests that it is premature to
527 consider such simultaneous analyses reliable at this point in time.

528
529 Our MrBayes analyses sample tree topologies with branch lengths in units of substitutions per
530 site and so ignore temporal information inherent in phylogenetic analysis of non-
531 contemporaneous tips or external information about relative branch lengths that can be provided

532 by the fossil record. We therefore also performed a series of Bayesian tree searches assuming a
533 molecular clock under the fossilized birth-death framework⁸⁰⁻⁸², as implemented in BEAST
534 v2.5.1⁸³. Briefly, this framework allowed us to sample from the posterior distribution of time-
535 scaled trees for taxa in our proteomic dataset, inferred using their sequences and stratigraphic
536 ages, while using phylogenetically constrained fossil taxa that lack amino acid data to provide
537 additional information on relative branch lengths and divergence times. Our choice of fossil taxa
538 and topological constraints broadly followed the approach undertaken in ref. 34 for sloth
539 mitogenomes. However, our proteomic topologies raise questions about the phylogenetic
540 positioning of some fossil folivorans that have previously been considered on morphological
541 grounds as early representatives of Pleistocene and Holocene families. For example, some
542 extinct folivorans, such as the Huayquerian nothrotheriid *Mionothropus*⁸⁴ can be plausibly
543 assigned to a specific terminal branch in our proteomic topology. Others, however, are
544 customarily assigned to clades that we failed to recover. This applies to the Santacrucian taxon
545 *Eucholoeops*, usually interpreted as a basal megalonychid^{24, 85, 86} and therefore as a member of a
546 clade not found to be monophyletic in our analyses. Such issues inevitably affect efforts to
547 calibrate the proteomic + genomic data clock and to infer divergence times. Acknowledging this,
548 we employed a minimal set of constraints (see Supplemental information, Fig. S4) on the
549 positioning of fossil folivorans in our Bayesian estimation of topology and divergence times,
550 integrating over all possible placements of phylogenetically uncertain fossils using stratigraphic
551 context alone when necessary. We performed analyses with and without a monophyly constrain
552 on *Bradypus* and results did not differ at unaffected nodes.

553 The use of a Bayesian approach requires the specification of prior probabilities on model
554 parameters. We used default priors on substitution model parameters but specified the following:

555 net diversification $\sim \text{Exp}(1)$, yielding a broad, vague prior; turnover $\sim \text{beta}(2,1)$, yielding high
556 prior weight on extinction \cong speciation; sampling probability $\sim \text{beta}(2,2)$ yielding a humped
557 distribution that placed most prior weight on sampling probabilities of 0.5; origin $\sim U(61.5, 150)$
558 yielding a flat prior on ages older than 61.5 Ma up to 150 Ma. In addition, the analysis was
559 conditioned on the number of extant taxa sampled ($\rho = 0.129$ in the xenarthran proteomic
560 analyses, $\rho = 0.333$ in the folivoran proteomic analyses, $\rho = 0.266$ in the combined analyses).
561 Based on comparisons of marginal likelihoods computed via Path Sampling (see Supplementary
562 Information, Table S3), we employed a relaxed uncorrelated clock with log-normally distributed
563 rates for proteomic and combined analyses, with an exponential prior (mean=0.1) placed on the
564 mean of log-normal distribution and the default gamma $\Gamma(0.5396, 0.3819)$ on the standard
565 deviation. Two MCMC analysis were run for 10 million generations each, sampling every 1000
566 generations, after which fossils without data were pruned from the trees, the first 20% of the
567 retained samples were discarded as burn-in, the samples combined, and maximum clade
568 credibility trees constructed using the tree annotator software accompanying the BEAST suite.
569 Runs from the prior using a fixed topology (the maximum clade credibility tree based on the pre-
570 pruning sample) were used to confirm that divergence time estimates were not simply returning
571 the prior.

572 **Data availability** Mass spectrometry proteomics data have been deposited to the
573 ProteomeXchange Consortium via the PRIDE partner repository with the dataset
574 identifier PXD012859. Collagen sequences are available on the Uniprot website
575 (<https://www.uniprot.org/>); the complete list can be found in Supplemental Information, Table
576 S5. Phylogenetic datasets have been deposited on DataDryad (doi:10.5061/dryad.7dd64gs).

577

578 **Supplementary Information**

579 [URL tk]

580

581 **Acknowledgements**

582 We thank the curatorial staffs of the following museums and private collections for permission to
583 sample specimens in their care: AMNH-M, American Museum of Natural History
584 (Mammalogy), New York, USA; AMNH-P, American Museum of Natural History
585 (Paleontology), New York, USA; CIV, Iota Quatro faunal collection, courtesy of Lazaro Vinola;
586 El Trebol faunal collection, Bariloche, Argentina; FR, Forest Reserve (Trinidad) faunal
587 collection currently housed in Department of Mammalogy, AMNH, New York, USA;
588 IANIGLA-PV, Instituto Argentino de Nivología, Glaciología y Ciencias Ambientales, CCT-
589 CONICET-Mendoza, Mendoza, Argentina; MACN-PV, Museo Argentino de Ciencias Naturales
590 "Bernardino Rivadavia" (Sección Paleovertebrados), Buenos Aires, Argentina; MAPBAR,
591 Museo de la Asociación Paleontológica Bariloche (APB), prov. Río Negro, Argentina; MMP,
592 Museo Municipal de Ciencias Naturales "Lorenzo Scaglia" Mar del Plata, prov. Buenos Aires,
593 Argentina; MNHN SAO, Muséum national d'Histoire naturelle, Paris, France; MPS, Museo
594 Paleontológico "Fray Manuel de Torres", San Pedro, prov. Buenos Aires, Argentina; MUSM,
595 Museo de Historia Natural de la Universidad Nacional Mayor de San Marcos, Lima, Peru;
596 NYSM VP, New York State Museum (Vertebrate Paleontology), Albany, New York, USA; RM,
597 Cuban faunal collection currently housed in Department of Mammalogy, AMNH, New York,
598 USA; UF, University of Florida, Natural History Museum of Florida (Vertebrate Paleontology),
599 Gainesville, USA; UMAG ah, Instituto de La Patagonia, Universidad de Magallanes, Punta

600 Arenas, Chile; USNM, United States National Museum of Natural History (Paleobiology),
601 Washington DC, USA. SP would like to thank Beatrice Demarchi for useful discussion and
602 support. The authors thank the National Science Foundation for grants OPP 0636639 to RDEM
603 and DEB 1547414 to RDEM, MC, and KP.

604

605 **Author contributions**

606 RDEM, MC, and SP conceived the project. SP undertook AAR and proteomic analysis and
607 concatenated collagen sequences, with laboratory and technical assistance from RF, JO, KM,
608 MM, MC, KP, and BC. GJS conducted phylogenetic analyses. FP and AMF supplied
609 paleontological information. AK, MT, FS, ML, AH, AI, RF, JB, JLL, FMM, RSG, MR, AG,
610 CdM, and GB supplied fossil samples, locality information, species identifications, and
611 commentary on the manuscript. RDEM, SP, and GJS wrote the manuscript, with input from all
612 authors.

613

614 **Competing interests.** The authors declare no competing interests.

615

616 **Materials and Correspondence.** Address inquiries to macphee@amnh.org

617 **REFERENCES CITED**

618 ¹ Gardner, A. L. in *Mammals of South America*, **1** (ed Gardner, A. L.) 157-176 (University of
619 Chicago Press, 2007).

620

621 ² Nowak, R. *Walker's Mammals of the World: Monotremes, Marsupials, Afrotherians,*
622 *Xenarthrans, and Sundatherians* (Johns Hopkins, 2018).

623

624 ³ Kraglievich, L. Descripción de dos cráneos y otros restos del género “*Pliomorphus*”
625 Ameghino, procedentes de la formación entrerriana de las barrancas del río Paraná. *Anal. Mus.*
626 *Nac. Hist. Nat. Buenos Aires* **33**, 1-56 (1923).

627

628 ⁴ Hoffstetter, R. in *Traité de Paléontologie*, **6.2** (ed Piveteau, J.) 535-636 (Masson, 1958).

629

630 ⁵ MacPhee, R. D. E. & Iturralde-Vinent, M. A. Origin of the Greater Antillean land mammal
631 fauna 1: New Tertiary land mammals from Cuba and Puerto Rico. *Amer. Mus. Novitates* **3141**, 1-
632 31 (1995). <http://hdl.handle.net/2246/3657>

633

634 ⁶ Iturralde-Vinent, M. A. & MacPhee, R. D. E. Paleogeography of the Caribbean region:
635 implications for Cenozoic biogeography. *Bull. Amer. Mus. Nat. Hist.* **238**, 1-95 (1999).
636 <http://hdl.handle.net/2246/1642>

637

638 ⁷ White, J. & MacPhee, R.D.E. in *Biogeography of the West Indies: Patterns and Perspectives*,
639 2nd ed (eds Woods, C.A. & Sergile, F. E.) 201-236 (CRC Press, 2001).

640

641 ⁸ Gaudin T. J. & McDonald, H. G. in *The Biology of Xenarthra* (eds Vizcaíno, S. F. & Loughry,
642 W. J.) 24-36 (University Press of Florida, 2008).

643

644 ⁹ Varela, L., Tambusso, P. S., McDonald, H. G. & Fariña, R. A. Phylogeny, macroevolutionary
645 trends and historical biogeography of sloths: Insights from a Bayesian morphological clock
646 analysis. *Syst. Biol.* (2018). doi: 10.1093/sysbio/syy058

647

648 ¹⁰ Pujos, F., De Iuliis, G. & Cartelle, C. A paleogeographic overview of tropical forest sloths:
649 Towards an understanding of the origin of extant suspensory sloths? *J. Mammal. Evol.* **24** (1),
650 19-38 (2017). doi:10.1007/s10914-016-9330-4

651

652 ¹¹ Patterson, B. & Pascual, R. Evolution of mammals on southern continents. *Q. Rev. Biol.* **43** (4),
653 409-451 (1968). <https://www.jstor.org/stable/2819014>

654

655 ¹² Engelmann, G. F. in *The Evolution and Ecology of Armadillos, Sloths, and Vermilinguas* (ed
656 Montgomery, G. G.) 195-203 (Academic Press, 1985).

657

658 ¹³ Webb, S. D. in *The Evolution and Ecology of Armadillos, Sloths, and Vermilinguas* (ed
659 Montgomery, G. G.) 105-112 (Smithsonian Institution Press, 1985).

660

661 ¹⁴ White, J. Indicators of locomotor habits in xenarthrans: evidence of locomotor heterogeneity
662 among fossil sloths. *J. Vert. Paleontol.* **13** (2), 230-242 (1993).
663 <https://www.jstor.org/stable/4523502>
664

665 ¹⁵ Delsuc, F., Catzeflis, F. M., Stanhope, M. J. & Douzery, E. J. P. 2001. The evolution of
666 armadillos, anteaters and sloths depicted by nuclear and mitochondrial phylogenies: implications
667 for the status of the enigmatic fossil *Eurotamandua*. *Proc. Roy. Soc. B* **268**, 1605-1615 (2001).
668 doi: 10.1098/rspb.2001.1702
669

670 ¹⁶ Gaudin, T. J. Phylogenetic relationships among sloths (Mammalia, Xenarthra, Tardigrada): the
671 craniodental evidence. *Zool. J. Linn. Soc.* **140**, 255-305 (2004). doi: 10.1111/j.1096-
672 3642.2003.00100.x
673

674 ¹⁷ McDonald, H. G. & De Iuliis, G. in *The Biology of Xenarthra* (eds Vizcaino, S. F. & Loughry,
675 W. J.) 39-55 (University Press of Florida, 2008).
676

677 ¹⁸ Pujos, F., Gaudin, T. J., De Iuliis, G. & Cartelle, C. Recent advances on variability, morpho-
678 functional adaptations, dental terminology, and evolution of sloths. *J. Mamm. Evol.* **19**, 159-169
679 (2012). doi: 10.1007/s10914-012-9189-y
680

681 ¹⁹ Nyakatura, J. A. The convergent evolution of suspensory posture and locomotion in tree sloths.
682 *J. Mamm. Evol.* **19** (3), 225-234 (2012). doi: 10.1007/s10914-011-9174-x
683

684 ²⁰ McKenna, M. C. & Bell, S. K. *Classification of Mammals above the Species Level* (Columbia
685 University Press, 1997).

686

687 ²¹ Patterson, B., Turnbull, W. D., Segall, W. & Gaudin, T. J. The ear region in xenarthrans (=
688 Edentata: Mammalia). Part II. Pilosa (sloths, anteaters), palaeanodonts, and a miscellany.
689 *Fieldiana Geol.* **24**, 1-79 (1992). doi: 10.5962/bhl.title.3466

690

691 ²² Pujos, F. *Megatherium celendinense* sp. nov. from the Pleistocene of Peruvian Andes and the
692 megatheriine phylogenetic relationship. *Palaeontology* **49** (2): 285-306 (2006). doi:
693 10.1111/j.1475-4983.2006.00522.x

694

695 ²³ Pujos, F., De Iuliis, G. & Mamani Quispe, B. *Hiskatherium saintandrei*, gen. et sp. nov.: An
696 unusual sloth from the Santacrucian of Quebrada Honda (Bolivia) and an overview of Middle
697 Miocene, small megatherioids. *J. Vert. Paleontol.* **31** (5), 1131-1149 (2011). doi:
698 10.1080/02724634.2011.599463

699

700 ²⁴ McDonald, H. G., Rincón, A. D. & Gaudin, T. J. A new genus of megalonychid sloth
701 (Mammalia, Xenarthra) from the late Pleistocene (Lujanian) of Sierra De Perija, Zulia State,
702 Venezuela. *J. Vert. Paleontol.* **33** (5), 1226-1238 (2013). doi: 10.1080/02724634.2013.764883

703

704 ²⁵ McDonald, H. G. & Carranza-Castaneda, O. Increased xenarthran diversity of the Great
705 American Biotic Interchange: a new genus and species of ground sloth (Mammalia, Xenarthra,

706 Megalonychidae) from the Hemphillian (Late Miocene) of Jalisco, Mexico. *J. Paleontol.* **91** (5),
707 1-14 (2017). doi: 10.1017/jpa.2017.45
708
709 ²⁶ Brandoni, D. A new genus of Megalonychidae (Mammalia, Xenarthra) from the Late Miocene
710 of Argentina. *Rev. Bras. Paleontol.* **17** (1), 33-42 (2014). doi: 10.4072/rbp.2014.1.04
711
712 ²⁷ Brandoni, D. The Megalonychidae (Xenarthra, Tardigrada) from the late Miocene of Entre
713 Ríos Province, Argentina, with remarks on their systematics and biogeography. *Geobios* **44**, 33-
714 44 (2011). doi: 10.1016/j.geobios.2010.06.005
715
716 ²⁸ De Iuliis, G., Gaudin, T. J. & Vicars, M. J. A new genus and species of nothrotheriid sloth
717 (Xenarthra, Tardigrada, Nothrotheriidae) from the Late Miocene (Huayquerian) of Peru.
718 *Palaeontology* **54** (1), 171-205 (2011). doi: 10.1111/j.1475-4983.2010.01001.x
719
720 ²⁹ Gaudin, T. J. & Croft, D. Paleogene Xenarthra and the evolution of South American mammals.
721 *J. Mamm.* **96** (4), 622-634 (2015) doi: 10.1093/jmammal/gyv073
722
723 ³⁰ Rincón, A. D., Solórzano, A., McDonald, H. G. & Montellano-Ballesteros, M. Two new
724 megalonychid sloths (Mammalia: Xenarthra) from the Urumaco Formation (Late Miocene), and
725 their phylogenetic affinities. *J. Syst. Palaeontol.* **17** (5), 409-421 (2019).
726 doi:10.1080/14772019.2018.1427639
727

728 ³¹ Boscaini, A., Gaudin, T. J., Mamani Quispe, B., Antoine, P.-O. & Pujos, F. New well-
729 preserved craniodental remains of *Simomylodon uccasamamensis* (Xenarthra, Mylodontidae)
730 from the Pliocene of the Bolivian Altiplano: phylogenetic, chronostratigraphic and
731 paleobiogeographic implications. *Zool. J. Linnean Soc.* **185** (2), 459-486 (2019). doi:
732 10.1093/zoolinnea/zly075
733
734 ³² McDonald, H. G. & De Iuliis, G. in *The Biology of Xenarthra* (eds Vizcaino, S. F. & Loughry,
735 W. J.) 39-55 (University Press of Florida, 2008).
736
737 ³³ Delsuc, F. & Douzery, E. J. P. in *The Biology of Xenarthra* (eds Vizcaino, S. F. & Loughry,
738 W. J.) 11-23 (University Press of Florida, 2008).
739
740 ³⁴ Slater, G., Cui, P., Forasiepi, A. M., Lenz, D., Tsangaras, K., Voirin, B., de Moraes, N.,
741 MacPhee R. D. E. & Greenwood, A. D. Evolutionary relationships among extinct and extant
742 sloths: the evidence of mitogenomes and retroviruses. *Genome Biol. Evol.* **8** (3), 607-621 (2016).
743 doi: 10.1093/gbe/evw023
744
745 ³⁵ Delsuc, F., Kuch, M., Gibb, G. C., Hughes, J., Szpak, P., Southon, J., Enk, J., Duggan, A. T. &
746 Poinar, H. N. Resolving the phylogenetic position of Darwin's extinct ground sloth (*Mylodon*
747 *darwinii*) using mitogenomic and nuclear exon data. *Proc. Roy. Soc. B* **285**, 20180214 (2018).
748 doi: 10.1098/rspb.2018.0214
749

750 ³⁶ Moraes-Barros, N., Silva, J. A. & Morgante, J. S. Morphology, molecular phylogeny, and
751 taxonomic inconsistencies in the study of *Bradypus* sloths (Pilosa: Bradypodidae). *J. Mammal.*
752 **92** (1), 86-100 (2011). doi: 10.1644/10-MAMM-A-086.1
753

754 ³⁷ Höss, M., Dilling, A., Carrant, A. & Pääbo, S. Molecular phylogeny of the extinct ground
755 sloth *Mylodon darwini*. *Proc. Nat. Acad. Sci. USA* **93** (1), 181-185 (1996). PMID: 8552600
756

757 ³⁸ Poinar, H. N., *et al.* (1998). Molecular coproscopy: Dung and diet of the extinct ground sloth
758 *Nothrotheriops shastensis*. *Science* **281** (5375), 402-406. PMID: 9665881
759

760 ³⁹ Greenwood, A. D., Castresana, J., Feldmaier-Fuchs, G. & Pääbo, S. A molecular phylogeny of
761 two extinct sloths. *Mol. Phylogen. Evol.* **18** (1), 94-103 (2001). doi: 10.1006/mpev.2000.0860
762

763 ⁴⁰ Hofreiter, M., Betancourt, J.L., Sbriller A. P., Markgraf, V. & McDonald, H.G. Phylogeny,
764 diet, and habitat of an extinct ground sloth from Cuchillo Cura, Neuquen Province, southwest
765 Argentina. *Quat. Res.* **59** (3): 364-378 (2003). doi: 10.1016/S0033-5894(03)00030-9
766

767 ⁴¹ Welker, F., Smith, G. M., Hutson, J. M., Kindler, L., Garcia-Moreno, A., Villaluenga, A.,
768 Turner, E. & Gaudzinski-Windheuser, S. Middle Pleistocene protein sequences from the
769 rhinoceros genus *Stephanorhinus* and the phylogeny of extant and extinct Middle/Late
770 Pleistocene Rhinocerotidae. *PeerJ* **5**, e3033 (2017). doi: 10.7717/peerj.3033
771

772 ⁴² Welker, F., *et al.* Palaeoproteomic evidence identifies archaic hominins associated with the
773 Châtelperronian at the Grotte du Renne. *Proc. Nat. Acad. Sci. USA* **113** (40), 11162-11167
774 (2016). doi:org/10.1073/pnas.1605834113

775

776 ⁴³ Welker, F., *et al.* Ancient proteins resolve the evolutionary history of Darwin's South-
777 American ungulates. *Nature* **522**, 81-84 (2015). doi: 10.1038/nature14249

778

779 ⁴⁴ Buckley, M., Fariña, R. A., Lawless, C., Tambusso, P. S., Varela, L., Carlini, A. A., Powell, J.
780 E. & Martinez, J. G. Collagen sequence analysis of the extinct giant ground sloths *Lestodon* and
781 *Megatherium*. *PloS One* **10** (12), e0144793 (2015). doi: 10.1371/journal.pone.0144793

782

783 ⁴⁵ Dobberstein, R. C., Collins, M. J., Craig, O. E., Taylor, G., Penkman, K. E. H. & Ritz-Timme,
784 S. Archaeological collagen: why worry about collagen diagenesis? *Archaeol. Anthropol. Sci.* **1**
785 (1), 31-42 (2009). doi: 10.1007/s12520-009-0002-7

786

787 ⁴⁶ Buckley, M. & Collins, M. J. Collagen survival and its use for species identification in
788 Holocene-Lower Pleistocene bone fragments from British archaeological and palaeontological
789 sites. *Antiqua* **1** (1), e1 (2011). doi: 10.4081/antiqua.2011.e1

790

791 ⁴⁷ Buckley, M. & Wadsworth, C. Proteome degradation in ancient bone: diagenesis and
792 phylogenetic potential. *Palaeogeog. Palaeoclimat. Palaeoecol.* **416**, 69-79 (2014). doi:
793 10.1016/j.palaeo.2014.06.026

794

795 ⁴⁸ Rybczynski, N., Gosse, J. C., Harington, C. R., Wogelius, R. A., Hidy, A. J. & Buckley, M.
796 Mid-Pliocene warm-period deposits in the High Arctic yield insight into camel evolution. *Nat.*
797 *Comm.* **4**, 1550 (2013). doi:10.1038/ncomms2516
798
799 ⁴⁹ Allentoft, M. E., *et al.* The half-life of DNA in bone: measuring decay kinetics in 158 dated
800 fossils. *Proc. Roy. Soc. B* **279** (1748), 4724–4733 (2012). doi: 10.1098/rspb.2012.1745
801
802 ⁵⁰ Orlando, L., *et al.* Recalibrating *Equus* evolution using the genome sequence of an early
803 Middle Pleistocene horse. *Nature* **499** (7456), 74-78 (2013). doi: 10.1038/nature12323
804
805 ⁵¹ Presslee, S., *et al.* Radiocarbon dating and proteomic analysis of highly purified bone collagen
806 derived from Rancho La Brea mammal fossils. *J. Vert. Paleont.* Program & Abstracts 2016, 208
807 (2016).
808
809 ⁵² Tuross, N. & Stathoplos, L. in *Methods in Enzymology* **224** (ed Zimmer A., White, T. J., Cann,
810 R. L. & Wilson, A. C.) 121-129 (Academic Press, 1993).
811
812 ⁵³ Westbury, M., *et al.* A mitogenomic timetree for Darwin’s enigmatic “transitional” South
813 American mammal, *Macrauchenia patachonica*. *Nat. Commun.* **8**, 15951 (2017) doi:
814 10.1038/ncomms15951.
815

816 ⁵⁴ Hautier L., Gomes Rodrigues H., Billet G. & Asher R. J. The hidden teeth of sloths:
817 evolutionary vestiges and the development of a simplified dentition. *Sci. Rep.* **6**, 27763 (2016).
818 doi:10.1038/srep27763
819
820 ⁵⁵ Cione, A. L. & Tonni, E. P. in *Quaternary of South America Antarctic Península* (eds Tonni,
821 E.P. & Cione, A.L.) 23-51 (Balkema,1999).
822
823 ⁵⁶ Cartelle, C., De Iuliis, G. & Ferreira, R. L. Systematic revision of tropical Brazilian
824 scelidotheriines sloths (Xenarthra, Mylodontoidea). *J. Vertebr. Paleontol.* **29** (2), 555–566
825 (2009). doi:org/10.1671/039.029.0231
826
827 ⁵⁷ Guth, C. *La Région Temporale des Edentés* (Imprimerie Jeanne d’Arc Le Puy, 1961).
828
829 ⁵⁸ Guilherme, E., Bocquentin, J. & Porto, A. S. A new specimen of the genus *Octodontobradys*
830 (*Orophodontidae*, *Octodontobradynae*) from the Late Miocene-Pliocene of the southwestern
831 Amazon Basin, Brazil. *Anuár. Inst. Geociências* **34** (1), 64-71 (2011). www.anuario.igeo.ufrj.br
832
833 ⁵⁹ Varona, L. *Catálogo de los Mamíferos Vivientes y Extinguidos de las Antillas* (Academia de
834 Ciencias de Cuba, 1974).
835
836 ⁶⁰ Webb, S. D. & Perrigo, S. in *The Evolution and Ecology of Armadillos, Sloths, and*
837 *Vermilinguas* (ed Montgomery, G. G.) 113-120 (Smithsonian Institution Press, 1985).
838

839 ⁶¹ MacPhee, R. D. E., Iturralde-Vinent, M. A. & Gaffney, E. S. Domo de Zaza: An Early
840 Miocene vertebrate locality in south-central Cuba, with notes on the tectonic evolution of Puerto
841 Rico and Mona Passage. *Amer. Mus. Novitates* **3394**, 1-42 (2003).
842 <http://hdl.handle.net/2246/2820>

843

844 ⁶² Tong, Y. F., Binford, G., Rheims, C. A., Kuntner, M., Liu, J., Agnarsson, I. Huntsmen of the
845 Caribbean: Multiple tests of the GAARlandia hypothesis. *Mol. Phylogenet. Evol.* **130**, 259-268
846 (2019). doi:10.1016/j.ympev.2018.09.017

847

848 ⁶³ Steadman, D. W., Martin, P. S., MacPhee, R. D. E., Jull, A. J. T., McDonald, H. G., Woods, C.
849 A., Iturralde-Vinent, M. A. & Hodgins, G. W. L. Asynchronous extinction of late Quaternary
850 sloths on continents and islands. *Proc. Nat. Acad. Sci. USA* **102** (33), 11763-11768 (2005). doi:
851 10.1073/pnas.0502777102

852

853 ⁶⁴ Hoorn, C., *et al.* Amazonia through time: Andean uplift, climate change, landscape evolution,
854 and biodiversity. *Science* **330** (6006), 927-931 (2010). doi: 10.1126/science.1194585

855

856 ⁶⁵ Tejada-Lara, J. V., *et al.* Life in proto-Amazonia: Middle Miocene mammals from the
857 Fitzcarrald Arch (Peruvian Amazonia). *Palaeontology* **58** (2), 341-378 (2015). doi:

858 10.1111/pala.12147

859

860 ⁶⁶ Penkman, K., Kaufman, D. S., Maddy, D. & Collins, M. J. Closed-system behaviour of the
861 intra-crystalline fraction of amino acids in mollusc shells. *Quat. Geochronol.* **3** (1-2), 2-25
862 (2008). doi: 10.1016/j.quageo.2007.07.001.

863

864 ⁶⁷ Kaufman, D. S. & Manley, W. F. A new procedure for determining DL amino acid ratios in
865 fossils using reverse phase liquid chromatography. *Quat. Sci. Rev.* **17** (11), 987-1000 (1998). doi:
866 10.1016/S0277-3791(97)00086-3

867

868 ⁶⁸ Demarchi, B., *et al.* Protein sequences bound to mineral surfaces persist into deep time. *eLife*
869 **5**, e17092 (2016). doi: 10.7554/eLife.17092

870

871 ⁶⁹ Kontopoulos, I., Presslee, S., Penkman, K. & Collins, M. J. Preparation of bone powder for
872 FTIR-ATR analysis: the particle size effect. *Vibrational Spectroscopy* **99**, 167–177 (2018). doi:
873 10.1016/j.vibspec.2018.09.004

874

875 ⁷⁰ Van Doorn, N. L., Hollund, H. & Collins, M. J. A novel and non-destructive approach for
876 ZooMS analysis: Ammonium bicarbonate buffer extraction. *Archaeol. Anthropol. Sci.* **3** (3), 281-
877 289 (2011). doi: 10.1007/s12520-011-0067-y

878

879 ⁷¹ Chambers, M. C., *et al.* A cross-platform toolkit for mass spectrometry and proteomics. *Nature*
880 *Biotechnol.* **30**, 918-920 (2012). doi: 10.1038/nbt.2377

881

882 ⁷² Kearse, M., *et al.* Geneious Basic: an integrated and extendable desktop software platform for
883 the organization and analysis of sequence data. *Bioinformatics* **28** (12), 1647-1649 (2012). doi:
884 10.1093/bioinformatics/bts199

885

886 ⁷³ Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high throughput.
887 *Nucl. Acids Res.* **32** (5), 1792-1797 (2004). doi: 10.1093/nar/gkh340

888

889 ⁷⁴ Swofford, D. L. PAUP* *Phylogenetic Analysis Using Parsimony (and Other Methods)*.
890 Version 4 (Sinauer Associates, 2002). doi: 10.1002/0471650129.dob0522

891

892 ⁷⁵ Lanfear, R., Calcott, B., Ho, S. Y. W. & Guindon, S. PartitionFinder: combined selection of
893 partitioning schemes and substitution models for phylogenetic analyses. *Mol. Biol. Evol.* **29** (6),
894 1695-1701 (2012). doi: 10.1093/molbev/mss020

895

896 ⁷⁶ Lanfear, R., Frandsen, P. B., Wright, A. M., Senfeld, T. & Calcott B. PartitionFinder 2: New
897 methods for selecting partitioned models of evolution for molecular and morphological
898 phylogenetic analyses. *Mol. Biol. Evol.* **34** (3), 772-773 (2017). doi: 10.1093/molbev/msw260

899

900 ⁷⁷ Dayhoff, M. O., Schwartz, R. M. & Orcutt, B. C. in *Atlas of Protein Sequence and Structure* **5**
901 (ed Dayhoff, M. O.) 345-352 (National Biomedical Research Foundation, 1978).

902

903 ⁷⁸ Ronquist, F., Teslenko, M., Van Der Mark, P., Ayres, D.L., Darling, A., Höhna, S., Larget, B.,
904 Liu, L., Suchard, M.A. & Huelsenbeck, J.P. MrBayes 3.2: Efficient Bayesian phylogenetic

905 inference and model choice across a large model space. *Sys. Biol.* **61** (3), 539-542 (2012). doi:
906 10.1093/sysbio/sys029
907
908 ⁷⁹ Kass, R.E. & Raftery, A. E. Bayes Factors. *J. Amer. Stat. Assoc.* **90** (430), 773-795 (1995).
909
910 ⁸⁰ Heath, T. A, Huelsenbeck, J. P. & Stadler, T. The fossilized birth-death process for coherent
911 calibration of divergence-time estimates. *Proc. Nat. Acad. Sci. USA.* **111** (29), 2957-2966 (2014).
912 doi: 10.1073/pnas.1319091111
913
914 ⁸¹ Gavryushkina, A., Welch, D., Stadler, T. & Drummond, A. J. Bayesian inference of sampled
915 ancestor trees for epidemiology and fossil calibration. *PLoS Comput. Biol.* **10** (12), e1003919
916 (2014). doi: 10.1371/journal.pcbi.1003919
917
918 ⁸² Gavryushkina, A., Heath, T. A., Ksepka, D. T., Stadler, T., Welch, D. & Drummond, A. J.
919 Bayesian total-evidence dating reveals the recent crown radiation of penguins. *Syst. Biol.* **66** (1),
920 57-73 (2017). doi: 10.1093/sysbio/syw060
921
922 ⁸³ Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C.-H., Xie, D., Suchard, M. A.,
923 Rambaut, A. & Drummond, A. J. BEAST 2: a software platform for Bayesian evolutionary
924 analysis. *PLoS Comput. Biol.* **10** (4) e1003537 (2014). doi: 10.1371/journal.pcbi.1003537
925
926 ⁸⁴ De Iuliis, G., Gaudin, T. J. & Vicars, M. J. A new genus and species of nothrotheriid sloth
927 (*Xenarthra*, Tardigrada, Nothrotheriidae) from the Late Miocene (Huayquerian) of Peru.
928 *Palaeontology* **54**, 171-205 (2011). doi: 10.1111/j.1475-4983.2010.01001.x

929

930 ⁸⁵ De Iuliis, G., Pujos, F., Toledo, N., Bargo, M. S. & Vizcaíno, S. F. *Eucholoeops* Ameghino,
931 1887 (Xenarthra, Tardigrada, Megalonychidae) from the Santa Cruz Formation, Argentine
932 Patagonia: implications for the systematics of Santacrucian sloths. *Geodiversitas* **36** (2), 209-255
933 (2014). doi: 10.5252/g2014n2a2

934 ⁸⁶ Hirschfeld, S. E. & Webb, S. D. Plio-Pleistocene megalonychid sloths of North America. *Bull.*
935 *Florida Mus. Nat. Hist. Biol. Sci.* **12** (5), 213-294 (1968).

937 ⁸⁷ Delsuc, F. *et al.* Ancient mitogenomics rewrites the evolutionary history and biogeography of
938 sloths. *Current Biology*. [tk]

939

940 **FIGURE AND TABLE LEGENDS (MAIN TEXT)**
941

942 **Figure 1:** Phylogenetic relationships among major folivoran taxa based on morphological
943 evidence (mostly after ref. 8, 16), with existence of unallocated taxa acknowledged. In this
944 framework, the three-toed tree sloth *Bradypus* is sister to other sloths (grouped here as
945 Eutardigrada), while the two-toed tree sloth *Choloepus* is included within Megalonychidae.

946

947 **Figure 2:** Geographical locations of sequenced samples. Sequences for *Cyclopes* and *Lestodon*
948 (**in bold**) taken from the literature; others, this paper (Table 1 and Supplementary Information,
949 Fig. S1).

950

951 **Figure 3:** 50% majority rule consensus tree from Bayesian analysis of the proteomic data
952 without temporal information, as performed in MrBayes. Values below nodes are posterior
953 probabilities for the descendant clade (see Results). Values above nodes are bootstrap support
954 derived from 10,000 bootstrap replicates. A dash (–) indicates that a node was not represented in
955 the 50% majority rule bootstrap consensus. Extant *Dasypus* and extinct *Doedicurus* and
956 *Glyptodon* are members of the order Cingulata; extant *Cyclopes* is a representative of
957 Vermilingua, which together with Folivora comprise order Pilosa. Cingulates and pilosans
958 together comprise superorder Xenarthra (see also Fig. 4).

959

960 **Figure 4:** Time scaled maximum clade credibility tree from BEAST analysis of 24 extant and
961 extinct xenarthran collagen sequences plus published mitochondrial genomes (see text). Branch
962 lengths are the mean values from the retained posterior sample, while blue bars represent 95%
963 highest posterior density intervals. Values at nodes are posterior probabilities (note that the
964 monophyly of *Bradypus* is constrained here). Vertical shaded bars correspond to South American
965 land mammal ages (SALMAs), two of which are emphasized: Deseadan (**), 29–21 Ma, during
966 which the first generally-accepted representatives of traditional Megatherioidea and
967 Mylodontoidea appear paleontologically; and the Santacrucian (*), 17.5–16.3 Ma, the SALMA
968 during which mylodontids maintained substantial taxonomic diversity but megalonychids and
969 megatheriids declined⁹. On the right (grey boxes), folivoran species used in analyses are
970 associated with their traditional family names, but with superfamily contents organized
971 according to phylogenetic conclusions in text. Megalocnidae is placed outside traditional
972 superfamily structure in its own (unnamed) box. The tree implies that the fundamental split
973 within Folivora is not between Megatherioidea and Mylodontoidea vs. Bradypodoidea as

974 classically understood, but instead between redefined *Megatherioidea and *Mylodontoidea vs.
975 Megalocnidae.

976

977 **Table 1:** Collagen peptides and per cent coverage of the sequenced ancient and modern samples.

978

979 **Table 2:** Selected divergence time estimates from BEAST analyses using different combinations
980 of taxa and data (see Results and Supplementary Information). Note that, although consistently

981 recovered as monophyletic, the position of Megalocnidae shifted among analyses, falling

982 alternately as sister to all other Folivora (Xenarthra) or Megatherioidea (Folivora).

983

Fig. 1





Fig. 2






Fig. 3

Fig. 4




Table 1 Collagen peptides and per cent coverage

Museum reference ¹	ID	Species	# Collagen Peptides	% Coverage
MMP 5672	15191	<i>Doedicurus sp.</i>	867	90
MACN-PV 7	15194	<i>Glyptodon sp.</i>	731	84
UF 76796	15559	<i>Acratocnus ye</i>	696	86
UF 76385	15565	<i>Acratocnus ye</i>	629	87
AMNH 20820	16265	<i>Bradypus variegatus</i>	793	88
AMNH 139772	17009	<i>Choloepus hoffmanni</i>	1109	94
MACN-PV 2652	15216	<i>Glossotherium robustum</i>	837	88
UF 169931	15564	<i>Megalocnus zile</i> ²	6	6
NYSM VP-46	16849	<i>Megalonyx jeffersonii</i> ³	874	85
MAPBAR 3965	15225	<i>Megatherium americanum</i>	520	81
UMAG ah 5854	16222	<i>Mylodon darwini</i>	1371	96
UF 171347	15548	<i>Neocnus comes</i>	699	84
UF 170210	15780	<i>Neocnus comes</i>	591	84
UF 75469	15781	<i>Neocnus dousman</i>	614	74
USNM 244372	14723	<i>Nothrotheriops shastensis</i>	528	79
USNM 3000	14715	<i>Paramylodon harlani</i>	642	87
UF 75526	15556	<i>Parocnus serus</i>	575	82
MUSM 1386	17480	<i>Scelidodon sp.</i>	1324	92
MACN-PV 1791	15202	<i>Scelidothierium sp.</i>	475	76

¹Institutional acronyms:

AMNH-M, American Museum of Natural History (Mammalogy), New York, USA

MACN-PV, Museo Argentino de Ciencias Naturales "Bernardino Rivadavia", Buenos Aires, Argentina

[Document title]

MAPBAR, Museo de la Asociación Paleontológica Bariloche, Bariloche, Argentina

MMP, Museo Municipal de Ciencias Naturales "Lorenzo Scaglia" Mar del Plata, Buenos Aires, Argentina

MUSM, Museo de Historia Natural, Universidad Nacional Mayor de San Marcos, Lima, Peru

NYSM VP, New York State Museum (Vertebrate Paleontology), Albany, New York, USA

UF, University of Florida, Natural History Museum of Florida, Gainesville, USA

UMAG ah, Instituto de La Patagonia, Universidad de Magallanes, Punta Arenas, Chile

USNM, United States National Museum of Natural History (Paleobiology), Washington DC, USA

²Mainly contaminants; not sequenced.

³SDS/PAGE protein extraction

Table 2: Selected divergence time estimates from BEAST analyses using different combinations of taxa and data (see Results and Supplementary Information). Note that, though consistently recovered as monophyletic, the position of Megalocnidae shifted among analyses, falling alternately as sister to all other Folivora (Xenarthra) or Megatherioidea (Folivora).

Clade	Protein only		mtDNA + Protein
	Xenarthra	Folivora	Xenarthra
Crown Xenarthra	62.0 (57.6 - 62.8)	-	62.6 (58.0 - 70.2)
Pilosa	50.4 (37.4 - 62.8)	-	52.1 (35.8 - 64.8)
Folivora	26.4 (18.0 - 36.0)	23.4 (14.9 - 33.9)	31.2 (21.1-41.4)
Megalocnidae	9.9 (3.8 - 17.8)	7.7 (3.4 - 13.0)	12.7 (4.4-22.6)
Megatherioidea + Megalocnidae	-	19.4 (12.8 - 27.8)	-
Megatherioidea + Mylodontoidea	22.7 (16.1 - 31.0)	-	26.9 (17.2 - 34.4)
Megatherioidea	15.7 (10.7 - 21.8)	13.9 (9.4 - 19.4)	23.0 (14.0 - 30.1)
<i>Megalonyx + Bradypus</i>	11.1 (8.4 - 15.0)	10.5 (8.4 - 14.1)	18.4 (8.4 - 25.2)
<i>Bradypus</i> spp.	-	-	16.0 (1.5 - 22.1)
<i>Megatherium + Nothrotheriops</i>	12.3 (8.4 17.7)	10.9 (7.8 - 15.1)	16.7 (9.9 - 23.6)
Mylodontoidea	15.3 (9.8 - 21.4)	15.4 (8.9 - 23.4)	22.2 (14.7 - 30.0)
<i>Choloepus + Mylodontidae</i>	12.03 (7.3 - 17.2)	10.5 (6.2 - 15.9)	20.5 (13.8 - 27.9)
<i>Choloepus</i> spp.	-	-	6.8 (2.6 - 11.8)