



Geodesi

2008-08-01

Gauss Conformal Projection (Transverse Mercator)

Krüger's Formulas

Symbols and Definitions

a	semi-major axis of the ellipsoid
f	flattening of the ellipsoid
e^2	first eccentricity squared
φ	geodetic latitude, positive north
λ	geodetic longitude, positive east
x	grid coordinate, positive north
y	grid coordinate, positive east
λ_0	longitude of the central meridian
k_0	scale factor along the central meridian
$\delta\lambda$	difference $\lambda - \lambda_0$
FN	false northing
FE	false easting

All angles are expressed in radians. Please note that the x-axis is directed to the north and the y-axis to the east.

The following variables are defined out of the ellipsoidal parameters a and f :

$$e^2 = f(2 - f)$$

$$n = \frac{f}{(2 - f)}$$

$$\hat{a} = \frac{a}{(1 + n)} \left(1 + \frac{1}{4} n^2 + \frac{1}{64} n^4 + \dots \right)$$

Conversion from geodetic coordinates (φ, λ) to grid coordinates (x, y) .

Compute the conformal¹ latitude φ^*

$$\varphi^* = \varphi - \sin \varphi \cos \varphi (A + B \sin^2 \varphi + C \sin^4 \varphi + D \sin^6 \varphi + \dots)$$

The coefficients A, B, C , and D are computed using the following formulas:

$$A = e^2$$

$$B = \frac{1}{6} (5e^4 - e^6)$$

$$C = \frac{1}{120} (104e^6 - 45e^8 + \dots)$$

$$D = \frac{1}{1260} (1237e^8 + \dots)$$

Let $\delta\lambda = \lambda - \lambda_0$ and

$$\xi' = \arctan(\tan \varphi^* / \cos \delta\lambda)$$

$$\eta' = \operatorname{arctanh}(\cos \varphi^* \sin \delta\lambda)$$

then

$$x = k_0 \hat{a} \left(\begin{array}{l} \xi' + \beta_1 \sin 2\xi' \cosh 2\eta' + \beta_2 \sin 4\xi' \cosh 4\eta' + \beta_3 \sin 6\xi' \cosh 6\eta' + \\ + \beta_4 \sin 8\xi' \cosh 8\eta' + \dots \end{array} \right) + FN$$

$$y = k_0 \hat{a} \left(\begin{array}{l} \eta' + \beta_1 \cos 2\xi' \sinh 2\eta' + \beta_2 \cos 4\xi' \sinh 4\eta' + \beta_3 \cos 6\xi' \sinh 6\eta' + \\ + \beta_4 \cos 8\xi' \sinh 8\eta' + \dots \end{array} \right) + FE$$

¹ Older Swedish literature refers to this quantity as the isometric latitude. Today the term isometric latitude is applied to the quantity

$\psi = \ln \{ \tan(\pi/4 + \varphi/2) [(1 - e \sin \varphi)/(1 + e \sin \varphi)]^{e/2} \}$. The isometric latitude is related to the conformal latitude by $\psi = \ln \tan(\pi/4 + \varphi^*/2)$. Cf. John P. Snyder: Map Projections - A Working Manual, U.S. Geological Survey Professional Paper 1395.

where the coefficients $\beta_1, \beta_2, \beta_3$ and β_4 are computed by

$$\beta_1 = \frac{1}{2}n - \frac{2}{3}n^2 + \frac{5}{16}n^3 + \frac{41}{180}n^4 + \dots$$

$$\beta_2 = \frac{13}{48}n^2 - \frac{3}{5}n^3 + \frac{557}{1440}n^4 + \dots$$

$$\beta_3 = \frac{61}{240}n^3 - \frac{103}{140}n^4 + \dots$$

$$\beta_4 = \frac{49561}{161280}n^4 + \dots$$

Conversion from grid coordinates (x,y) to geodetic coordinates (φ,λ)

Introduce the variables ξ and η as

$$\xi = \frac{x - FN}{k_0 \cdot \hat{a}}$$

$$\eta = \frac{y - FE}{k_0 \cdot \hat{a}}$$

Let

$$\begin{aligned}\xi' &= \xi - \delta_1 \sin 2\xi \cosh 2\eta - \delta_2 \sin 4\xi \cosh 4\eta - \delta_3 \sin 6\xi \cosh 6\eta \\ &\quad - \delta_4 \sin 8\xi \cosh 8\eta - \dots\end{aligned}$$

$$\begin{aligned}\eta' &= \eta - \delta_1 \cos 2\xi \sinh 2\eta - \delta_2 \cos 4\xi \sinh 4\eta - \delta_3 \cos 6\xi \sinh 6\eta \\ &\quad - \delta_4 \cos 8\xi \sinh 8\eta - \dots\end{aligned}$$

where

$$\delta_1 = \frac{1}{2}n - \frac{2}{3}n^2 + \frac{37}{96}n^3 - \frac{1}{360}n^4 + \dots$$

$$\delta_2 = \frac{1}{48}n^2 + \frac{1}{15}n^3 - \frac{437}{1440}n^4 + \dots$$

$$\delta_3 = \frac{17}{480}n^3 - \frac{37}{840}n^4 + \dots$$

$$\delta_4 = \frac{4397}{161280}n^4 + \dots$$

The conformal latitude φ^* and the difference in longitude $\delta\lambda$ are obtained by the formulas

$$\varphi^* = \arcsin(\sin \xi' / \cosh \eta')$$

$$\delta\lambda = \arctan(\sinh \eta' / \cos \xi')$$

Finally, the latitude φ and the longitude λ are obtained by the formulas

$$\lambda = \lambda_0 + \delta\lambda$$

$$\varphi = \varphi^* + \sin \varphi^* \cos \varphi^* (A^* + B^* \sin^2 \varphi^* + C^* \sin^4 \varphi^* + D^* \sin^6 \varphi^* + \dots)$$

where

$$A^* = (e^2 + e^4 + e^6 + e^8 + \dots)$$

$$B^* = -\frac{1}{6}(7e^4 + 17e^6 + 30e^8 + \dots)$$

$$C^* = \frac{1}{120}(224e^6 + 889e^8 + \dots)$$

$$D^* = -\frac{1}{1260}(4279e^8 + \dots)$$

Worked example:

ELLIPSOID GRS 1980

Semi-major axis (a)	6378137.0000 m.
Flattening (f)	1/298.257222101

TRANSVERSE MERCATOR PARAMETERS

Longitude of the central meridian	13 35 7.692000 degr. min. sec.
Scale factor on the central meridian	1.000002540000
False northing	-6226307.8640 m.
False easting	84182.8790 m.

Latitude and longitude	66 0 0.0000	24 0 0.0000 degr. min. sec.
Grid coordinates	1135809.413803	555304.016555 m